

Lichens of neglected habitats in Eastern and East-Central European lowlands

JURGA MOTIEJŪNAITĖ

Laboratory of Mycology, Institute of Botany, Žaliųjų ežerų str. 49,
LT 08406 Vilnius 21, mikojm@botanika.lt

Motiejūnaitė J.: *Lichens of neglected habitats in Eastern and East Central European lowlands*. Acta Mycol. 41 (1): 145-154, 2006.

Situation of lichens of aquatic and transient habitats in Eastern and East-Central European lowlands is discussed basing on example of several selected species: *Leptogium biatorinum*, *Sarcosagium campestre*, *Steinia geophana*, *Verrucaria aquatilis*, *V. hydrela*, *V. praetermissa*, *V. xyloxena*. Both habitat types are generally very much neglected in the region and all species show large spatial gaps in recording, which makes it difficult to judge both about their true distribution limits and spreading dynamics. On the other hand, targeted search through the suitable habitats and abundance of such indicate that many of these lichens are probably not uncommon in the region.

Key words: Lichens, biogeography, aquatic habitats, transient habitats

INTRODUCTION

Rare lichens often make significant parts of national floras and provide the basis for red lists. However, “rare” might comprise both declining species and stable or spreading ones. Declining species are usually well known, more or less historically documented and confined to the habitats and substrates that are traditionally investigated by lichenologists. Meanwhile situation of many species defined as “rare” in national floras often remain obscure. Especially this is true for Eastern and East-Central (EEC) Europe. The reasons for some lichen to be defined as rare can be several:

1. The species are “new” – relatively recently described.
2. The species are taxonomically “difficult”, they are often reported as complexes or not separated from similar species.
3. The species are connected to particular habitats or niches that are neglected by lichenologists.

“New”, recently described species comprise lichens, which have been described during the last two decades. Presence of such species in the national lists generally indicates degree of recent lichenological activity (see e.g. Motiejūnaitė 2005). Taxonomically „difficult“ species are either scarce or totally excluded from the floras or inventory lists of eastern Europe (e.g., most of sterile crustose lichens) or poorly distinguished, e.g. see notes for *Acrocordia cavata* in Motiejūnaitė (2005) or *Anisomeridium polypori* in Czyżewska et al. (2005). Species associated with habitats neglected by lichenologists make a weird group, indicating, to certain extent, activities of lichenologists, the variability of fieldwork and nature of lichenological coverage. Species of this group are often mentioned in discussions about lichen flora dynamics, as most of them are considered to be rare in many European countries; a sudden boost in locality numbers after specific habitat studies might result in a discussion as to whether the species is spreading or not, whether the species is a neophyte or a long-present member of the national flora.

Typical examples of neglected habitats are lowland freshwater bodies and transient habitats, both of which receive only sporadic attention from lichenologists. The purpose of this paper was to review situation of lichens associated with both these habitats in EEC Europe and to attract attention to the importance of study of these habitats. For the aims of the paper several species should be characteristic to aquatic and transient habitats were selected. Their known distribution and situation in EEC Europe were analysed mainly on the basis of all available literature sources and to some extent of herbarium collections (mainly Lithuanian).

TERRITORY UNDER DISCUSSION

For the review temperate and cool temperate forest zone of lowland EEC Europe (between ca. 20° and 55° longitude and 53° and 63° latitude) was chosen. Though the area is biogeographically complex, it still holds certain homogeneity pertaining conditions for lichen distribution. There are no large natural obstacles for lichen dispersal in this area, the only limiting factors for their spread being climatic and edaphic. Though land-use history and intensity vary from country to country, there are still many similarities connected with long periods of similar administrative history. Many traditional rural activities persist, such as use of local building materials resulting in sand-, gravel- and loam-pits, and of untreated timber for buildings (although the latter tendency is decreasing lately at least in part of the territory).

Administratively the territory includes Polish lowlands (P), Baltic countries – Lithuania (Li), Latvia (La), Estonia (E), Belarus, and the forest zones of the Ukrainian lowlands (U) and of European Russia (R).

RESULTS AND DISCUSSION

Lichens of aquatic habitats. Aquatic habitats, though locally well studied in Western and Central Europe (see e.g. Gilbert 1996; Gilbert, Giavarini 1997; Molitor, Diederich 1997; Keller 2000; Thüs 2002) are very much under-recorded in EEC Europe, especially in its lowland parts. Both numbers of recorded species are low and the records themselves are sparse, though when well studied, aquatic environs show to be rich in lichens. Crustose lichens are espe-

cially understudied, among which perithecioid species prevail. *Verrucaria aquatilis*, *V. hydrela* and *V. praetermissa* are the most common freshwater aquatic perithecioid lichens in the region, their distribution and status reflects situation of whole group.

Distribution of these lichens in EEC Europe reflects solely regional activity of lichenologists and gives no clear pattern of geographical ranges. Most of them (except *V. hydrela*) are attributed to suboceanic species, occurring mainly in temperate zone. Generally, these lichens are quite well known in Western and Central Europe, but are very much under-recorded in the east. Aquatic *Verrucaria* species are found almost exclusively in southeastern and eastern region of the Baltic sea (Figs. 1-3). Though there is no shortage of suitable habitats in Eastern Europe, aquatic lichens, especially crustose species are almost utterly omitted from lichen inventories. Judging by the known distribution and tendencies of recording, aquatic *Verrucaria* species should be found also further eastwards in pristine streams of forested areas, as it is demonstrated by a record of *V. hydrela* outside of the discussed area, in southern part of the Baikal lake region (Urbanavichene, Urbanavichus 1999). Besides, these lichens are source of taxonomic confusion, which is discernible even among scarce records (e.g., the record of *V. praetermissa* in Kaliningrad region, Russia (Andreev 2002) suggests *V. elaeina* (see Orange 2000).

Frequencies of these lichens in the discussed region are as hypothetical, as their distribution ranges. In most cases they range from very rare to rather rare in all countries (Tab. 1), though it is admitted that overlooking is the main reason for the record scarcity (e.g. for *Verrucaria hydrela* in Northern Poland (Jando, Kukwa 2003).

Dynamics of these species as well as conservational value are even less clear. Generally, aquatic lichens are connected with vulnerable habitats, as most of them inhabit pristine streams, little influenced by eutrophication (see Gilbert 1996; Gilbert, Giavarini 1997; Thüs 2002; Motiejūnaitė 2003). This is well expressed by the fact that aquatic *Verrucaria* species are red-listed in West-European areas of intensive urban and agricultural land use: *V. aquatilis* and *V. hydrela* – in Denmark (Søchting 1997) and lowland lands in Germany (variable categories, Wirth et al. 1996); *V. praetermissa* – in German lowland lands (Critically endangered to Rare). Thus, they could be attributed to habitat-demanding and decreasing species. In EEC Europe, aquatic *Verrucaria* species are red-listed only in Poland (Tab. 1).




Fig. 1. Distribution of *Verrucaria aquatilis* in Eastern and East Central Europe

- records more than 50 year old,
- modern records




Fig. 2. Distribution of *Verrucaria hydrela* in Eastern and East Central Europe

- records more than 50 year old,
- modern records




Fig. 3. Distribution of *Verrucaria praetermissa* in Eastern and East Central Europe
● modern records

Lichens of transient habitats. Lichens of transient habitats are ephemeral, generally inconspicuous and rarely recorded, therefore are presumed to be nationally rare in almost all European countries. Most often such habitats are man-made, like bare soil of roadside scarps, edges of dirtroads, gravel-, sand- or loam-pits. Such places bear diverse albeit very much understudied lichen flora (see e.g. Ernst 1993, 1995). *Leptogium biatorinum*, *Sarcosagium campestre*, *Steinia geophana* and *Verrucaria xyloxytina* represent typical members of pioneer lichen communities inhabiting transient man-made habitats in EEC European lowlands. All these species are probably widely distributed both regarding climatic zones and oceanic gradient.

Distribution of these lichens in EEC Europe even more strongly reflects regional activity of lichenologists, as these species are most often recorded during targeted studies of specific habitats. To date, the widest distribution in west-east direction is demonstrated by *Sarcosagium campestre* (Fig. 4), *Verrucaria xyloxytina* (Fig. 5), and *Leptogium biatorinum* (Fig. 6), two latter species being recorded also outside the discussed region of EEC Europe, in the steppe zone in Russia (Vedeneev 1999; Muchnik 2001; Shustov 2002). *Steinia geophana* is more widely distributed in a south-north direction (Fig. 7): outside the discussed area its range reaches Kola Peninsula in Russia (BILAS herbarium, specimens No. 5304 and 5339). However, not all of this data is totally reliable, as at least two species can be source of taxonomic confusion. In some references (Western European as well) *L. biatorinum* is not differentiated from *L. byssinum* (Jørgensen 1994). In earlier Russian and Ukrainian references (Oksner 1956; Inashvili 1975), the description of *L. byssinum*




Fig. 4. Distribution of *Sarcosagium campestre* in Eastern and East Central Europe
● modern records




Fig. 5. Distribution of *Verrucaria xyloxytina* in Eastern and East Central Europe
○ records more than 50 year old,
● modern records

Table 1

Distribution, frequency and conservation status of the selected lichen species in Eastern and East Central European lowlands

Frequency abbreviations: vr (very rare), r (rare), rr (rather rare), rf (rather frequent), f (frequent), vf (very frequent). Frequency evaluations for Lithuania are given according to Motiejūnaitė (2002a), for Estonia according to Rändlane and Saag (1999), for Latvia according to Piterāns (2002) (in part). In some cases parts of countries are indicated (e.g., NE north eastern, C central). **Conservation status** abbreviations: VU (Vulnerable), NT (Near threatened), DD (Data deficient). References for conservation information: Cieśliński (2003b), Cieśliński et al. (2003). For country abbreviations see chapter "Territory under discussion".

Species	Countries, frequency	References	Conservation status
<i>Leptogium biatorinum</i> (Nyl.) Leight.	P (rr), Li (rr), Lv (vr), R (vr)	Lettau 1912; Ceynowa Gieldon 2001; Cieśliński 2003a; Motiejūnaitė et al. 2006	
<i>Sarcosagium campestre</i> (Fr.) Poetsch & Schied.	P (rr), Li (rr), Lv (vr), U(vr), E(vr)	Liśnicki 1991; Fałtynowicz 1992, 2003; Kondratyuk et al. 1998; Ceynowa Gieldon 1999; Rutkowski, Słowiak 1999; Cieśliński 2003a; Sparrius 2003; Suija et al. 2005; Motiejūnaitė et al. 2006; Kukwa, pers. com.	
<i>Steinia geophana</i> (Nyl.) Stein	P (r), Li (rr), E (r), R (vr)	Fałtynowicz 1993; 2003; Fadeeva et al. 1997; Ceynowa Gieldon 1998, 1999; Zavarzin et al. 1999; Halonen et al. 2000; Kuznetsova, Himelbrant 2002; Cieśliński 2003a; Aptroot et al. 2005	P (NT in NE)
<i>Verrucaria aquatilis</i> Mudd	P (vr), Li (rr), La (vr), R (vr)	Kopachevskaya 1977; Fałtynowicz 1992; Rutkowski 1993; Fadeeva et al. 1997; Fałtynowicz et al. 2000; Kowalewska et al. 2000; Kukwa 2000; Cieśliński 2003a; Motiejūnaitė et al. 2006	P (VU, DD in NE)
<i>Verrucaria hydrella</i> Ach.	P (rr), Li (f), La (r), E (vr), R (vr)	Lettau 1912; Piterāns 1982; Fadeeva et al. 1997; Fałtynowicz et al. 2000; Halonen et al. 2000; Kowalewska et al. 2000; Kukwa 2000; Fałtynowicz, Królik 2001; Cieśliński 2003a; Jando, Kukwa 2003; Motiejūnaitė et al. 2006	P (VU, not red listed in NE)
<i>Verrucaria praetermissa</i> (Trevis.) Anzi	P (vr), Li (rf), La (vr), E (vr), R (vr)	Halonen et al. 2000; Andreev 2002; Cieśliński 2003a; Motiejūnaitė et al. 2006	P (NT, DD in NE)
<i>Verrucaria xyloxena</i> Norman	P (rf, common in C), Li (rr), E (vr), R (vr)	Orange 1991; Ceynowa Gieldon 1998; Rändlane, Saag 1999; Cieśliński 2003a; Kuznetsova, pers. com.	




Fig. 6. Distribution of *Leptogium biatorinum* in Eastern and East Central Europe
 ○ records more than 50 year old,
 ● modern records




Fig. 7. Distribution of *Steinia geophana* in Eastern and East Central Europe
 ○ records more than 50 year old,
 ● modern records

can accomodate both species. In Estonia only *L. byssinum* is recorded (Randlane, Saag 1999) basing only on literature data, thus the identity of the species is questionable. Illustrations and descriptions in Polish references (Nowak, Tobolewski 1975; Ceynowa-Giełdon 1999) define both lichens very well. Lettau (1912) also circumscribed these two species. Part of the reports of *V. xyloxena* southeastwards of the discussed area (Voronezh, Volgograd, Samara, Saratov, Uljanovsk regions of Russia) might be doubtful – the species is reported as saxicolous (Muchnik 2001; Shustov 2002).

Frequencies of transient habitat species in Eastern Europe reflect general recording – locally they are reported as quite common, but data is totally absent from most of the territory. In most countries they are reported to be rare or very rare (Tab. 1). Regardless record scarcity, it can be assumed that they are much more frequent, as traditionally there are numerous habitats suitable for these lichens, such as sand-, gravel- and loam-pits which are present in almost every village, dirt-roads, road scarps, etc. and recording does not show any tendencies of decrease. Even a cursory search through such habitats can result in a variety of finds (e.g. Motiejūnaitė 2002b; Sparrius 2003), and intensive targeted search can totally change frequency patterns (Ceynowa-Giełdon 1998, 1999, 2001). There are different opinions concerning the dynamics of these species in Europe. In Jørgensen (1994) and Jørgensen and Motiejūnaitė (2005) species of transitional habitats are presumed to be decreasing due to modern agricultural methods. Woods and Coppins (2003) express opinion that species of transient habitats should be excluded from the red-lists, though they evaluate *Verrucaria xyloxena* as Critically Endangered in the same reference. However, in some countries lichens of transient habitats are given threat status: *Leptogium biatorinum* is red listed in Denmark, Finland (Vitikainen et al. 1997) and in lowland lands of Germany (categories range from Extinct to Poorly known); *Sarcosagium campestre* and *Steinia geophana* – in Denmark and in lowland lands of Germany (variable categories); *Verrucaria xyloxena* is red-listed in Germany as Data deficient (not recorded in lowland lands). In EEC Europe, lichens of transient habitats can hardly be qualified as threatened though *Steinia geophana* is red listed in NE Poland. On the contrary, part of these lichens might belong to spreading species, as they occupy niches that are naturally scarce in the discussed region and appear mainly in connection with human activities.

CONCLUSIONS

As the examples in this paper have shown, there are large spatial gaps in recording lichens of aquatic and transitional habitats in EEC Europe, making these lichens nationally rare in almost all countries in the discussed region. Also there exists certain degree of taxonomic confusion pertaining these species.

Lichen floras, resulting from species inventories make base for environmental evaluation and monitoring programmes, land management policy making, etc. Disregard of certain habitats or species, or on the contrary, detailed study of all habitat types and all species present might lead to very different results of lichen inventories (see e.g. Lõhmus et al. 2003) and subsequently, to erroneous interpretations of monitoring evaluation results. In wider scale, both historical and present understudy of some habitats leads to the dearth of knowledge on true lichen distribution areas and tendencies of their decrease or spreading and often disable biodiversity monitoring projects in larger geographical scale.

Acknowledgements: I would like to express my sincere thanks to my colleagues prof. K. Czyżewska (Łódź), dr. M. Kukwa (Gdańsk), dr. V. Golubkov (Grodn), Mr. D. Himelbrant and Ms. E. Kuznetsova (St. Petersburg) for the data they supplied and allowed to use in the paper and their valuable comments on the manuscript.

REFERENCES

- Andreev M.P. 2002. Dopolnenije k likhenoflore Kaliningradskoi oblasti Rossii: lishainiki kamen istykh substratov. Novosti Sist. Nizsh. Rast. 36: 68–72.
- Aptroot A., Czarnota P., Jüriado I., Kocourková J., Kukwa M., Lõhmus P., Palice Z., Randlane T., Saag L., Sérusiaux E., Sipman H., Sparrius L. B., Suija A., Thüs H. 2005. New or interesting lichens and lichenicolous fungi found during the 5th IAL Symposium in Estonia. Folia Cryptog. Estonica 41: 13–22.
- Ceynowa Giełdon M. 1998. Nazemne gatunki porostów z rodzaju *Verrucaria* (Lichenes, Verrucariaceae) na Kujawach i niektórych terenach sąsiednich. Fragm. Flor. Geobot. Ser. Polonica 5: 243–249.
- Ceynowa Giełdon M. 1999. *Leptogium byssinum* (Lichenes, Collemataceae) na Kujawach i terenach sąsiednich nad dolną Wisłą. Fragm. Flor. Geobot. Ser. Polonica 6: 227–234.
- Ceynowa Giełdon M. 2001. Kalcyfilne porosty naziemne na Kujawach. Wyd. Uniwersytetu Mikołaja Kopernika, Toruń, 44 pp.
- Cieśliński S. 2003a. Atlas rozmieszczenia porostów (Lichenes) w Polsce Północno Wschodniej. Phytocoenosis 15 (N.S.), Suppl. Cartogr. Geobot. 15: 1–430.
- Cieśliński S. 2003b. Czerwona lista porostów zagrożonych w Polsce Północno Wschodniej. Monogr. Bot. 91: 91–106.
- Cieśliński S., Czyżewska K., Fabiszewski, J. 2003. Czerwona lista porostów wymarłych i zagrożonych w Polsce. Monogr. Bot. 91: 13–49.
- Czyżewska K., Motiejūnaitė J., Cieśliński S. 2005. New and noteworthy species of lichens and allied fungi from North Eastern Poland. Acta Mycol. 40 (2): 277–291.
- Ernst G. 1993. Zur Ökologie und Verbreitung von *Geisleria sycnogonioides*, einer bislang kaum bekannten terricolen Flechte. Herzogia 9: 321–337.
- Ernst G. 1995. *Vezdaea leprosa* Spezialist am Straßenrand. Herzogia 11: 175–188.
- Fadeeva M.A., Golubkova N.S., Vitikainen O., Ahti T. 1997. Predvaritel'nyj spisok lishajnikov Karel'ii i obitajushchikh na nikh gribov. Karel'skij nauchnyj centr RAN. Petrozavodsk.
- Fałtynowicz W. 1992. The lichens of Western Pomerania (NW Poland). An ecogeographical study. Polish Bot. Stud. 4: 1–182.

- Fałtynowicz W. 1993. *Steinia geophana* (Nyl.) B. Stein. (In:) S. Cieśliński, W. Fałtynowicz (eds.), *Atlas of the geographical distribution of lichens in Poland. Part 1. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków*, p. 65–67.
- Fałtynowicz W. 2003. The lichens, lichenicolous and allied fungi of Poland. An annotated checklist. *Biodiversity of Poland 6. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków*, 435 pp..
- Fałtynowicz W., Marcinkowska E., Rutkowski P. 2000. Porosty rezerwatu „Dolina Zagórskiej Strugi” koło Rumii na Pojezierzu Kaszubskim. *Acta Bot. Cassubica* 1: 119–126.
- Fałtynowicz W., Królak D. 2001: Porosty rezerwatu „Jar Rzeki Raduni” na Pojezierzu Kaszubskim (północna Polska). *Acta Bot. Cassubica* 2: 133–141.
- Gilbert O.L. 1996. The lichen vegetation of chalk and limestone streams in Britain. *Lichenologist* 28: 145–159.
- Gilbert O.L., Giavarini V.J. 1997. The lichen vegetation of acid watercourses in England. *Lichenologist* 29: 347–367.
- Halonen P., Kukwa M., Motiejunaite J., Löhmus P., Martin, L. 2000. Notes on lichens and lichenicolous fungi found during the XIV Symposium of Baltic Mycologists and Lichenologists in Järvselja, Estonia. *Folia Cryptog. Estonica* 36: 17–21.
- Inashvilli C.N. 1975. Sem. *Collemataceae*. (In:) O. B. Blum, A. V. Dombrovskaya, C. N. Inashvilli, A. V. Piterāns, E. G. Roms, V. P. Savic (eds), *Opredelitel lishainikov SSSR. 3. Caliciaceae Gyalectaceae*. Nauka, Leningrad: 80–118.
- Jando K., Kukwa M. 2003. Porosty, grzyby naporostowe i nażywicze projektowanego rezerwatu „Wiszące Torfowiska nad jeziorem Jaczno” oraz terenów przyległych do jeziora Jaczno w Suwalskim Parku Krajobrazowym (Północno Wschodnia Polska). *Parki nar. Rez. przyr.* 22 (1): 3–17.
- Jørgensen P.M. 1994. Further notes on European taxa of the lichen genus *Leptogium*, with emphasis on the small species. *Lichenologist* 26: 1–29.
- Jørgensen P.M., Motiejūnaitė J. 2005. *Lemmopsis pelodes* (Körb. ex Stein) Ellis found in Lithuania. *Graphis Scripta* 17 (1): 17–19.
- Keller C. 2000. Die Wasserflechten der Teigtisch zwischen der Landmannsperrre und dem kraftwerk Arnstein (Steiermark, Österreich). *Herzogia* 14: 49–58.
- Kondratyuk S.Ya., Khodosovtsev A.Ye., Zelenko S.D. 1998. The second checklist of lichen forming, lichenicolous and allied fungi of Ukraine. *Phytosociocentre*, Kiev.
- Kopachevskaya E.G. 1977. Sem. *Verrucariaceae*. In: Kopachevskaya E.G., Makarevich M.F., Oksner, A.N. *Opredelitel lishainikov SSSR. 4. Verrucariaceae Pilocarpaceae*: 7–54. Nauka, Leningrad.
- Kowalewska A., Jando K., Kukwa M. 2000. Lichens and lichenicolous fungi of the protected nature reserve “Przyjaźń”, protecting the forest support surrounded by the cultivated fields (Kaszu by lakes district, N. Poland). (In:) *Pervaja Rosijskaya Likhenologicheskaya shkola i mezhdunarodnyj simpozium molodykh likenologov “Arkoalpijskaja flora. Okhrana lishainikov”*. Programma i tezisy dokladov. Apatity: 75.
- Kukwa M. 2000: Porosty i grzyby naporostowe zachodniej części Pojezierza Iławskiego (Polska północna). *Fragmenta Florae Geobotanicae Ser. Polonica* 7: 281–297.
- Kuznetsova E.S., Himelbrant D. E. 2002. Dopolnenye k flore lishainikov Nizhesvirskogo zapovednika. *Novosti Sist. Nizhsh. Rast.* 36: 144–150.
- Lettau G. 1912. Beiträge zur Lichenenflora von Ost und Westpreussen. *Festschrift zum 50 jährigen bestehen des preussischen botanischen vereins Königsberg* 53: 17–91.
- Lipnicki L. (ed.). 1991. Porosty (Lichenes) Pszczewskiego Parku Krajobrazowego. (In:) V Meeting of Polish lichenologists: 5–32. Inst. Bad. Ekspertyzy Nauk., Gorzów Wlkp.
- Löhmus P., Saag L., Löhmus A. 2003. Is there merit in identifying leprarioid crusts to species in ecological studies? *Lichenologist* 35: 187–190.
- Molitor F., Diederich P. 1997. Les pyrénotriches aquatiques du Luxembourg et leurs champignons lichénicoles. *Bulletin de la Société Nationale luxembourgeoise* 98: 69–92.
- Motiejūnaitė J. 2002a. Lapiškosios ir krūmiškosios kerpės (Ascomycetes lichenisati. Species folioseae et fruticosae). *Lietuvos grybai* 13(1). Valstiečių laikraštis, Vilnius.
- Motiejūnaitė J. 2002b. Diversity of lichens and lichenicolous fungi in the transboundary region of Marijampolė district (southern Lithuania). *Botanica Lituanica* 8: 277–294.

- Motiejūnaitė J. 2003. Aquatic lichens in Lithuania. Lichens on submerged alder roots. Herzogia 16: 113–121.
- Motiejūnaitė J. 2005. Distribution of some rare and declining lichen species in lowland eastern and eastern central Europe. Biologia (Bratislava) 60 (4): 357–363
- Motiejūnaitė J., Piterāns A., Suija A., Lōhmuus P., Kuznetsova E., Tórra T., Pri godina Lukošienė I. 2006. New or noteworthy lichens, lichenicolous and allied fungi found during the 16th Symposium of Mycologists and Lichenologists in Latvia. Botanica Lithuanica (in prep.).
- Muchnik E.E. 2001. Konspekt lishainikov stepnykh i ostepnennykh mestoobitanij central'nogo Chernozemja. Novosti Sist. Nizhsh. Rast. 35: 183–195.
- Nowak J., Tbolewski Z. 1975. Porosty polskie. Państwowe Wydawnictwo Naukowe, Warszawa Kraków.
- Orange A. 1991. Notes on some terricolous species of *Verrucaria*. Lichenologist 23: 3–10.
- Orange A. 2000. *Verrucaria elaeina*, a misunderstood European lichen. Lichenologist 32: 411–422
- Oksner A.M. 1956. Flora lishainikiv Ukrainskoi. Tom I. Akademiya Nauk Ukrainskoi RSR, Institut Botaniki, Kiev.
- Piterāns A.V. 1982. Lishainiki Latvii. Zinatne, Riga.
- Piterāns A. 2002. Lichens of Latvia. <http://latvijas.daba.lv/scripts/db/saraksti/saraksti.cgi?d=keerpij&l=en>. [Last modified: 20.05.2002].
- Randlane T., Saag A. (eds) 1999. Second checklist of lichenized, lichenicolous and allied fungi of Estonia. Folia Cryptog. Estonica 35: 1–132.
- Rutkowski P. 1993. Flora i ekologia porostów rezerwatu „Jar Rzeki Reknicy” na Pojezierzu Kaszubskim. Parki nar. Rez. przyr. 12 (4): 29–40.
- Rutkowski P., Słowik J. 1999. Flora porostów doliny Drwęcy w okolicach Nowego Miasta Lubawskiego. Frgm. Flor. Geobot. Ser. Polonica 6: 211–217.
- Shustov M. V. 2002. Lishainiki Privilzhskoi vozyshennosti. Novosti Sist. Nizhsh. Rast. 36: 185–203.
- Søchting U. 1997. Laver. [In:] Rødliste 1997 over planter og dyr i Danmark. <http://www.sns.dk/net/pub/rodliste/72.htm>
- Sparrius L.B. 2003. Contribution to the lichen floras of the Białowieża Forest and the Biebrza Valley (Eastern Poland). Herzogia 16: 155–160.
- Suija A., Nomm M., Boch S. 2005. New Estonian records. Lichens and lichenicolous fungi. Folia Cryptog. Estonica 41: 135–136.
- Thüs H. 2002. Taxonomie, Verbreitung und Ökologie silicicoler Süßwasserflechten im außeralpinen Mitteleuropa. Bibl. Lichenol. 83: 1–214.
- Urbanavichene I.N., Urbanavichus G.P. 1999. K flore lishainikov khrebtov Chamar Daban (Juzhnoe pribajkalje). Novosti Sist. Nizhsh. Rast. 33: 161–171.
- Vedeneev A.M. 1999. Lishainiki Sarepty (Volgogradskaja oblastj). Bot. Zhurn. 84 (11): 100–107.
- Vitikainen O., Ahti T., Kuusinen M., Lommi S., Ulvinen T. 1997. Checklist of lichens and allied fungi of Finland. Norrlinia 6: 1–123.
- Wirth V., Schöller H., Scholtz P., Ernst G., Feuerer T., Gnüchtel A., Hauck M., Jacobsen P., John V., Litterski B. 1996. Rote Liste der Flechten (Lichenes) der Bundesrepublik Deutschland. Schriftenreihe für Vegetationskunde 28: 307–368.
- Woods R.G., Coppins B.J. 2003. A Conservation Evaluation of British Lichens. British Lichen Society, London.
- Zavarzin A.A., Katenina O.A., Kotlov Yu.V., Sokolova S.V. 1999. Lichens of St. Petersburg and Leningrad region. (In:) N.B. Balashova, A. A. Zavarzin (eds.), Biodiversity of the Leningrad region (Algae. Fungi. Lichens. Bryophytes. Invertebrates. Fishes and Pisciformes): 205–260. St. Petersburg University Press, St. Petersburg.

Porosty słabo poznanych siedlisk na niżu Europy Środkowo-Wschodniej i Wschodniej

Streszczenie

W pracy przedstawiono rozmieszczenie, frekwencję i stopień zbadania porostów siedlisk wodnych i krótkotrwałych w obrębie strefy leśnej niżu Europy Środkowo Wschodniej i Wschodniej: Polski, Litwy, Łotwy, Estonii i Białorusi oraz Ukrainy i europejskiej części Rosji. Oba typy siedlisk są bardzo słabo rozpoznane w tej części Europy. Zagadnienie to omówiono na przykładzie wybranych gatunków efemerycznych: *Leptogium biatorinum*, *Sarcosagium campestre*, *Steinia geophana* i *Verrucaria xylooxena* oraz wodnych: *Verrucaria aquatilis*, *V. hydrela* i *V. praetermissa*.

W skali wielkoprzestrzennej widoczne są luki w rozmieszczeniu badanych gatunków (Figs 1–7). Ustalenie ich rzeczywistego rozmieszczenia i dynamiki rozprzestrzenienia jest, jak na razie, trudne. Celowe poszukiwanie odpowiednich siedlisk i charakterystycznych im gatunków sygnalizuje, że mogą one być nawet częste w regionie.