

## ***Fusarium equiseti* as one of the main *Fusarium* species causing wilt and root rot of chickpeas in Morocco**

**El Hazzat N<sup>a</sup>, Adnani M<sup>a</sup>, Msairi S<sup>b</sup>, El Alaoui MA<sup>a,\*</sup>, Mouden N<sup>c</sup>, Chliyah M<sup>a</sup>, Boughribil S<sup>d</sup>, Selmaoui K<sup>a</sup>, Ouazzani Touhami A<sup>a</sup>, Douira A<sup>a</sup>**

<sup>a</sup>Laboratoire des Productions Végétales, Animales et Agro-industrie, Equipe de botanique, Biotechnologie et Protection des Plantes, Département de Biologie, Faculté des Sciences, Université Ibn Tofail, Kénitra (Morocco).

<sup>b</sup> Laboratory of Phytobiotechnology, National Agency of Medicinal and Aromatic Plants, Taounate, Morocco

<sup>c</sup> Laboratoire de Chimie Moléculaire et Molécules de l'Environnement, Faculté Pluridisciplinaire de Nador-Université Mohammed 1er Oujda, Maroc (Morocco).

<sup>d</sup>Laboratoire de Virologie Microbiologie, Qualité et Biotechnologie/ Ecotoxicology et Biodiversité, Faculté des Sciences et Techniques – Mohammedia, Hassan II Université de Casablanca (Morocco).

\*Corresponding author; e-mail: [manal.adnani@gmail.com](mailto:manal.adnani@gmail.com)

### **Abstract**

Fungal isolates of *Fusarium* were collected from symptomatic chickpea (*Cicer arietinum* L.) plants growing in fields within Souk Tlat commune in the Gharb region. Morphological and molecular characterizations were performed of the fungal isolate N3 obtained from a chickpea plant. PCR amplification and sequencing of the internal transcribed spacer using the primers ITS1 and ITS4 was applied to identify the fungal isolate N3. The maximum similarity index of the fungus was found to be 99.33% with *Fusarium equiseti* (accession no. MT111122). In the pathogenicity test, both chickpea seed dip inoculation and soil infestation by the spore suspension of *Fusarium* isolate were adopted. Four weeks after chickpea seed inoculation, few plants emerged and those that emerged were stunted. A high percentage of inoculated seeds did not emerge and

showed accentuated rot symptoms.. Eight weeks after sowing seeds in infested soil, the obtained chickpea seedlings displayed root necrosis, browning at the crown, and wilting. In addition, these plants showed a foliar alteration index of 0.395. The re-isolation was positive for different parts of chickpea plants for both seed and soil inoculation. *Fusarium equiseti* isolate decreased the length of the root and aerial parts, and number of leaves and branches of the inoculated chickpea plants either by seed inoculation or soil infestation with values of 0.91 cm and 19.73 cm, 1.29 cm and 19.44 cm, 1.11 and 18.66, and 0.0 and 2.08 respectively, whereas the corresponding values for the control plants were 27.16 and 28.33 cm, 29.05 and 31.05 cm, 24.21 and 25.66, and 3.50 and 3.11, respectively. To the best of our knowledge, this is the first report of *F. equiseti* on chickpea (*Cicer arietinum* L.) in Morocco.

**Keywords** – Chickpeas – inoculation – *Fusarium equiseti* – rot – wilting

Running head: *Fusarium equiseti* in chickpea: First report in Morocco

## 1. Introduction

Chickpea (*Cicer arietinum* L.) is one of the oldest cultivated protein legumes in the world. It is mainly used for human consumption and is an essential constituent of the Mediterranean diet and basic food in Pakistan and India (Millan et al., 2010). Domesticated in association with other crops such as wheat and barley, *Cicer arietinum* L. is believed to be a part of the agricultural revolution, and in terms of consumption, ranks second after broad bean (Gupta et al., 2014).

In Morocco, the cultivated chickpea area covers an acreage of 54000 ha (MAPMDREF, 2020a), with a production of 49700 T recorded in the 2019/2020 agricultural season (MAPMDREF, 2020b). Most of the agricultural systems of chickpea production in the country suffer from several constraints, mainly biotic and abiotic factors that cause serious damage before and after harvest. The widespread high temperature and drought stress in different regions of chickpea production

can affect flowering and pod setting stages which lead to decreases in chickpea yield (Houasli et al., 2020). Previous reports on the chickpea crop have recorded the presence of *Ascochyta rabiei*, the causal agent of blight disease in Morocco, in all chickpea areas (Grewal, 1984; Singh, 1984) Bencheqroun et al. (2022) pointed out that *Didymella rabiei* (Kovatsch.) Arx. is the most devastating fungal infection of chickpea crops inflicting considerable yield and quality losses. Moreover, numerous fungal species associated with the chickpea diseases of wilt and root rot have been reported in Morocco, including *Fusarium oxysporum* f sp. *ciceris* (El Aoufir, 2001; El Bouazaoui et al., 2018); *Fusarium redolens* (Jiménez-Fernández et al., 2011); and *Rhizoctonia bataticola*, *R. solani*, and *Pythium* sp. (El Bouazaoui et al., 2018).

*Fusarium equiseti* has also been reported in other crop species, especially melon (*Cucumis melo*), soybean (*Glycine max*), cumin (*Cuminum cyminum*), cauliflower (*Brassica oleracea*), winter rapeseed (*Brassica napus*), tomato plant (*Solanum lycopersicum*), pepper (*Capsicum annuum*), and the cabbage *Brassica oleracea* (Adams et al., 1987; Chen et al., 2014; Gally et al., 1998; Goswami et al., 2008; Li et al., 2014; Ramchandra & Bhat 2012). The pathogen is also responsible for pre and postharvest decay of zucchini fruits (*Cucurbita pepo* L.) (Ezrari et al., 2019). It was recently isolated from the seeds of the fragrant wallflower (*Matthiola longipetala*) showing rot (Ivanović et al., 2020). Khan et al. (2021) noted that *F. equiseti* was responsible for seedling death in sugar beet.

Isolates of *Fusarium* and *F. solani* were recovered from necrotic lesions of chickpea roots in different chickpea growing areas (El Hazzat et al., 2019). Nevertheless, taxonomic confirmation of which species of *Fusarium* causes this necrosis is lacking because numerous species are important plant pathogens (Austwick, 1982). Additionally, the differentiation of *Fusarium* species

through morphological characters is imprecise; hence, the use of molecular techniques become more efficient and accurate for the discrimination of fungal species (Steenkamp et al., 2000).

Therefore, the present study was carried out to identify an isolate of *Fusarium* sp. collected for the first time in chickpea fields in Morocco from diseased chickpea plants based on morphological characters and molecular and pathological characterization via fulfillment of Koch's postulates.

## **2. Materials and methods**

### *2.1 Fungal material*

*Fusarium* isolates was obtained from necrotic lesions associated with infected stem samples of chickpea plants that were grown in different fields in Souk Tlat in Gharb Province of the Rabat-Sale-Kenitra region, Morocco. One hundred plants were chosen at random from the fields. One stem base sample from each plant was analyzed.

Pieces of diseased tissues were rapidly disinfected with 90% alcohol for 5 min, rinsed three times with sterile distilled water, and dried with sterile filter paper. Samples were then placed onto potato dextrose agar plates (BIOKAR Diagnostics) and incubated at 25 °C for 7 days. The colonies formed were transferred to potato sucrose agar (PSA) medium containing potato, sucrose, agar-agar, and distilled water. Agar plates were incubated in the same conditions and then observed for species determination. The *Fusarium* sp. isolate was cultivated in Petri dishes containing PSA medium. The medium was poured into Petri dishes containing 100 mg/L of chloramphenicol at a rate of 30 to 40 mL per dish. Incubation of cultures was performed in the dark at 25 °C for 7 days followed by macroscopic and microscopic characterization depending on the age of the cultures.

### *2.2 Morphological characterization*

Macroscopic examination of *Fusarium* sp. was carried out according to the development of the cultures on PSA medium. Observations focused on colony appearance, mycelium density, the

presence of the pinkish color of the colony, as well as growth and spore production. The microscopic characteristics of the *Fusarium* N3 isolate were determined under an optical microscope to confirm the species identity of this pathogen.

### 2.3 Molecular analysis and identification

Molecular identification of the *Fusarium* N3 isolate was performed after 5 days of culture on PSA medium. DNA extraction was performed according to the method described by Murray and Thompson (1980) and Doyle et al. (1987). DNA amplification of the internal transcribed spacer (ITS) rDNA region was performed using polymerase chain reaction (PCR) using universal primers ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') (White et al., 1990). The PCR reaction was carried out in a reaction mixture of 25 µL containing 5 µL of 5× buffer (MyTaq Reaction Buffer, Bioline, London, UK,), 1 µL of each primer (10 µM), 0.2 µL of MyTaq DNA polymerase (Bioline, London, UK) (5 U µL<sup>-1</sup>), 1–2 µL of template DNA (100 ng) and Milli-Q water to complete the volume. A Veriti thermal cycler (Applied Biosystems) was used for the PCR with the following conditions: initial denaturation at 95 °C for 1 min; 35 cycles of denaturation at 95 °C for 15 s, annealing at 52 °C for 20 s, and extension at 72 °C for 15 s; and a final elongation of 72 °C for 3 min. The quality of the PCR products was verified by electrophoresis on 1% agarose gel in the presence of a 100 bp molecular weight marker. Sequencing was performed using an ABI PRISM BigDye Terminator v.3.1 Ready Reaction Cycle Sequencing Kit and primer set ITS1 and ITS4. The sequencing products were run on an ABI PRISM 3130XL Genetic Analyzer (Applied Biosystems) using the POP-7 polymer. The sequence resulting from this study was submitted to GenBank under the accession no. MT111122,

The obtained ITS sequence was then compared with the homologous nucleotides sequence in the GenBank database using the Basic Local Alignment Search Tool (BLAST)

(<http://www.ncbi.nlm.nih.gov/BLAST>).

The seeds of a local variety of chickpea intended for pathogenicity testing were surface sterilized by soaking for 5 min in a 10% NaOCl solution, rinsed in sterile distilled water, and then dried on filter paper. The pathogenicity of the N3 isolate was checked through two inoculation techniques.

**Technique 1:** the surface-disinfected chickpea seeds were inoculated by soaking in water containing a conidial suspension of  $10^6$  spores/mL of the *Fusarium* isolate N3 at room temperature (20 °C) for 1 h. The control seeds were treated with only sterile distilled water. The inoculated and control chickpea seeds were sown in plastic pots (13 cm × 13.5 cm) containing autoclaved sieved Mamora forest soil at the rate of 5 seeds/pot. The Mamora forest soil used is loose, very sandy, and slightly basic pH (7.27), with an organic carbon content of 0.35% to 0.6% (Mouria, 2009). The soil was sieved and sterilized three times at an interval of 24 h at 200 °C for 2 h, and then distributed in the plastic pots (13 cm × 13.5 cm) at the rate of 2 kg of soil per pot.

**Technique 2:** The culture substrate was inoculated by pouring 15 mL of the conidial suspension of the N3 isolate (*Fusarium*) at a concentration of  $10^6$  spores/mL into each pot containing sterile soil. The fungus was allowed to settle for 48 h in the growing medium. The previously disinfected chickpea seeds were sown into these pots at the rate of 5 seeds per pot. Three replicates were prepared for each of the treatments (each pot was a replicate). Two lots of chickpea cultivation pots (control and inoculated) were brought back to the greenhouse to promote seed germination, plant growth, and symptom development monitoring.

After 4 weeks, plant emergence and plant survival were determined in pots containing inoculated seeds. In the pots with inoculated culture substrate according to the second inoculation technique, plant emergence and disease symptoms on chickpea plants were noted after 8 weeks, followed by

assessment of disease severity by calculating the leaf damage index according to the scale established by Douira and Lahlou (1989).

Notes	Appearance of leaves
0	Healthy appearance
1	Cotyledonary leaf: wilting or yellowing
2	Cotyledonary leaf: fall
3	True leaf: wilting or yellowing
4	True leaf: necrosis
5	True leaf: fall

The scores related to the number of leaves constitute the foliar alteration index, calculated according to the formula below (Douira & Lahlou, 1989):

$$\text{FAI} = [\sum(i \times X_i)] / 6 \times \text{NtF}$$

**FAI:** Foliar alteration index.

**i:** Leaf appearance notes 0 – 5.

**X<sub>i</sub>:** Number of leaves with note i.

**NtF:** Total number of leaves.

An average index was then calculated for each batch of plants.

At the end of the trial, the pots were brought back to the laboratory to re-isolate the pathogen from the different parts of the plants (roots, crown, stems, and leaf petioles) obtained either from the inoculated seeds (technique 1) or inoculated growing substrate (technique 2). The different parts were separated and disinfected with 95% alcohol for 2 min. Samples were then rinsed several times with sterile distilled water, dried quickly on sterile filter paper, transferred to PSA medium, and incubated in the dark at 25 °C. The microscopic observation was performed after 1 week.

The re-isolation percentage (RP%) was calculated using the following formula:  $RP = (N_s PX / N_t) \times 100$ , where  $N_s PX$  is the number of segments containing the fungal species X and  $N_t$  is the total number of segments used for re-isolation.

After 8 weeks, the lengths of the aerial and root parts of the chickpea plant were measured with a double decimeter and the number of leaves and pods of each plant was counted. The fresh weights of the aerial and root parts were recorded using a precision balance and the dry weight of these parts was recorded after being dried in the oven at 70 °C for 48 h.

### 3. Results

The isolate N3 of *Fusarium equiseti* was obtained from symptomatic samples among a complex of fusarium species including *F. solani* and *F. oxysporum*. The percentage of isolation was 3% in the case of *F. equiseti*.

On the PSA medium, the isolate N3 developed a colony with an abundant aerial mycelium which was fluffy and beige-white colored (Figure 1). Under a microscope, the mycelial filaments were septate. Chlamydospores were present (7 to 13 µm in diameter, spherical, globular, most often intercalary, solitary, or in pairs, and frequently as short chains). Macroconidia were numerous, slightly curved, usually with 5 to 6 septa and 31 to 45 µm long. The description was identical to that of *F. equiseti* (Corda) Saccardo, reported by Leslie and Summerell (2006) and Rafique et al. (2019). The identity of the isolate N3 was evaluated and confirmed. After identification, the isolate N3 was registered in the national database under voucher ID: RAB111030 and submitted to GenBank as accession no. MT111122. This sequence was 99.33% identical to *Fusarium equiseti*. The results of the pathogenicity tests using two inoculation methods demonstrated the pathogenic capacity of the isolate N3 of *F. equiseti* towards a local variety of chickpea. Few plants emerged

from seeds inoculated with the isolate N3. The majority of inoculated seeds turned rotten (Figure 2) compared to un-inoculated seeds which grew normally under greenhouse conditions.

Furthermore, chickpea plants obtained from seeds inoculated with *F. equiseti* showed a disruption in the growth parameters. The length of the plants, average number of leaves produced as well as the average number of branches formed were much lower in plants from inoculated seeds than those presented by the control plants; these parameters were 1.29 and 29.05 cm, 1.11 and 24.21 leaves, and 0 and 3.50 twigs, respectively. *Fusarium equiseti* isolate also adversely affected the growth of roots which exhibited a root length of 0.91 cm, and 0.78 and 0.13 g as fresh and dry weight of roots, respectively, in comparison with 27.16 cm, 7 g, and 5 g respectively, in the control plants.

The symptoms observed in chickpea plants developed on a culture substrate inoculated with *F. equiseti* according to technique 2 were variable: necrosis of the roots and crown followed by wilting of the plants. After 8 weeks of cultivation, the aggressiveness of infection induced by the *F. equiseti* isolate as estimated by calculating the foliar alteration index was 0.395 in plants grown in the culture substrate inoculated with a conidial suspension of the *Fusarium* isolate N3.

Regarding growth parameters, chickpea plants grown in the inoculated substrate displayed a length of 19.44 cm, an average of 18.66 leaves, and 2.08 twigs per plant compared to a length of 31.05 cm, 25.66 leaves, and 4.11 twigs per plant in control chickpea plants.

Plants which were grown in the culture substrate inoculated with *F. equiseti* also presented a lower root length, fresh and dry weight of root or aerial parts attaining 19.73 cm, 5.33 and 4.12 g, and 5.47 and 4.02 g, respectively, than plants grown in uninoculated soil (28.33 cm, 7.33 and 6.60 g, and 6.66 and 5.40 g, respectively).

The re-isolation performed on the plants grown either from inoculated seeds or on soil infested with *F. equiseti* confirmed the presence of the fungus in different parts of the plants. Using the

first technique, the highest percentage of the pathogen re-isolation was registered in the chickpea root with 70.22% followed by collar (70%), stem (60%), and leaf petiole (34.33 %), whereas percentages recorded in chickpea plants after soil inoculation were 84.77% (collar), 72.77% (root), 64.44% (stem), and 36.11% (petiole). *Fusarium equiseti* was consistently reisolated from infected seeds and collar tissues, satisfying Koch's postulates. To our knowledge, this is the first report of *Fusarium equiseti* causing seed rot and discoloration at the level of the crown and wilting of chickpea plants.




Fig. 1. Characteristic of *Fusarium equiseti* colony (A); chlamydospores (B); macroconidia and microconidia (C) Magnification, 400 $\times$ ; mounting medium, lactophenol cotton blue.




Fig. 2. Symptoms of stunting in chickpea plants (**a**) derived from seeds inoculated with *Fusarium equiseti* isolate (F.e) and control seedlings from non-inoculated chickpea seeds (T); **b**: symptoms of rot in inoculated chickpea seeds; **c**: brownish discoloration of crown in chickpea plants grown in *F. equiseti*-infested soil (F.e); **d**: chickpea plants grown in un-infested soil (T).

#### 4. Discussion

The *Fusarium* complex responsible for root rot and wilt diseases in chickpeas is diverse (Zemouli-

Benfreha 2014). One of the representative candidates of this complex is the *F. equiseti* that was isolated for the first time in Morocco from the roots of diseased chickpea plants and identified using morphological characters in addition to molecular characterization. In the current study, Koch's postulates were verified by inoculation of chickpea seeds and culture substrate.

The inoculated seeds expressed weak germination and growing ability with a high degree of rotting and decomposition. Chickpea plants from seeds sown in soil infested with a suspension of *F. equiseti* spores showed a range of deterioration symptoms such as root and crown necrosis, vascular browning, yellowing leaves, and stunting and wilting of the plants.

The fungus was re-isolated from the roots, crown, stems, and leaf petioles of the inoculated plants through seed dip inoculation or soil infestation with the spore suspension of *F. equiseti*. The ability of the pathogen to invade the upper levels of plants can be inferred from these results. Colonization of chickpea plants tissues by *F. equiseti* after inoculation affected their growth, leading to root and vegetative growth retardation including the production of leaves and twigs. All these types of symptoms were observed in chickpea plants inoculated with *F. solani* (El Hazzat et al., 2019) as well as in lentil and cumin (*Cuminum cyminum*) plants infected with *F. equiseti* (Rafique et al., 2019; Ramchandra et al., 2011). *Fusarium equiseti* is among the fungal species capable of attacking several legumes species, in which it can induce damping-off and root rot disease (Rubella et al., 2008). These authors have reported decayed seeds and reddish brown to black lesions on hypocotyl and roots of kidney bean (*Phaseolus vulgaris*), pea (*Pisum sativum*), and chickpea (*Cicer arietinum*) following inoculation with *F. equiseti* isolates originating from fields of ginseng. Similarly, the seeds that germinated in soil infected with *F. equiseti* resulted in plants that showed browning at the crown and stem base followed by wilting. Some symptoms on leguminous hosts bore a resemblance to those stated herein. On wild pigeon pea, this fungal species provokes foliar

chlorosis, browning and black discoloration of the stem, plant drying, and ultimately plant death (Mishra et al., 2021). Fernandez and Jefferson (2004) observed a discoloration in the subcrown stems and roots of cereal species.

This fungal species is capable of infecting seeds, roots, tubers, and fruits of several species of cultivated plants, such as cucurbits (Joffe & Palti 1967), cotton (Chimbekujwo, 2000), cowpea (Rodrigues & Menezes 2005), lentils (Chaudhary & Kaur 2002), sugar beet (Stojšin et al., 2001), potatoes (Rai, 1979; Theron & Holz 1989), citrus (Sukmawati & Miarsyah 2017), pine (Ocamb & Juzwik 1995), and even nursery plants (Bloomberg, 1981).

*Fusarium equiseti* has also been isolated from cereals such as corn, wheat, and barley (Ballois, 2012). This fungal species is responsible for rotting stems and premature wilting of corn plants (Swamy et al., 2020) and root rot in winter wheat (Booth 1971). Sometimes, this pathogen is associated with blight of wheat ears (Shaner, 2003; Gale, 2003; Tekauz et al., 2005; Wing et al., 1993; Xue et al., 2006) and rice panicles. Infection can also occur during grain storage or afterward (Hashem et al., 2010). In Morocco, several *Fusarium* species are associated with symptoms of root rot in cereals; the most common are *F. equiseti*, *F. culmorum*, *F. oxysporum*, and *F. solani* (Lyamani 1988).

*Fusarium equiseti*, isolated for the first time in Morocco, was isolated from the fungal complex associated with chickpea roots. Pathogenicity tests conducted in the greenhouse showed that this fungal species is endowed with significant pathogenicity towards this host plant, and the symptoms developed were similar to those observed in chickpea plants inoculated with *F. solani* (El Hazzat et al., 2019). Extending the surveys to other regions of Morocco is important to build a population of *F. equiseti* isolates and to determine the amplitude of the variation in the pathogenicity of this pathogen via other chickpea varieties grown in Morocco. With time and in the absence of an

effective control program, *Fusarium equiseti* will probably become an important pathogen and take its place among the other known diseases of chickpea.

## 5. Conclusion

To our knowledge, the present study showed for the first time that *Fusarium equiseti* can play serious role in the disease process of chickpea in Morocco causing seed rot, root and crown necrosis and leaf yellowing.

## Authors' Contributions:

**All authors** contributed to manuscript preparation: laboratory work, manuscript writing, identification of the fungus, translation of the manuscript, and result analysis

## References

- Adams, G. C.Jr., Gubler, W. D., & Grogan, R. G.,1987. Seedling disease of muskmelon and mixed melons in California caused by *Fusarium equiseti*. *Plant disease*, 71(4), 370-374.
- Austwick, P.K.C., 1982. *Fusarium* infections in man and animals. In MO Moss and JE Smith, ed. *The applied mycology of Fusarium*. Cambridge: Cambridge University Press. pp. 129-140.
- Ballois, N., 2012. Characterization of the diversity of *Fusarium* species and their mycotoxigenic potential on French cereals. *Sciences de l'environnement*. Université de Lorraine URL : [https://www.researchgate.net/publication/332369909\\_Characterisation\\_of\\_the\\_diversity\\_of\\_Fusarium\\_species\\_and\\_their\\_mycotoxigenic\\_potential\\_on\\_French\\_cereals](https://www.researchgate.net/publication/332369909_Characterisation_of_the_diversity_of_Fusarium_species_and_their_mycotoxigenic_potential_on_French_cereals).
- Bencheqroun, S. K., Ahmed, S., Imitiaz, M., Hamwieh, A., Udupa, S.M., Sahri, A., Aouzal, S., Kehel, Z., 2022. Pathogen diversity and mating types of *Didymella rabiei* isolates collected from Morocco. *Current Plant Biology* 29, 100231.

- Bloomberg, W.J., 1981. Diseases caused by *Fusarium* in forest nurseries. In: *Fusarium: Diseases, biology, and taxonomy*. In: Nelson PE, Toussoun TA, Cook R. editors. University Park, USA: The Pennsylvania State University Press pp. 178–187.
- Booth, C., 1971. *The genus fusarium*. Commonwealth Mycological Institute, Kew, Surrey, England. 237 pp.
- Chaudhary, R.G., Kaur, A., 2002. Wilt disease as a cause of shift from lentil cultivation in Sangod Tehsil of Kota (Rajasthan). *Indian Journal of Pulses Research* 15, 193-194.
- Chen, Y., Zhou, Q., Strelkov, S. E., & Hwang, S. F., 2014. Genetic diversity and aggressiveness of *Fusarium* spp. isolated from canola in Alberta, Canada. *Plant disease*, 98(6), 727-738.
- Chimbekujwo, I.B., 2000. Frequency and pathogenicity of fusarium wilts (*Fusarium solani* and *Fusarium equiseti*) of cotton (*Gossypium hirsutum*) in Adamawa in Nigeria. *Revista de Biología Tropical* 48, 01-05.
- Douira, A., & Lahlou, H., 1989. Variabilité de la spécificité parasitaire chez *Verticillium albo-atrum* Reinke et Berthold, forme à microsclérotés. *Cryptogamie. Mycologie*, 10(1), 19-32.
- Doyle, J. J., & Doyle, J. L., 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue (No. Research).
- El Aoufir, A., 2001. Étude du flétrissement vasculaire du pois chiche (*Cicer arietinum*) causé par *Fusarium oxysporum* f. sp. *ciceri*. Évaluation de la fiabilité de l'analyse isoenzymatique et de la compatibilité végétative pour la caractérisation des races physiologiques. **Thèse de doctorat**, Université Laval Québec P,161.

- El bouazaoui, A., Maafa, I., Ahmed Kemal, S., Douira, A., Udupa, S.M., 2018. Status of wilt and root rot diseases of Kabuli chickpea in some regions of Morocco. <https://repo.mel.cgiar.org/handle/20.500.11766/9158>
- ELHazzat, N., Ouazzani Touhami, A., Chliyeh, M., Selmaoui, K., Douira, A., 2019. Effet d'un inoculum endomycorhizien composite sur la manifestation de la fusariose du pois chiche (*Cicer arietinum*), causée par *Fusarium solani*. *Plant Cell Biotechnology and Molecular Biology* 20, 486–500.
- Ezrari, S., Lahlali, R., Radouane, N., Tahiri, A., & Lazraq, A., 2020. First report of *Fusarium equiseti* causing pre-and postharvest fruit rot on zucchini in Morocco. *Journal of Plant Pathology*, 102(1), 251-251.
- Fernandez, M.R., Jefferson, P.G., 2004. Fungal populations in roots and crowns of common and durum wheat in Saskatchewan. *Can. J. Plant Pathol.* 26:325-334.
- Gale, L.R., 2003. Population biology of *Fusarium* species causing head blight of grain crops. *Fusarium head blight of wheat and barley*, Alberta Agriculture and Forestry, Government of Alberta. P120-143.
- Gally, T., Gonzalez, B., Sobero y Rojo, M. P., & Lori, G., 1998. Plumele soft rot caused by *Fusarium equiseti*, *F. oxysporum*, and *F. pallidoroseum* on soybean seedlings in Argentina. *Plant disease*, 82(9), 1063-1063.
- Goswami, R. S., Dong, Y., & Punja, Z. K., 2008. Host range and mycotoxin production by *Fusarium equiseti* isolates originating from ginseng fields<sup>1</sup>. *Canadian Journal of Plant Pathology*, 30(1), 155-160.

Gupta, S., Nadarajan, N., & Gupta, D. S. (Eds.), 2014 . Legumes in the omic era (Vol. 348).

Springer India 2014, DOI: <https://doi.org/10.1007/978-81-322-2023-7>

Grewal, J. S. (1984). Evidence of physiologic races in *Ascochyta rabiei* of chickpea. In M.

Saxena & K. Singh (Eds.), Proceedings of the Workshop on Ascochyta Blight and

Wintering of Chickpeas (pp. 55–65). Aleppo, Syria: ICARDA.

Hashem, M., Moharam, A. M., Zaied, A. A., & Saleh, F. E. M., 2010. Efficacy of essential oils in

the control of cumin root rot disease caused by *Fusarium* spp. Crop protection, 29(10), 1111-

1117.

Houasli, C., Idrissi, O., Nsarellah, N., 2020. Chickpea genetic improvement in Morocco: State of

the art, progress and prospects. Moroccan Journal of Agricultural Sciences, 1(1). in California

caused by *Fusarium equiseti*. Plant disease 71, 370-374.

Ivanović, Ž., Milošević, D., Ignjatov, M., Marjanović-Jeromela, A., Karaman, M., & Grahovac,

M., 2020. First Report of *Fusarium equiseti* as the Causal Agent of Seed Rot of *Matthiola*

*longipetala* in Serbia. Plant Disease, 104(9), 2516.

Joffe, A.Z., Palti, J., 1967. *Fusarium equiseti* (Cda) Sacc. Isr J Bot Basic Appl Plant Sci, P 1–18.

Khan, M. F. R., Liu, Y., Bhuiyan, M. Z. R., Lakshman, D., Liu, Z., & Zhong, S. (2021). First

Report of *Fusarium equiseti* Causing Seedling Death on Sugar Beet in Minnesota, USA. Plant

Disease, 105(7).

Leslie, J.F., & Summerell, B.A., 2006. The *Fusarium* Laboratory Manual. Blackwell Publishing.

Ames, Iowa, USA , P. 388.

Li, P. P., Cao, Z. Y., Wang, K., Zhai, H., Jia, H., Liu, N., ... & Dong, J. G., 2014. First report of

*Fusarium equiseti* causing a sheath rot of corn in China. Plant disease, 98(7), 998-998.

Lyamani, A., 1988. Wheat root rot in West Central Morocco and effects of *Fusarium culmorum* and *Helminthosporium sativum* seed and soil-borne inoculum on root rot development, plant emergence and crop yield. Ph.D. Thesis Iowa State University P 270.

MAPMDREF., 2020a, Emblavement des légumineuses principales. Retrieved from [https://www.onicl.org.ma/portail/sites/default/files/FichierPage/StatEmblLeg\\_20-01-2020\\_0.pdf](https://www.onicl.org.ma/portail/sites/default/files/FichierPage/StatEmblLeg_20-01-2020_0.pdf)

MAPMDREF., 2020b. Production des légumineuses principales. Retrieved from [https://www.onicl.org.ma/portail/sites/default/files/FichierPage/Prod\\_Statleg\\_20-01-2020\\_0.pdf](https://www.onicl.org.ma/portail/sites/default/files/FichierPage/Prod_Statleg_20-01-2020_0.pdf)

Millan, T., Winter, P., Jüngling, R., Gil, J., Rubio, J., Cho, S., ... & Muehlbauer, F. J., 2010. A consensus genetic map of chickpea (*Cicer arietinum* L.) based on 10 mapping populations. Euphytica, 175(2), 175-189.

Mishra, R. K., Mishra, M., Bohra, A., Naik SJ, S., Saabale, P. R., Kumar, K., ... & Singh, N. P., 2021. First Report of *Fusarium equiseti* Causing Wilt of *Cajanus scarabaeoides*, a Wild Relative of Pigeonpea, in India. Plant Disease, 105(9), 2735.

Mouria, B., 2009. Contribution à la lutte biologique contre la pourriture grise et la verticilliose de la tomate cultivée sous serre par utilisation du compost et les *Trichoderma* spp. seul ou en combinaison (Doctoral dissertation, Thèse de Doctorat National. Université Ibn Tofail. Faculté des Sciences, Kénitra, Maroc).

- Murray, M. G., & Thompson, W., 1980. Rapid isolation of high molecular weight plant DNA. *Nucleic acids research*, 8(19), 4321-4326.
- Ocamb, C.M., Juzwik, J., 1995. *Fusarium* species associated with rhizosphere soil and diseased roots of eastern white pine seedlings and associated nursery soil. *Can. J. Plant Pathol* 17, 325–330.
- Rafique, K., Rauf, C. A., & Kang, S., 2019. First report of *Fusarium equiseti* causing wilt on lentils (*Lens culinaris* Medik.) in Pakistan. *Journal of Plant Pathology*, 102(2), 571-571.
- Rai, R.P., 1979. *Fusarium equiseti* (Corda) Sacc. causing dry rot of potato-tuber a new report. *Curr.Sci.* P.1043–1045.
- Ramchandra, S., & Bhatt, P. N., 2012. First report of *Fusarium equiseti* causing vascular wilt of cumin in India. *Plant disease*, 96(12), 1821-1821.
- Ramchandra, S., Suthar Bhatt, P.N., 2011. Morphological and molecular identification of *Fusarium* isolated from cumin wilt. *Internat. J. Plant Protec* 4, 359-362.
- Rodrigues, A.A.C., Menezes, M., 2005. Identification and pathogenic characterization of endophytic *Fusarium* species from cowpea seeds. *Mycopathologia* 159, 79–85.
- Rubella, S.G., Yanhong, D., Zamir, K.P., 2008. Host range and mycotoxin production by *Fusarium equiseti* isolates originating from ginseng fields. *Canadian Journal of Plant Pathology* 2, 155-160
- Shaner, G.E., 2003. Epidemiology of *Fusarium* Head Blight of small grain cereals in North America. In: *Fusarium Head Blight of Weath and Barley*. Leonard, KJ , Bushnell, WR. (eds). APS Press, Phytopathology, St. Paul, MN, USA. P. 84-119.

- Steenkamp, E.T., Britz, H., Coutinho, T., Wingfield, B., Marasas, W., Wingfield, M., 2000. Molecular characterization of *Fusarium subglutinans* associated with mango malformation. *Molecular Plant Pathology* 1, 3, 187- 193.
- Stojšin, V., Balaz, F., Bagi, F., and Keljacki, I., 2001. Pathogenicity of *Fusarium* spp. isolates from sugar beet root. *Zastita- Bilja*, 52, 241–249.
- Sukmawati, D., Miarsyah, M., 2017. Pathogenic activity of *Fusarium equiseti* from plantation of citrus plants (*Citrus nobilis*) in the village Tegal Wangi, Jember Umbulsari, East Java, Indonesia. *Asian J Agri & Biol.* 5, 202-213.
- Singh, K. B. (1984). International Testing of Chickpeas for Resistance to Ascochyta Blight. *Plant Disease*, Vol. 68, p. 782. <https://doi.org/10.1094/pd-69-782>
- Swamy, S.D., Mahadevakumar, S., Hemareddy, H.B., Amruthesh, K.N., Mamatha, S., Kunjeti S.G., Swapnil, R., Kumard, T.V., Lakshmidēvi, N., 2020. First report of *Fusarium equiseti*–the incitant of post flowering stalkrot of maize (*Zea mays* L.) in India. *Crop Protection*.129, 105-035.
- Tekauz, A., Stulzer, M., Beyene, M., 2010. *Fusarium* head blight of winter wheat in Manitoba in 2009. *Canadian plant disease survey = Inventaire des maladies des plantes au Canada*, P. 47–48.
- Theron, D.J., Holz, G., 1989. *Fusarium* species associated with dry and stem-end rot of potatoes in South Africa. *Phytophylactica* 21, 175–181.
- White, T.J., Bruns, T., Lee, S., Taylor, J., 1990. Amplification and direct sequencing of fungal ribosomal RNA for phylogenetics. In: MA Innes, DH Gelfand, JJ Sninsky, TJ White eds,

PCR protocols. A guide to methods and applications, Academic Press, Inc., San Diego, California, 315-322.

Xue, A.G., Ho, K.M., Butler, G., Vigier, B.J, Babcock, C., 2006. Pathogenicity of *Fusarium* species causing head blight in barley. *Phytoprotection* 87, 55–61.

Zemouli-Benfreh, F., Henni, D-e., Merzoug, A., 2014. *Fusarium* wilt of Chickpea (*Cicer arietinum* L.) in North-west Algeria. *African Journal of Agricultural Research* 9, 168-175.

POSTPRINT