

Appendix S1

Supplementary Information

List of localities and ecological data (habitats, substrata and months of observation)

ASCOMYCOTA

***Aleuria aurantia* (Pers.) Fuckel**

Kaczawskie Mts.: Bystrzyca 1.1 km S (WIF-132b), Wleń 1.1 km E (WIF-136f); IX-X; forest road; soil.

Karkonosze Mts.: Szklarska Poręba; X; mixed forest; soil. **Kaczawskie Foothills:** Bystrzyca 1.9 km N (WIF-14j), Wleń 1.5 km N (WIF-31a), Wleń 2.6 km N (WIF-25a); IX-X; deciduous forest (log landing), logging road, spruce forest; soil.

***Aleuria splendens* (Quél.) J. Breitenb. & F. Kränz. ex Bon & Courtec.**

Sudetes Foreland: Szklary 0.5 km W; X; overgrown spoil tip; soil.

***Ascobolus stercorarius* (Bull.) J. Schröt. (= *A. furfuraceus* Pers.)**

Kaczawskie Foothills: Przeździec 1.7 km E (WIF-12k); IV; mixed forest; *Capreolus capreolus* dung.

***Ascocoryne albida* (Berk.) Seifert**

Kaczawskie Foothills: Bystrzyca 2.1 km NW (WIF-26c); X; deciduous forest; ?*Populus tremula* wood.

***Ascocoryne cylichnium* (Tul.) Korf**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); X; oak-hornbeam forest; *Carpinus betulus* wood.

***Ascocoryne* sp. (*A. inflata* sensu Baral)**

Kaczawskie Foothills: Bystrzyca 2.1 km N (WIF-14f); XI; mixed sapling stand; wood.

***Ascodichaena rugosa* Butin**

Izerskie Foothills: Pilchowice 1.1 km S (MaF – Dziki Gully), Wleń 1.4 km SW (ŁuF-58m); II-V; beech forest, deciduous forest; *Fagus sylvatica* trunk.

***Ascotremella faginea* (Peck) Seaver**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VI-XII; spruce forest; *Fraxinus* sp. log, deciduous wood. **Kaczawskie Foothills:** Belczyna 0.7 km S (WIF-123g), Belczyna 0.8 km S (WIF-123k); XI; spruce forest; *Quercus* sp. branch, *Fraxinus* sp. log.

***Bisporrella citrina* (Batsch) Korf & S.E. Carp.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); XI; deciduous forest; ?ash wood. **Kaczawskie Foothills:** Bystrzyca 1.7 km N (WIF-16); X; spruce forest; *Betula* sp. wood.

***Bulgaria inquinans* (Pers.) Fr.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; *Quercus* sp. log. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), VI; deciduous forest; *Quercus* sp. wood. **Kaczawskie Foothills:** Bystrzyca 1.5 km N (WIF-19a), Bystrzyca 2.1 km N (WIF-14f), Bystrzyca 2.4 km NW (WIF-22), Radomiłowice 1 km SW (near WIF-11b), Wleń 1.1 km N (WIF-31j); IX-XI; deciduous forest, mixed forest; *Quercus* sp. log and trunk, *Carpinus betulus* log, stump.

***Calycina* sp. (*C. subherbarum* sensu Baral)**

Kaczawskie Foothills: Bystrzyca 2.5 km NW (WIF-26k); X; coniferous forest margin; decaying stems of herbaceous plants.

***Chaetosphaeria myriocarpa* (Fr.) C. Booth**

Kaczawskie Foothills: Wleń 2 km N (WIF-30f); VII; under lime tree; *Tilia* sp. wood.

***Cheilymenia vitellina* (Pers.) Dennis**

Kaczawskie Foothills: Belczyna 1 km SE (WIF-122b); VIII; beech forest; soil.

***Cordyceps militaris* (L.) Link**

Kaczawskie Mts.: Bystrzyca 1.2 km S (WIF-133a); IX; ash riparian forest; pupa. **Kaczawskie Foothills:** Bystrzyca 1.1 km SE (WIF-123m); IX-X; deciduous forest; pupa of *Lepidoptera*, unidentified insect.

***Cudoniella acicularis* (Bull.) J. Schröt.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-58m); XII; deciduous forest; ?soil.

***Cyathicula coronata* (Bull.) De Not. (= *Crocicreas coronatum* (Bull.) S.E. Carp.)**

Kaczawskie Foothills: Bystrzyca 2.5 km NW (WIF-26k); X; coniferous forest margin; decaying stems of herbaceous plants.

***Cyathicula cyathoidea* (Bull.) Thüm. (= *Crocicreas cyathoideum* (Bull.) S.E. Carp.)**

Kaczawskie Foothills: Radomiłowice 0.7 km SW (WIF-11a); V; mixed forest; decaying stems of herbaceous plants.

***Diatrype disciformis* (Hoffm.) Fr.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); II; deciduous forest; deciduous branch.

***Dumontinia tuberosa* (Bull.) L.M. Kohn**

Kaczawskie Mts.: Bystrzyca 1 km SE (WIF-131a); IV; riparian forest; *Anemone* sp. roots.

***Elaphomyces asperulus* Vittad.**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt), Wleń 2.2 km SW (ŁuF – Gniazdo Mt); IV-V; spruce-pine forest, mixed forest; soil.

***Elaphomyces granulatus* Fr.**

Karkonosze Mts.: Szklarska Poręba (SzPF-299a); X; spruce forest; soil. **Izerskie Foothills:** Wleń 1.7 km SW (ŁuF – Gniazdo Mt), Wleń 2.3 km NW (WIF-46i); IV-VIII; mixed forest, spruce forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.1 km S (WIF-14h); mixed forest; soil.

***Elaphomyces muricatus* Fr.**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt); XI; spruce-pine forest; soil.

***Encoelia furfuracea* (Roth) P. Karst.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); II; deciduous forest; *Corylus avellana* branch. **Kaczawskie Foothills:** Bełczyna 0.8 km N (WIF-122s), Bełczyna 1.3 km E (WIF-122d), Bystrzyca 1.2 km SE (WIF-126n, 126r), Bystrzyca 1.8 km E (WIF-126m), Bystrzyca 2.2 km N (WIF-15b), Radomiłowice 1.4 km W (WIF-12Ak), Wleń 1.8 km NE (WIF-33d), Wleń 2.6 km N (WIF-25a), near Wąwóz Lipy Res.; XI-II; riparian forest, oak-hornbeam forest, deciduous sapling stand, pine forest, mixed forest; *Corylus avellana* branch, *Alnus incana* branch.

***Erinella aeruginosa* Henn.**

Kaczawskie Mts.: Wleń 1.3 km NE (WIF-136b); IX; mixed forest; *Quercus* sp. wood.

***Geoglossum cookeanum* Nannf. ex Minter & P.F. Cannon**

Kaczawskie Mts.: Góra Miłek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Geoglossum fallax* E.J. Durand**

Kaczawskie Mts.: Buki Sudeckie Res.; IX; *Dentario enneaphylli-Fagetum*; soil.

***Geoglossum glabrum* Pers.**

Izerskie Foothills: Zalesie 0.8 km NE (LuF); VIII; mire; *Sphagnum* sp.

***Geoglossum umbratile* Sacc.**

Izera Mts.: by the tourist trail between Jakuszyce and Orle; IX; soil. **Kaczawskie Mts.:** Góra Miłek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Glutinoglossum glutinosum* (Pers.) Hustad, A.N. Mill., Dentinger & P.F. Cannon**

Kaczawskie Mts.: Buki Sudeckie Res.; IX; *Dentario enneaphylli-Fagetum*; soil.

***Geopyxis carbonaria* (Alb. & Schwein.) Sacc.**

Kaczawskie Mts.: Bystrzyca 1.4 km SE (WIF-130f); X; spruce forest (old campfire); charcoal. **Karkonosze Mts.:** Jakuszyce 1 km S; VIII; coniferous forest (old campfire); charcoal. **Kaczawskie Foothills:** Radomiłowice 1.1 km SW (WIF-11g), Wleń 1.7 km NE (WIF-27h); VII; coniferous forest (old campfire), mixed forest (old campfire); charcoal.

***Gyromitra esculenta* (Pers.) Fr.**

Karkonosze Mts.: Szklarska Poręba (by the yellow tourist trail); V; roadside; soil.

***Gyromitra infula* (Schaeff.) Quél.**

Karkonosze Mts.: Szklarska Poręba (Kilińskiego St.); X; ruderal place; wood. **Kaczawskie Foothills:** Bystrzyca 2.5 km NE (WIF-15h); X; deciduous forest; wood.

***Helvella acetabulum* (L.) Quél.**

Izerskie Foothills: Radomice (limestone quarry); V; overgrown quarry; soil. **Sudetes Foreland:** Szklary 0.4 km W; IV; overgrown spoil tip; soil.

***Helvella crispa* (Scop.) Fr.**

Kaczawskie Foothills: Wleń 2 km N (WIF-30f); VII; under lime tree; soil.

***Helvella elastica* Bull.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; deciduous forest; soil.

***Helvella ephippium* Lév.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest (logging road); soil.

***Helvella lacunosa* Afzel.**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry); IX; overgrown quarry; soil. **Kaczawskie Foothills:** Radomiłowice 0.7 km S (WIF-11c), Wleń 2.6 km N (WIF-25a); VIII-X; oak-hornbeam forest, forest margin (log landing); soil.

***Helvella latispora* Boud.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF-64 – Gniazdo Mt); VI; deciduous forest; soil.

***Helvella leucomelaena* (Pers.) Nannf.**

Sudetes Foreland: Szklary 17, Szklary 0.4 km W; IV; orchard, overgrown spoil tip; soil.

***Helvella macropus* (Pers.) P. Karst.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.7 km SW (ŁuF – Gniazdo Mt); VIII-X; deciduous forest, spruce forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.5 km NW (WIF-22p); VIII; mixed forest; soil.

***Helvella pezizoides* Afzel.**

Sudetes Foreland: Szklary 0.5 km W; X; overgrown spoil tip; soil.

***Helvella fibrosa* (Wallr.) Korf (= *H. villosa* (Hedw.) Dissing & Nannf.)**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; deciduous forest (logging road); soil.

***Holwaya mucida* (Schulzer) Korf & Abawi**

Kaczawskie Foothills: Bystrzyca 1.8 km N (WIF-16d); X; oak-hornbeam forest; *Tilia* sp. log.

***Humaria hemisphaerica* (Hoffm.) Fuckel**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1.4 km SE (WIF-130f); VII; oak-hornbeam forest, spruce forest; soil. **Izerskie Foothills:** Wleń 1.5 km SW (ŁuF-64 – Gniazdo Mt), Wleń 1.4 km NW (WIF-46); VII-VIII; deciduous forest, mixed forest; soil. **Kaczawskie Foothills:** Belczyna 1.8 km SE (WIF-124a), Wleń 2.6 km N (WIF-25a); VIII; deciduous forest; soil.

***Hyaloscypha daedaleae* Velen.**

Kaczawskie Foothills: Bystrzyca 2.1 km N (WIF-14f); XI; mixed sapling stand; wood.

***Hymenoscyphus imberbis* (Bull.) Dennis**

Kaczawskie Mts.: Buki Sudeckie Res.; X; *Carici remotae-Fraxinetum*; wood.

***Hymenoscyphus pseudoalbidus* V. Queloz, C.R. Grünig, R. Berndt, T. Kowalski, T.N. Sieber & O. Holdenrieder**

Kaczawskie Foothills: Bystrzyca 2.4 km NW (WIF-211); VI; deciduous forest; *Fraxinus* sp. petioles.

***Hymenoscyphus scutula* (Pers.) W. Phillips**

Kaczawskie Mts.: Milek Mt; X; meadow; decaying herbaceous plants.

***Hypocrea auranteffusa* Jaklitsch**

Kaczawskie Foothills: Bystrzyca 2.5 km NE (WIF-15h); VIII; deciduous forest; deciduous wood (near rizomorphs of ?*Armillaria*).

***Hypocrea citrina* (Pers.) Fr.**

Kaczawskie Mts.: Bystrzyca 1.5 km SE (WIF-130d); VIII; wet spruce forest; litter, stump. **Kaczawskie Foothills:** Bystrzyca 2.1 km N (WIF-14h), Wleń 1.9 km NE (WIF-28d); VIII-IX; mixed forest; litter, *Betula* sp. stump.

***Hypocrea gelatinosa* (Tode) Fr.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); IX; oak-hornbeam forest; deciduous wood. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); XI; riparian forest; deciduous wood. **Kaczawskie Foothills:** Przeździec 1.4 km E (WIF-12d), Wleń 2.3 km NE (WIF-25c); VII-VIII; mixed forest, spruce forest; wood, ?ash wood.

***Hypocrea schweinitzii* (Fr.) Sacc.**

Kaczawskie Foothills: Wleń 2.6 km N (WIF-25a); VIII; mixed forest; *Picea abies* wood.

***Hypomyces rosellus* (Alb. & Schwein.) Tul. & C. Tul.**

Kaczawskie Mts.: Bystrzyca 1.2 km S (WIF-133b); IX; mixed forest; deciduous wood.

***Hypoxylon howeanum* Peck**

Kaczawskie Foothills: Bystrzyca 1.8 km N (WIF-16a); X; oak-hornbeam forest; *Quercus* sp. branch.

***Jackrogersella multiformis* (Fr.) L. Wendt, Kuhnert & M. Stadler (= *Annulohypoxylon multiforme* (Fries) Y.M. Yu, Rogers & Hsieh var. *multiforme*)**

Kaczawskie Foothills: Przeździec 0.9 km S (WIF-22k); IV; riparian forest; ?*Quercus* sp. log.

***Kretzschmaria deusta* (Hoffm.) P.M.D. Martin**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt); IV; beech forest; *Fagus sylvatica* stump.

***Lachnellula occidentalis* (G.G. Hahn & Ayers) Dharne**

Kaczawskie Foothills: Wleń 1.9 km NE (WIF-28d); V; mixed forest; *Larix* sp. twigs.

***Lasiobolus ciliatus* (J.C. Schmidt) Boud. (= *L. equinus* (O. F. Müll.) P. Karst.)**

Kaczawskie Foothills: Przeździec 1.7 km E (WIF-12k); IV; mixed forest; *Capreolus capreolus* dung.

***Leotia lubrica* (Scop.) Pers.**

Kaczawskie Mts.: Radzimowice, Tarczyn 1.1 km NE (WIF-134b); VIII-IX; spruce forest, under spruce; soil. **Izerskie Foothills:** Wleń 2.3 km NW (WIF-46j); VIII; mixed forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.5 km NE (WIF-15h), Wleń 1.9 km NE (WIF-27d), Wleń 2.3 km NE (WIF-26m); VIII-IX; deciduous forest, mixed forest near stream; wood, soil.

***Melastiza boudierii* (Höhn.) Le Gal**

Kaczawskie Foothills: Przeździec 0.8 km SW (WIF-22l); spruce forest (logging road); soil.

***Microglossum fuscrobens* Boud.**

Kaczawskie Mts.: Buki Sudeckie Res.; VIII; *Dentario enneaphylli-Fagetum*; soil.

***Microglossum nudipes* Boud.**

Kaczawskie Mts.: Góra Milek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Microglossum viride* (Pers.) Gillet**

Karkonosze Mts.: Szklarska Poręba (Złoty Potok Valley); XI; beech forest (helocrene); soil.

***Mitrula paludosa* Fr.**

Karkonosze Mts.: Karkonosze NP (near red tourist trail to Śnieżka Mt); VI; spruce forest (wet roadside); litter.

Izerskie Foothills: Zalesie 0.8 km N (LuF); V-VII; mire; soil, *Sphagnum* sp.

***Mollisia ventosa* P. Karst.**

Kaczawskie Mts.: near Bystrzyca (WIF-134); VII; riparian forest; deciduous wood.

Morchella esculenta* (L.) Pers. f. *esculenta

Kaczawskie Mts.: Buczyna Storczykowa na Białych Skałach Res.; V; beech forest; soil. **Kaczawskie Foothills:** Leszczyna 1 km N (near Ciche Szczęście mine); IV; oak-hornbeam forest; soil. **Sudetes Foreland:** Szklary 17; IV; orchard; soil.

***Neonectria coccinea* (Pers.) Rossman & Samuels (= *Nectria coccinea* (Persoon) Fries)**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); XI; deciduous forest; *Acer pseudoplatanus* branch.

***Neonectria ditissima* (Tul. & C. Tul.) Samuels & Rossman (= *Neonectria galligena* (Bresadola) Rossman & Samuels)**

Kaczawskie Foothills: Bystrzyca 1.4 km N (WIF-19c); IV; mixed forest; *Fraxinus* sp. branch.

***Nigrograna fuscidula* (Sacc.) Jaklitsch & Voglmayr**

Kaczawskie Foothills: Bystrzyca 2.2 km N (WIF-13d); III; deciduous forest; *Salix* sp. branch.

***Ophiocordyceps entomorrhiza* (Dicks.) G.H. Sung, J.M. Sung, Hywel-Jones & Spatafora**

Kaczawskie Mts.: Tarczyn 0.8 km E (WIF-134h), Góra Milek Res., Buki Sudeckie Res.; VIII-IX; *Cephalanthero-Fagetum*, *Dentario enneaphylli-Fagetum*, mixed forest; imago of *Coleoptera*. **Izerskie Foothills:** Wleń 2.7 km NW (WIF-44k); VI; riparian forest; imago of *Coleoptera*. **Kaczawskie Foothills:** Bystrzyca 1.4 km N (WIF-19b); X; oak-hornbeam forest; imago of *Coleoptera*.

***Orbilia xanthostigma* (Fr.) Fr.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133); VI; riparian forest; wood. **Kaczawskie Foothills:** Bystrzyca 1.8 km N (WIF-16a), Wleń 1.4 km N (WIF-31c); VIII-X; oak-hornbeam forest, mixed forest; wood, deciduous wood.

***Otidea alutacea* (Pers.) Massee**

Izerskie Foothills: Wleń 1.4 km NW (WIF-46g); IX; mixed forest; soil. **Kaczawskie Foothills:** Bełczyna 0.7 km S (WIF-123d), Wleń 2.5 km N (WIF-25a); VII-IX; oak-hornbeam forest, deciduous forest; soil.

***Otidea mirabilis* Bolognini & Jamoni**

Izerskie Foothills: Wleń 1.6 km NW (WIF-46f); VIII; spruce forest on riparian habitat (logging road); soil. **Kaczawskie Foothills:** Bystrzyca 1.4 km N (WIF-16k), Wleń 2.6 km N (WIF-25a); VIII-IX; oak-hornbeam forest; soil.

***Otidea onotica* (Pers.) Fuckel**

Izerskie Foothills: Wleń 1.3 km SW (ŁuF – Gniazdo Mt); IX; oak-hornbeam forest; soil. **Kaczawskie Foothills:** Bystrzyca (near WIF-20); Wleń 1.9 km NE (WIF-27f), Wleń 2.5 km N (WIF-25a); VIII-IX; deciduous forest, mixed forest, oak-hornbeam forest; soil.

***Otidea tuomikoskii* Harmaja**

Kaczawskie Mts.: Tarczyn 0.9 km NE (WIF-134c); IX; spruce forest; soil.

***Pachyella babingtonii* (Berk. & Broome) Boud.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VII; oak-hornbeam forest; wood. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest; *Betula* sp. wood. **Kaczawskie Foothills:** Bystrzyca 2 km NE (WIF-15s); X; deciduous forest; deciduous wood.

***Peziza arvernensis* Boud.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 2.1 km NE (WIF-45a); V-X; deciduous forest; wood, soil. **Kaczawskie Foothills:** Bełczyna 1 km SE (WIF-122f), Bystrzyca 1.4 km N (WIF-16k), Radomiłowice 1 km SW (WIF-11b); IX-XI; beech forest, oak-hornbeam forest, mixed forest; soil, wood, *Fagus sylvatica* sawdust.

***Peziza badia* Pers.**

Karkonosze Mts.: Jakuszyce 1 km S, Szklarska Poręba (Złoty Potok Valley), Szklarska Poręba (central station of the ski lift on Szrenica Mt); VIII-XI; coniferous forest, spruce forest, roadside; soil. **Kaczawskie Foothills:** Wleń 1.4 km N (WIF-31c); X; mixed forest; soil.

***Peziza echinospora* P. Karst.**

Kaczawskie Foothills: Radomiłowice 1.1 km SW (WIF-11g); VII; mixed forest (old campfire); charcoal.

***Peziza gerardii* Cooke**

Kaczawskie Mts.: Góra Milek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Peziza micropus* Pers.**

Kaczawskie Foothills: Rząśnik 1.3 km NW (WIF-125g); X; mixed forest; wood.

***Peziza saniosa* Schrad. ex J.F. Gmel.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; soil. **Izerskie Foothills:** Wleń 1.4 km NW (WIF-46a); VIII; deciduous sapling stand; soil.

***Peziza succosa* Berk.**

Izerskie Foothills: Wleń 1.6 km NW (WIF-46f), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VII-IX; spruce forest, deciduous forest; soil.

***Peziza vesiculosa* Bull.**

Kaczawskie Foothills: Bystrzyca 1.6 km N (WIF-16b); IX; forest; straw.

***Phaeohelotium monticola* (Berk.) Dennis**

Kaczawskie Foothills: Bystrzyca 2 km S (WIF-126w); VII; deciduous forest; acorn cupules.

***Plicaria endocarpoides* (Berk.) Rifai (= *P. leiocarpa* (Curr.) Boud.)**

Kaczawskie Foothills: Wleń 1.7 km NE (WIF-27h); I; spruce-pine forest (old campfire); charcoal.

***Propolis farinosa* (Pers.) Fr.**

Kaczawskie Mts.: Milek Mt; X; abandoned orchard; *Malus* sp. branch. **Izerskie Foothills:** Wleń 2.2 km SW (ŁuF – Gniazdo Mt); I; mixed forest; ?hornbeam log.

***Ptychoverpa bohémica* (Krombh.) Boud. (= *Verpa bohémica* (Krombh.) J. Schröt.)**

Kaczawskie Foothills: Bełczyna 0.6 km W (WIF-130); IV; under willows; soil. **Sudetes Foreland:** Szklary 0.4 km W; IV; overgrown spoil tip; soil.

***Rhizina undulata* Fr.**

Kaczawskie Mts.: Bystrzyca 1.4 km SE (WIF-130f); VIII-X; spruce forest (old campfire); charcoal. **Kaczawskie Foothills:** Radomiłowice 0.7 km SW (WIF-11f); VIII; spruce forest (old campfire); charcoal.

***Rhytisma acerinum* (Pers.) Fr.**

Kaczawskie Foothills: Leszczyna 1 km N (near Ciche Szczęście mine); IV; oak-hornbeam forest; *Acer platanoides* leaves, *Acer pseudoplatanus* leaves.

***Sarcoscypha austriaca* (Beck ex Sacc.) Boud.**

Izerskie Foothills: Pilchowice 1 km N (MaF); III; deciduous sapling stand; wood. **Kaczawskie Foothills:** Belczyna 0.8 km S (WIF-123l), Belczyna 0.9 km NW (WIF-13s), Bystrzyca 2.6 km NW (WIF-21k), Belczyna 2 km SE (WIF-124b), Belczyna 0.8 km S (WIF-123l), Leszczyna 1 km N (near Ciche Szczęście mine); II-IV; oak-hornbeam forest, mixed forest, young spruce forest, mixed wood margin; wood.

***Scutellinia barlae* (Boud.) Maire**

Kaczawskie Mts.: Buki Sudeckie Res.; X; *Carici remotae-Fraxinetum*; soil.

***Scutellinia crinita* (Bull.) Lambotte**

Kaczawskie Mts.: Bystrzyca 1.2 km SE (WIF-131d), Buki Sudeckie Res.; X; *Carici remotae-Fraxinetum*, mixed forest; deciduous wood, litter. **Kaczawskie Foothills:** Bystrzyca 2 km NE (WIF-15s), Wleń 2.6 km N (WIF-25a); VI-X; deciduous forest, riparian forest; wood, deciduous wood.

***Scutellinia nigrohirtula* (Svrček) Le Gal**

Kaczawskie Mts.: Bystrzyca (WIF-134); VII; deciduous forest; wood. **Izerskie Foothills:** Wleń 1.6 km NW (WIF-46f); VIII; spruce forest; soil. **Kaczawskie Foothills:** Wleń 2.3 km NE (WIF-25c); VII; mixed forest; wood, soil.

***Scutellinia scutellata* (L.) Lambotte**

Kaczawskie Mts.: Bystrzyca (WIF-134); VII; riparian forest; *Fraxinus* sp. branch. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF-64c – Gniazdo Mt); deciduous forest; wood. **Kaczawskie Foothills:** Bystrzyca 1.8 km N (WIF-16a), Bystrzyca 2.6 km NW (WIF-26f), Przeździec 1.3 km E (WIF-12), Radomiłowice 1.4 km W (WIF-12Ai), Nad Groblą Res.; VII-X; oak forest, oak-hornbeam forest, forest near a stream, lime forest; wood, *Quercus* sp. wood, *Tilia* sp. log.

***Scutellinia subhirtella* Svrček**

Izerskie Foothills: SW od Wlenia (ŁuF – Gniazdo Mt); VII; mixed forest; twigs. **Kaczawskie Foothills:** Przeździec 1.3 km E (near WIF-12); VII; forest near a stream; wood.

***Scutellinia trechispora* (Berk. & Broome) Lambotte**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF-64f – Gniazdo Mt); deciduous forest; soil.

***Thelebolus terrestris* Alb. & Schwein. (= *Byssonectria terrestris* (Alb. & Schwein.) Pfister)**

Kaczawskie Mts.: Bystrzyca 1.4 km SE (WIF-130g), Wleń 1.9 km E (WIF-136a); II-IV; under alder (near a pond), under spruce; litter.

***Tolypocladium ophioglossoides* (Ehrh. ex J.F. Gmel.) Quandt, Kepler & Spatafora**

Karkonosze Mts.: Szklarska Poręba (SzPF-299a); X; spruce forest; sporocarps of *Elaphomyces granulatus* Fr. **Izerskie Foothills:** Wleń 2 km SW (ŁuF – Gniazdo Mt); IX; spruce-pine forest; sporocarps of *Elaphomyces* sp.

***Tricharina praecox* (P. Karst.) Dennis**

Kaczawskie Foothills: Wleń 1.7 km NE (WIF-27h); I; spruce-pine forest (old campfire); charcoal.

***Trichoglossum hirsutum* (Pers.) Boud.**

Kaczawskie Mts.: Buki Sudeckie Res.; IX; *Dentario enneaphylli-Fagetum*; soil.

Verpa conica* (O.F. Müll.) Sw. f. *conica

Izerskie Foothills: Wleń; V; pot with *Saxifraga* sp. plant (purchased in Jelenia Góra); soil.

***Vibrissea flavovirens* (Pers.) Korf & J.R. Dixon**

Kaczawskie Mts.: near Bystrzyca (WIF-134); VII; riparian forest; deciduous wood.

BASIDIOMYCOTA

***Abortiporus biennis* (Bull.) Singer**

Kaczawskie Mts.: Bystrzyca 1.3 km SE (WIF-130c); levee of a forest pond; wood.

***Agaricus augustus* Fr.**

Kaczawskie Mts.: Tarczyn 0.9 km E (WIF-134c); IX; spruce forest; soil. **Kaczawskie Foothills:** Wleń 2 km N (WIF-30f); IX; oak-hornbeam forest; soil.

***Agaricus comtulus* Fr.**

Kaczawskie Mts.: Tarczyn 1.2 km NE (WIF-133f); IX; mixed forest; soil. **Kaczawskie Foothills:** Bełczyna 1.9 km SE (WIF-124g); IX; forest margin; soil.

***Agaricus essettei* Bon**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; roadside of the forest road; soil. **Kaczawskie Foothills:** Górczyca 1.1 km W (WIF-24g); IX; mixed forest; soil.

***Agaricus huijsmanii* Courtec. (= *A. niveolutescens* Huijsman)**

Kaczawskie Mts.: Bystrzyca 1.5 km SE (WIF-130c); VIII; mixed forest; soil.

***Agaricus porphyrhizon* P.D. Orton**

Kaczawskie Mts.: Tarczyn 1.2 km E (WIF-133g); VIII; spruce forest; soil.

***Agaricus semotus* Fr.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; deciduous forest; soil. **Kaczawskie Foothills:** Bełczyna 1.8 km SE (WIF-124a); IX; mixed forest; soil.

***Agaricus sylvaticus* Schaeff.**

Izerskie Foothills: Łupki (near tourist hostel); IX; lawn (under spruce and larch); soil. **Kaczawskie Foothills:** Wleń 1.8 km N (WIF-30c); VIII; mixed forest; soil.

***Agaricus sylvicola* (Vittad.) Peck s.str.**

Izerskie Foothills: Wleń 2.1 km SE (WIF-45a), Wleń 2.3 km NW (WIF-45j); IX-X; deciduous forest, mixed forest; soil.

***Agaricus urinascens* (F.H. Möller & J. Schäff.) Singer (= *A. macrosporus* F.H. Moller & Jul. Schäff.) Pilát)**

Kaczawskie Foothills: Radomiłowice 0.8 W (near WIF-12Ar); X; meadow; soil.

***Agaricus xanthodermus* Genev.**

Kaczawskie Foothills: at the Pilchowice Lake; XI; deciduous forest; soil.

***Agrocybe arvalis* (Fr.) R. Heim & Romagn.**

Kaczawskie Foothills: Wleń 2.5 km NE (near WIF-25c); IX; field; soil.

***Agrocybe dura* (Bolton) Singer**

Kaczawskie Foothills: Bystrzyca 2.4 km NW (near WIF-22); IX; field; soil.

***Agrocybe elatella* (P. Karst.) Vesterh. (= *A. paludosa* (J.E. Lange) Kühner & Romagn)**

Izerskie Foothills: Wleń; VI; lawn (under fir); soil.

***Agrocybe erebia* (Fr.) Singer**

Izerskie Foothills: Wleń 1.3 km NW (WIF-47s), Wleń 1.4 km NW (WIF-46g); IX; deciduous forest, mixed forest; litter, soil. **Kaczawskie Foothills:** Bełczyna 1 km SE (WIF-122f), Wleń 2.3 km N (WIF-25a); X; beech forest, oak-hornbeam forest; wood.

***Agrocybe pediades* (Fr.) Fayod**

Kaczawskie Mts.: Rząśnik 1.5 km W (near WIF-290); IX; meadow; soil. **Izerskie Foothills:** Wleń; V-VII; lawn; soil. **Kaczawskie Foothills:** Bełczyna 1 km S (near WIF-123n); X; grazed meadow; soil.

***Agrocybe praecox* (Pers.) Fayod**

Kaczawskie Mts.: Płoszczyńska 0.5 km W (PiF-250n); V; meadow (log landing); soil. **Kaczawskie Foothills:** Radomiłowice 0.7 km SW (near WIF-12a), Ustronie 2.5 km W (UsF – Piekieleko wilderness); V; forest margin (under oak), levee of a pond; soil.

***Alloclavaria purpurea* (Fr.) Dentinger & D.J. MacLaughlin (= *Clavaria purpurea* O.F. Müll.)**

Izera Mts.: at the Izera river (0.4 km from Torfowiska Rzeki Izery Res.); IX; among mosses; soil.

***Amanita argentea* Huijsman**

Kaczawskie Foothills: Wleń 2.3 km N (WIF-25a); VII; wet oak-hornbeam forest; soil.

***Amanita battarae* (Boud.) Bon**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; soil. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); IX; mixed forest; soil.

***Amanita citrina* (Schaeff.) Pers. f. *alba* (Pers.) Quél.**

Kaczawskie Foothills: Bystrzyca 2.1 km N (WIF-14d); VIII; mixed forest; soil.

Amanita citrina* (Schaeff.) Pers. f. *citrina

Kaczawskie Foothills: Bystrzyca 2.1 km N (WIF-14h), Radomiłowice 1.1 km SW (WIF-11g), Wleń 1.1 km N (WIF-31j); X; deciduous forest, mixed forest; soil.

***Amanita crocea* (Quél.) Singer**

Kaczawskie Mts.: Tarczyn 0.9 NE (WIF-134c); VIII; spruce forest; soil. **Kaczawskie Foothills:** Bełczyna 1.8 km S (WIF-126j), Bystrzyca 2.1 km E (WIF-125d); VII-X; mixed forest, deciduous forest; soil.

***Amanita eliae* Quél.**

Kaczawskie Mts.: Tarczyn 1.2 km E (WIF-133); IX; mixed forest; soil.

Amanita excelsa* (Fr.) Bertill. f. *excelsa

Kaczawskie Mts.: Tarczyn 0.9 km NE (WIF-134c); VI; young spruce forest; soil. **Izerskie Foothills:** Wleń 1.7 km SW (ŁuF – Gniazdo Mt), Wleń 1.9 km SW (ŁuF – Gniazdo Mt), Wleń 2 km SW (ŁuF – Gniazdo Mt); VIII-IX; spruce-beech forest, spruce forest, mixed forest; soil.

***Amanita fulva* (Schaeff.) Fr.**

Karkonosze Mts.: Szklarska Poręba (SzPF-355g); IX; spruce forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.1 km E (WIF-125d); X; deciduous forest; soil.

***Amanita gemmata* (Fr.) Bertill.**

Kaczawskie Foothills: Bełczyna 2 km SE (WIF-125b); X; spruce forest; soil.

***Amanita mairei* Foley**

Kaczawskie Mts.: Bystrzyca 1.4 km SE (WIF-131i); XI; spruce forest; soil.

Amanita muscaria* (L.) Pers. var. *muscaria

Karkonosze Mts.: Szklarska Poręba (Złoty Potok Valley); X; spruce forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.2 km N (WIF-15d); IX-X; mixed sapling stand; soil.

***Amanita muscaria* (L.) Pers. var. *regalis* Fr. (= *A. regalis* (Fr.) Michael)**

Izera Mts.: Świeradów-Zdrój (ŚF-452b), Jagnięcy Jar, between tourist hostel and Torfowiska Doliny Izery Res.; VIII-X; spruce-larch forest, under spruce, spruce forest; soil. **Karkonosze Mts.:** Szklarska Poręba; VIII; mixed forest; soil.

***Amanita pachyvolvata* (Bon) Krieglst.**

Izerskie Foothills: Łupki (near tourist hostel); IX; lawn (under spruce and larch); soil.

***Amanita pantherina* (DC.) Krombh.**

Izerskie Foothills: Wleń 2 km NW (WIF-46d); VIII; coniferous forest; soil.

***Amanita phalloides* (Vaill.) Link var. *alba* Constantin & L.M. Dufour**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; soil. **Kaczawskie Foothills:** Wleń 1.6 km NE (WIF-28k); X; mixed forest; soil.

Amanita phalloides* (Vaill.) Link var. *phalloides

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); IX; oak-hornbeam forest; soil. **Izerskie Foothills:** Wleń 1.3 km SW (ŁuF-58m); VIII; mixed forest; soil. **Kaczawskie Foothills:** Bystrzyca 1.8 km N (WIF-16a), Bystrzyca 2.2 km N (WIF-13d), Wleń 2.6 km N (WIF-25a); VIII-IX; oak-hornbeam forest, deciduous forest; soil. **Sudetes Foreland:** Szklary, Szklary 0.6 km S; IX; oak forest, overgrown meadow; soil.

***Amanita porphyria* Alb. & Schwein.**

Izerskie Foothills: Wleń 1.7 km SW (ŁuF – Gniazdo Mt), Wleń 1.8 km SW (ŁuF – Gniazdo Mt), Wleń 2.1 km NW (WIF-47j); VIII-IX; coniferous forest, mixed forest, spruce forest; soil.

Amanita rubescens* Pers. f. *rubescens

Kaczawskie Foothills: Przeździec 0.8 km SW (WIF-22l), Wleń 2.3 km NE (WIF-25c); VIII-IX; mixed forest, spruce-pine forest; soil.

***Amanita spadicea* Pers.**

Karkonosze Mts.: Karkonosze NP (Czeska Ścieżka); VIII; spruce forest; soil.

***Amanita strobiliformis* (Vittad.) Bertill.**

Kaczawskie Mts.: Góra Milek Res.; VIII; *Cephalanthero-Fagetum*; soil.

Amanita submembranacea* (Bon) Gröger var. *submembranacea

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); IX; oak-hornbeam forest; soil.

Amanita vaginata* (Bull.) Lam. f. *vaginata

Kaczawskie Foothills: Bystrzyca 2.8 km NW (WIF-26i); VII; mixed forest; soil.

***Amphinema byssoides* (Pers.) J. Erikss.**

Sudetes Foreland: Szklary 0.5 km SW; X; overgrown spoil tip; litter.

***Ampulloclitocybe clavipes* (Pers.) Redhead, Lutzoni, Moncalvo & Vilgalys (= *Clitocybe clavipes* (Pers.) P. Kumm.)**

Kaczawskie Foothills: Bystrzyca 1.9 km N (WIF-14h); IX; spruce forest; soil.

***Amylostereum areolatum* (Chaillet ex Fr.) Boidin**

Karkonosze Mts.: Szklarska Poręba (SzPF-354a); spruce forest; *Picea abies* wood. **Izerskie Foothills:** Pokrzywnik 1.2 km S (MaF-176d), Pokrzywnik 1.5 km S (MaF-176c); I; spruce forest; *Picea abies* stump.

***Antrodia crassa* (P. Karst.) Ryvarden**

Kaczawskie Foothills: Bystrzyca 1.9 km N (WIF-18c); I; spruce forest; *Picea abies* log.

***Antrodia serialis* (Fr.) Donk**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); XI; deciduous forest; *Picea abies* log. **Kaczawskie Foothills:** Wleń 2 km N (WIF-29h), Wleń 2.3 km NE (WIF-2c); II, IX; spruce forest; log.

***Antrodiella serpula* (P. Karst.) Spirin & Niemelä (= *A. hoehnelii* (Bres.) Niemelä)**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Wleń 1.2 km SE (WIF-137d); VIII-IX; oak-hornbeam forest, alder forest; *Carpinus betulus* stump, *Alnus* sp. log. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt); XI; deciduous forest, alder forest; ?*Alnus incana* branch.

***Armillaria borealis* Marxm. & Korhonen**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); X; riparian forest; wood. **Izerskie Foothills:** Wleń 1.7 km SW (ŁuF – Gniazdo Mt); X; spruce forest; wood. **Kaczawskie Foothills:** Wleń 2.3 km N (WIF-29b); IX; mixed forest; wood.

***Armillaria lutea* Gillet**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); X; riparian forest; wood. **Kaczawskie Foothills:** Wleń 2.6 km N (WIF-25a); X; deciduous forest; wood.

***Armillaria ostoyae* (Romagn.) Herink**

Izera Mts.: Świeradów-Zdrój (ŚF-238i); spruce forest; wood. **Kaczawskie Foothills:** Wleń 1.4 km N (WIF-31c); X; mixed forest; wood.

***Arrhenia epichysium* (Pers.) Redhead, Lutzoni, Moncalvo & Vilgalys (= *Omphalina epichysium* (Pers.) Quél.)**

Kaczawskie Foothills: Bełczyna 0.7 km S (WIF-123g); X; spruce forest; wood.

***Arrhenia parvivelutina* (Cléménçon & Irlet) Redhead, Lutzoni, Moncalvo & Vilgalys**

Karkonosze Mts.: Karkonosze NP (Kocioł Wielkiego Stawu – near ruins of a tourist hostel); IX; alpine grassland; soil.

***Arrhenia retiruga* (Bull.) Redhead**

Izerskie Foothills: Radomice 0.4 km E; X-XII; limestone quarry; mosses. **Kaczawskie Foothills:** at the Pilchowice Lake; X; basalt quarry; mosses.

***Arrhenia spathulata* (Fr.) Redhead**

Izerskie Foothills: Radomice 0.4 km E; XII; limestone quarry; mosses.

***Asterophora lycoperdoides* (Bull.) Ditmar**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); IX; oak-hornbeam forest; decaying basidiocarps of agaricoid fungi. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; wet spruce forest; decaying basidiocarps of agaricoid fungi. **Kaczawskie Foothills:** Radomiłowice 0.9 km W (WIF-12Ac); VII; under lime tree; decaying basidiocarps of agaricoid fungi.

***Asterophora parasitica* (Bull.) Singer**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; spruce forest; decaying basidiocarps of agaricoid fungi.

***Auricularia auricula-judae* (Bull.) J. Schröt.**

Kaczawskie Mts.: Rząśnik 2.7 km W (WIF-128b); II; mixed forest; *Sambucus nigra* branch. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); II; deciduous forest; *Fraxinus* sp. wood. **Kaczawskie Foothills:** Bełczyna 0.8 km N (WIF-123m), Leszczyna 1 km N (near Ciche Szczęście mine), Przeździec 0.2 km towards Wleń; IV-X; oak-hornbeam forest, deciduous sapling stand, roadside; *Alnus* sp. wood, *Populus tremula* log, deciduous trunk.

***Auriscalpium vulgare* Gray**

Kaczawskie Mts.: Tarczyn 0.9 km NE (WIF-134c); XI; spruce forest; *Picea abies* cone. **Kaczawskie Foothills:** Bystrzyca 2.3 km NW (WIF-15f), Wleń 1.9 km NE (WIF-28d); X-XI; mixed forest, spruce forest; *Picea abies* cone.

***Baeospora myosura* (Fr.) Singer**

Kaczawskie Mts.: Bystrzyca 1.1 km S (WIF-132b); XI; spruce forest; *Picea abies* cone. **Kaczawskie Foothills:** Bełczyna 0.6 km S (WIF-123g); X; spruce forest; *Picea abies* cone.

***Basidioradulum radula* (Fr.) Nobles**

Izerskie Foothills: Wleń 2.8 km N (near WIF-44j); wastelands; *Alnus incana* branch. **Kaczawskie Foothills:** Belczyna 0.6 km S (WIF-123f), Wleń 2.1 km N (WIF-30c); I-II; deciduous forest, deciduous sapling stand; *Salix ?fragilis* branch, deciduous branch.

***Bjerkandera adusta* (Willd.) P. Karst.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; deciduous wood. **Kaczawskie Foothills:** Bystrzyca 1.2 km SE (WIF-126n), near Wąwóz Myśluborski Res.; I, XI; forest, riparian forest; *Fraxinus* sp. wood.

***Bolbitius pluteoides* M.M. Moser**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-59b – Gniazdo Mt); X; riparian forest; wood.

***Bolbitius reticulatus* (Pers.) Ricken f. *aleuriatus* (Fr.) Enderle**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; wood.

***Bolbitius titubans* (Bull.) Fr.**

Kaczawskie Mts.: Rząśnik 2.3 km W (WIF-128g), near Wleń (near WIF-136); V-IX; deciduous forest, log landing; straw, soil. **Izerskie Foothills:** Wleń 2.1 km SW (ŁuF – Gniazdo Mt); IX; mixed forest; soil. **Kaczawskie Foothills:** Bystrzyca 1.7 km N (WIF-16g), Bystrzyca 2.4 km NW (WIF-21l, near WIF-22o), Wleń 2.5 km N (WIF-25a); VII-X; deciduous forest, meadow, field, spruce forest; sawdust, litter, soil.

***Boletopsis leucomelaena* (Pers.) Fayod**

Kaczawskie Foothills: Bystrzyca 2.5 km NW (WIF-21g); VII; spruce forest; soil.

***Boletus calopus* Pers.**

Izera Mts.: Piechowice-Górzyniec 0.5 km W; VIII; mixed forest; soil.

***Boletus edulis* Bull.**

Izera Mts.: at the Izera river (0.4 km S from Torfowiska Doliny Izery Res.); VIII; spruce forest; soil. **Kaczawskie Mts.:** Bystrzyca 0.9 km S (WIF-127a); VIII; riparian forest; soil. **Kaczawskie Foothills:** Wleń 0.9 km NW (WIF-32b); X; oak forest; soil.

Boletus luridiformis* Rostk. var. *luridiformis

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Tarczyn 1.1 km NE (WIF-134b); VII-VIII; oak-hornbeam forest, mixed forest; soil. **Karkonosze Mts.:** Szklarska Poręba (Złoty Potok Valley); X; beech forest; soil. **Izerskie Foothills:** Wleń 1.5 km SW (ŁuF-64 – Gniazdo Mt); spruce forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.1 km N, Bystrzyca 2.4 km NW (WIF-22p), Zalesie 0.6 km N (LuF); VIII-IX; deciduous forest, pine forest; soil.

Boletus luridus* Schaeff. var. *luridus

Kaczawskie Mts.: Nowe Rochowice; VIII; mixed forest; soil. **Karkonosze Mts.:** Szklarska Poręba (SzPF-350g); IX; mixed forest, soil. **Kaczawskie Foothills:** Radomiłowice 0.7 km S (near WIF-11a); VII-IX; log landing; soil.

***Boletus pulverulentus* Opat.**

Izerskie Foothills: Łupki (near tourist hostel); IX-X; lawn (near spruce and apple tree); soil.

***Boletus radicans* Pers.**

Kaczawskie Mts.: Góra Miłek Res.; VIII; *Cephalanthero-Fagetum*; soil.

***Boletus reticulatus* Schaeff.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VII; oak-hornbeam forest; soil. **Kaczawskie Foothills:** Bystrzyca 1.2 km N (WIF-171); IV; beech forest; soil.

***Boletus satanas* Lenz**

Kaczawskie Mts.: Grudno (Bukowa Mt), Góra Miłek Res.; VIII; *Cephalanthero-Fagetum*; soil.

***Byssomerulius corium* (Pers.) Parmasto**

Kaczawskie Mts.: Wleń 1.4 km SE (near WIF-137h), Wleń 1.7 km SE (WIF-137o); I; lime forest, margin of pine sapling stand; *Tilia* sp. branch, *Populus tremula* branch. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); II; riparian forest; *Fraxinus* sp. branch. **Kaczawskie Foothills:** Bystrzyca 1.2 km SE (WIF-126n), near Wąwóz Lipy Res.; I, XI; riparian forest; *Tilia* sp. branch, *Alnus incana* branch.

***Calocera cornea* (Batsch) Fr.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); X; riparian forest; ?*Acer pseudoplatanus* wood. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2 km SW (ŁuF – Gniazdo Mt); II-IX; beech forest, deciduous forest; deciduous wood, *Alnus incana* wood. **Kaczawskie Foothills:** Górczyca 1 km E (WIF-12Ao), Górczyca 1.1 km W (WIF-24g); VII-IX; mixed forest, birch forest; ?*Sorbus aucuparia* wood, *Carpinus betulus* wood.

***Calocera furcata* (Fr.) Quél.**

Kaczawskie Mts.: Wleń 2.5 km E (WIF-134w); XI; mixed forest; *Picea abies* log. **Izerskie Foothills:** Wleń 2.1 km SW (ŁuF – Gniazdo Mt); IX; beech forest; *Picea abies* wood.

***Calocera viscosa* (Pers.) Fr.**

Karkonosze Mts.: Szklarska Poręba (Złoty Potok Valley); XI; spruce forest; wood. **Izerskie Foothills:** Wleń 2 km NW (WIF-46d), Zalesie 0.6 km N; VIII-IX; pine forest; stump, litter. **Kaczawskie Foothills:** Bystrzyca 2.1 km N (WIF-14h), Radomiłowice 1.1 km SW (WIF-11g); VIII-XI; mixed forest; wood, litter.

***Cantharellus amethysteus* (Quél.) Sacc.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1.2 km S (WIF-133b); VIII-X; oak-hornbeam forest, mixed forest; soil. **Kaczawskie Foothills:** Wleń 2 km N (WIF-30f); VIII; under lime tree; soil.

Cantharellus cibarius* Fr. f. *cibarius

Kaczawskie Foothills: Bystrzyca 1.8 km N (WIF-16a); X; oak-hornbeam forest; soil.

***Cantharellus ferruginascens* P.D. Orton**

Kaczawskie Foothills: Wleń 2.1 km N (WIF-30c), Wleń 2.6 km N (WIF-25a); VIII; deciduous forest; soil.

***Cantharellus friesii* Quél.**

Kaczawskie Mts.: Wleń 1.3 km NE (WIF-136b); spruce forest; soil. **Kaczawskie Foothills:** Wleń 1.4 km N (WIF-32c); VIII; mixed forest; soil.

***Cantharellus lutescens* (Pers.) Fr. (= *C. aurora* (Batsch) Kuyper)**

Izerskie Foothills: Zalesie 0.8 km N (LuF); VIII-X; mire (under birch, pine and alder); soil.

***Cantharellus pallens* Pilát**

Kaczawskie Foothills: Wleń 2 km N (WIF-30f); VII; under lime tree; soil.

***Cantharellus tubaeformis* Fr.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-132c); X; spruce forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.1 km N (WIF-14h); IX; mixed forest; soil.

***Carcinomyces polyporina* (D.A. Reid) A.M. Yurkov (= *Tremella polyporina* D.A. Reid)**

Kaczawskie Foothills: Belczyna 0.7 km S (WIF-123g); X; spruce forest; hymenium of *Postia caesia* growing on *Picea abies* stump.

***Ceriporia purpurea* (Fr.) Donk**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); II; riparian forest; ?*Fraxinus* sp. branch.

***Ceriporiopsis mucida* (Pers.) Gilb. & Ryvarden**

Izerskie Foothills: Wleń 2.1 km SW (ŁuF – Gniazdo Mt); XI; mixed forest; deciduous log.

***Chalciporus piperatus* (Bull.) Bataille**

Kaczawskie Mts.: Tarczyn 0.6 km E (Tarczynka Mt., slate quarry); XI; overgrown quarry; soil. **Karkonosze Mts.:** Szklarska Poręba (Złoty Potok Valley); XI; spruce forest; soil. **Izerskie Foothills:** Wleń 1.5 km SW (ŁuF – Gniazdo Mt); VIII; spruce forest; soil. **Kaczawskie Foothills:** Radomiłowice 0.9 km W (WIF-12Ac); X; mixed forest; soil.

***Chlorophyllum rachodes* (Vittad.) Vellinga (= *Macrolepiota rhacodes* Vittadini var. *rhacodes*)**

Kaczawskie Foothills: Bełczyna 0.5 km S (WIF-123), Bystrzyca 2 km W (WIF-126b), Bystrzyca 2 km N (WIF-13i), Wleń 2.5 km N (WIF-25a); IX-X; oak-hornbeam forest, spruce forest; soil.

***Chondrostereum purpureum* (Pers.) Pouzar**

Kaczawskie Foothills: Bystrzyca 2.6 km NW (WIF-26j), near Wąwóz Myśliborski Res. (aside from Myślibórz village); IX, XI; mixed forest, riparian forest; *Alnus incana* wood.

Chroogomphus rutilus* (Schaeff.) O.K. Mill. var. *utilus

Sudeckie Foreland: Szklary 0.5 km SW; VIII-IX; spoil tip afforested with pines; soil.

***Clathrus archeri* (Berk.) Dring**

Kaczawskie Mts.: Rząśnik 2.2 km W (WIF-128k), Rząśnik 2,9 km W (WIF-135f), Taczyn 0.8 km NE (near WIF-134f), Taczyn 0.9 km N (WIF-134fx), Taczyn 1.2 km NE (WIF-133f); VII-X; mixed forest, spruce sapling stand, under oak, forest clearing, forest margin; soil. **Izerskie Foothills:** Wleń 1.3 km NW (WIF-47l), Wleń 2.3 km NW (WIF-45j); VII-IX; mixed forest; soil. **Kaczawskie Foothills:** Bełczyna 1 km S, Bystrzyca 2.1 km E (near WIF-126b), Bystrzyca 2.5 km NW (WIF-22p), Bystrzyca 2,5-2.6 km NW (WIF-21d), Bystrzyca 2.8 km NW (WIF-21c), Górczyca 1.4 km W (WIF-24f), Przeździec 0.8 km S (WIF-22d), Przeździec 0.8 km SE (WIF-21n), Radomiłowice 0.7 km SW (near WIF-12a), Ustronie 2.5 km W (UsF – Piekiełko wilderness), Wleń 1.4 km N (WIF-32a), Wleń 1.6 km NW (WIF-28f), Wleń 1.7 km NE (WIF-27g), Wleń 1.9 km NE (WIF-27f), Wleń 2.3 km NE (WIF-26m); VI-X; levee of the pond, deciduous forest, mixed forest, deciduous forest near a stream, meadow, deciduous sapling stand, pine sapling stand, logging road, spruce forest, deciduous tree clump; soil.

Clavaria argillacea* Pers. var. *argillacea

Karkonosze Mts.: Karkonosze NP (Śnieżka Mt), Karkonosze NP (Przedział Mt, near the tourist trail from Jakuszyce to Hala Szrenicka); IX-X; alpine grassland, spruce forest; soil.

***Clavaria falcata* Pers.**

Kaczawskie Mts.: Bystrzyca 1-1.2 km S (WIF-133a), Nowe Rochowice, Buki Sudeckie Res., Góra Miłek Res.; VIII-X; *Dentario enneaphylli-Fagetum*, *Cephalanthero-Fagetum*, calcareous grassland, ash tree riparian forest; soil. **Kaczawskie Foothills:** Bystrzyca 1.5 km NW (WIF-26b), Wleń 1.5 km NW (WIF-31), Wleń 1.6 km NE (WIF-27d); IX-XI; mixed forest, deciduous forest, ash tree riparian forest; soil.

***Clavaria fragilis* Holmsk**

Kaczawskie Mts.: 1. Nowe Rochowice, Bystrzyca 1.2 km S (WIF-133a); VIII-IX; calcareous grassland, riparian forest; soil.

***Clavaria fumosa* Pers.**

Kaczawskie Mts.: Góra Milek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Clavaria incarnata* Weinm**

Kaczawskie Mts.: Buki Sudeckie Res.; IX; *Dentario enneaphylli-Fagetum*; soil.

***Clavariadelphus pistillaris* (L.) Donk**

Kaczawskie Mts.: Góra Milek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Clavulina cinerea* (Bull.) J. Schröt.**

Kaczawskie Mts.: near Bystrzyca (WIF-134x); X; forest near a stream; soil. **Kaczawskie Foothills:** Bystrzyca 1.2 km SE (WIF-126o), Bystrzyca 1.3 km N (WIF-17g), Bystrzyca 2 km N (WIF-16a), near Nad Groblą Res.; X; deciduous forest, riparian forest, alder forest; soil.

***Clavulina coralloides* (L.) J. Schröt.**

Jelenia Góra Valley: W from Jelenia Góra; VIII; mixed forest; soil. **Izera Mts.:** Jakuszyce 0.8 km NW, Sępik Mt (tourist trail between Szklarska Poręba and Hala Izerska); X; spruce forest, coniferous forest; soil. **Kaczawskie Mts.:** Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1.6 km SE (WIF-131i); VIII-IX; oak-hornbeam forest, mixed forest; soil. **Izerskie Foothills:** Wleń 1.4 km NW (WIF-46g), Wleń 2.4 km W (ŁuF-63b – Gniazdo Mt), Wleń 2.8 km N (WIF-44j); X-XI; oak-hornbeam forest, spruce forest, mixed forest; soil. **Kaczawskie Foothills:** Bełczyna 1.1 km E (near WIF-122), Bełczyna 1.7 km SE (near WIF-124a), Bystrzyca 2 km S (WIF-126w), Bystrzyca 1.7 km N (WIF-16g), Tarczyn 1.1 km E (WIF-133f), Wleń 2.6 km N (WIF-25a); VII-X; mixed forest, deciduous forest, spruce forest; soil.

***Clavulina rugosa* (Bull.) J. Schröt.**

Kaczawskie Mts.: Bystrzyca 0.8 km SE (WIF-127a), Bystrzyca 1.5 km SW (WIF-128i), Tarczyn 0.6 km E, Tarczyn 0.9 km E (WIF-134c), Tarczyn 1.1 km E (WIF-133f), Góra Milek Res.; IX-XI; *Cephalanthero-Fagetum*, coniferous forest, mixed forest, oak-hornbeam forest, spruce forest; soil. **Izera Mts.:** Jakuszyce 1.2 km NW; X; spruce forest; soil. **Izerskie Foothills:** Marczów 2 km SW (WIF-44d), Wleń 1.5 km NW (WIF-46k), Wleń 1.7 km SW (ŁuF – G. Gniado); VIII-XI; spruce forest, spruce-pine forest; soil. **Kaczawskie Foothills:** Bełczyna 0.6 km S (WIF-123g), Bełczyna 1 km SE (WIF-122l), Bełczyna 1 km E (near WIF-122), Bystrzyca 2 km W (WIF-126p), Bystrzyca 2.5 km NW (WIF-21l), Wleń 2.2 km N (WIF-29c), Wleń 2.6 km N (WIF-25a); VIII-X; mixed forest, coniferous forest, deciduous forest, spruce forest; soil.

***Clavulinopsis corniculata* (Schaeff.) Corner**

Kaczawskie Mts.: Buki Sudeckie Res.; VIII-IX; *Dentario enneaphylli-Fagetum*; soil.

***Clavulinopsis fusiformis* (Sowerby) Corner**

Kaczawskie Mts.: Buki Sudeckie Res.; IX; *Dentario enneaphylli-Fagetum*; soil.

***Clavulinopsis helvola* (Pers.) Corner**

Kaczawskie Mts.: Buki Sudeckie Res., Góra Milek Res., Milek Mt; IX-X; *Dentario enneaphylli-Fagetum*, meadow; soil. **Kaczawskie Foothills:** Bełczyna 1,1-1.2 km SE (WIF-123m), Radomiłowice 0.8 km W (WIF-12Ar), Sobota 1.5 km S; X-XI; meadow, pasture; soil.

***Clavulinopsis laeticolor* (Berk. & M.A. Curtis) R.H. Petersen**

Kaczawskie Mts.: Góra Milek Res., Nowe Rochowice; VIII-IX; *Cephalanthero-Fagetum*; soil. **Kaczawskie Foothills:** Wleń 1.6 km NE (WIF-27d); IX; deciduous forest; soil.

***Climacocystis borealis* (Fr.) Kotl. & Pouzar**

Karkonosze Mts.: Szklarska Poręba-Huta (SzPF-173a); IX; spruce forest; *Picea abies* stump. **Izerskie Foothills:** Pilchowice 1.1 km S (MaF – Dziki Valley); X; mixed forest; *Picea abies* log.

***Clitocybe agrestis* Harmaja**

Kaczawskie Foothills: Radomiłowice 0.8 km W (near WIF-12Ar); X; wet meadow; soil.

***Clitocybe albofragrans* (Harmaja) Kuyper**

Izerskie Foothills: Radomice (limestone quarry); IX; overgrown quarry; soil.

***Clitocybe diatrete* (Fr.) P. Kumm.**

Izerskie Foothills: Radomice (limestone quarry); XI; overgrown quarry; soil.

Clitocybe nebularis* (Barsch) P. Kumm. var. *nebularis

Izerskie Foothills: Wleń 2.1 km N (WIF-44b); X; deciduous forest; soil. **Kaczawskie Foothills:** Bełczyna 0.7 km S (WIF-123h), Bystrzyca 1.5 km E (WIF-126g), Górczyca 1 km E (WIF-12Ao), Przeździec 1 km N (WIF-24t); X-XI; birch forest, deciduous forest, mixed forest, spruce forest; soil.

Clitocybe odora* (Bull.) P. Kumm. var. *odora

Kaczawskie Mts.: Tarczyn 0.6 km E (Tarczynka Mt); IX; overgrown slate excavation; soil. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; spruce forest; litter. **Kaczawskie Foothills:** Radomiłowice 0.7 km S (near WIF-11a); VII; forest margin (log landing); litter.

***Clitocybe vibecina* (Fr.) Quél.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); XI; deciduous forest; soil.

***Clitocybula platyphylla* (Pers.) Malençon & Bertault (= *Megacollybia platyphylla* (Pers.) Kotl. & Pouzar)**

Izerskie Foothills: Wleń 1.8 km SW (ŁuF – Gniazdo Mt), Wleń 2 km SW (ŁuF – Gniazdo Mt); VII-VIII; spruce-pine forest, mixed forest; wood, stump. **Kaczawskie Foothills:** Bystrzyca 1.9 km N (WIF-14j), Bystrzyca 2.5 km NE (WIF-15h); VI-VII; deciduous forest, spruce forest; soil.

***Clitopilus cystidiatus* Hauskn. & Noordel.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; blackthorn shrubs at forest margin; soil.

***Clitopilus pinsitus* (Fr.) Joss.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); IX; riparian forest; wood.

***Clitopilus scyphoides* (Fr.) Singer f. *reductus* Noordel.**

Kaczawskie Foothills: Bystrzyca 2.1 km NW (WIF-26c); X; deciduous forest; wood.

Clitopilus scyphoides* (Fr.) Singer f. *scyphoides

Sudetes Foreland: Szklary 0.5 km SW; IX; overgrown outcrop; litter.

***Collybia cirrata* (Schumach.) Quél.**

Kaczawskie Foothills: Sobota 1.5 km S; X; pasture; decaying basidiocarps of agaricoid fungi.

***Collybia cookei* (Bres.) J.D. Arnold**

Kaczawskie Foothills: Rząśnik 1.3 km NW (WIF-125); X; mixed forest; decaying basidiocarps of agaricoid fungi.

***Collybia tuberosa* (Bull.) P. Kumm.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; decaying basidiocarps of agaricoid fungi.

***Coltricia perennis* (L.) Murrill**

Karkonosze Mts.: Szrenica Mt (NE slope, ca. 920 m a.s.l.); VIII; spruce forest; soil.

Conocybe albipes* (G.H. Otth) Hauskn. var. *albipes

Izerskie Foothills: Wleń; VII; lawn; soil. **Kaczawskie Foothills:** Wleń 2,9 km N (near WIF-25b); VII; meadow; soil.

***Conocybe antipus* (Lasch) Fayod**

Kaczawskie Foothills: Bystrzyca 1.6 km N (WIF-16b); VIII; forest margin; soil.

***Conocybe brunneidisca* (Murrill) Hauskn.**

Kaczawskie Foothills: Bystrzyca 0.5 km N (WIF-20l), Sobota 1.5 km S; X-XI; overgrown pasture, basalt quarry; soil.

***Conocybe echinata* (Velen.) Singer**

Kaczawskie Foothills: Sobota 1.5 km S; XI; overgrown pasture; soil.

Conocybe juniana* (Velen.) Hauskn. & Svrček var. *juniana

Izerskie Foothills: Wleń; VII; lawn; soil.

***Conocybe juniana* (Velen.) Hauskn. & Svrček var. *sordescens* (P.D. Orton) Hauskn.**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry); X; calcareous grassland; soil.

***Conocybe juniana* (Velen.) Hauskn. & Svrček var. *subsejuncta* Hauskn.**

Izerskie Foothills: Wleń; VI-IX; lawn; soil. **Kaczawskie Foothills:** Wleń 2.6 km N (WIF-25a); VII; deciduous forest; litter.

***Conocybe microrrhiza* Hauskn. var. *tetraspora* (Singer & Hauskn.) Hauskn.**

Izerskie Foothills: Wleń; VI; lawn; soil.

Conocybe microspora* (Velen.) Dennis var. *microspora

Jelenia Góra Valley: Jelenia Góra (Jana Pawła II St. – near TESCO market); IX; lawn; soil.

Conocybe moseri* Watling var. *moseri

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); spruce forest; soil. **Kaczawskie Foothills:** Sobota 1.5 km S, Wleń 2.4 km N (WIF-29b); IX-XI; pasture, mixed forest; soil.

***Conocybe moseri* Watling var. *robustior* Hauskn. & H.-J. Hübner**

Kaczawskie Foothills: Bystrzyca 2.4 km NW (near WIF-22); IX; field; soil.

***Conocybe pilosella* (Pers.) Kühner**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); IX; oak-hornbeam forest; soil.

***Conocybe pubescens* (Gillet) Kühner**

Kaczawskie Mts.: Bystrzyca 1.1 km S (WIF-131g); XI; spruce forest; soil. **Karkonosze Mts.:** Szklarska Poręba (SzPF-325g); IX; roadside; soil.

***Conocybe pulchella* (Velen.) Hauskn. & Svrček (= *C. pseudopilosella* (Kühner) ex Kühner et Watling)**

Kaczawskie Foothills: Wleń 2.6 km N (WIF-25a); VIII; deciduous forest margin; soil.

***Conocybe rickeniana* P.D. Orton**

Kaczawskie Mts.: Góra Miłek Res.; X; deciduous forest; soil. **Izerskie Foothills:** Wleń 1.7 km SW (ŁuF – Gniazdo Mt); mixed forest; soil.

Conocybe semiglobata* Kühner & Watling var. *semiglobata

Jelenia Góra Valley: Jelenia Góra (Jana Pawła II St. – near TESCO market); IX; lawn; soil. **Izerskie Foothills:** Wleń 2 km SW (ŁuF – Gniazdo Mt); VII; spruce forest; soil. **Kaczawskie Foothills:** Bełczyna 1.1 km SE (WIF-123p), Bystrzyca 2.5 km NW (WIF-15h); mixed forest; soil.

***Conocybe singeriana* Hauskn.**

Kaczawskie Foothills: Sobota 1.5 km S; XI; overgrown pasture; soil.

***Conocybe subovalis* Kühner & Watling**

Kaczawskie Foothills: Sobota 1.5 km S; XI; pasture; soil.

***Conocybe subpallida* Enderle**

Kaczawskie Foothills: Bystrzyca 0.5 km N (WIF-20l); basalt quarry; soil.

***Conocybe subpubescens* P.D. Orton**

Izera Mts.: Sępik Mt (N slope); IX; spruce forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.6 km N (WIF-22), Przeździec 0.8 km SW (WIF-22l), Przeździec 1.3 km E (WIF-12l); IV-VI; mixed forest, alder riparian forest; soil.

***Coprinellus angulatus* (Peck) Redhead, Vilgalys & Moncalvo (= *Coprinus angulatus* Peck non J.E. Lange)**

Kaczawskie Foothills: Bełczyna 2.2 km S (WIF-124b); X; mixed forest (old campfire); charcoal.

Coprinellus callinus* (M. Lange & A.H. Sm.) Vilgalys, Hopple & Jacq. Johnson var. *callinus

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt); X; under spruce; litter. **Kaczawskie Foothills:** Bystrzyca 2.2 km N (WIF-15a), Wleń; VIII-IX; lawn, ash tree sapling stand; soil.

***Coprinellus disseminatus* (Pers.) J.E. Lange (= *Coprinus disseminatus* (Pers.) Quél.)**

Izerskie Foothills: Wleń, Wleń (Południowe Es.), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VII; deciduous forest, lawn; wood. **Kaczawskie Foothills:** Bystrzyca 2.5 km NW (WIF-15h), Leszczyna 1 km N (near Ciche Szczęście mine); IV-IX; deciduous forest, oak-hornbeam forest; wood.

***Coprinellus domesticus* (Bolton) Vilgalys, Hopple & Jacq. Johnson (= *Coprinus domesticus* (Bolton) Gray)**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Wojcieszów 2.7 km SE (near Miłek Mt); IV-VII; mixed forest, meadow; wood. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest; wood. **Kaczawskie Foothills:** Bełczyna 1.8 km S (WIF-126j), Bystrzyca 1.2 km SE (WIF-126o), Górczyca 1.1 km W (WIF-24g), Leszczyna 1 km N (near Ciche Szczęście mine), Wleń 2.3 km N (WIF-25a); IV-X; mixed forest, alder forest, oak-hornbeam forest; wood, litter.

***Coprinellus flocculosus* (DC.) Vilgalys, Hopple & Jacq. Johnson (= *Coprinus flocculosus* (DC.) Fr.)**

Kaczawskie Foothills: Bystrzyca 1.6 km N (WIF-16b); IX; forest; straw.

***Coprinellus heterothrix* (Kühner) Redhead, Vilgalys & Moncalvo**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; deciduous forest; soil. **Kaczawskie Foothills:** Belczyna 1.8 km SE (WIF-124a); VIII; oak-hornbeam forest margin; soil.

***Coprinellus impatiens* (Fr.) J.E. Lange (= *Coprinus impariens* (Fr.) Quél.)**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2.1 km SE (WIF-45a); X; deciduous forest; litter, soil. **Kaczawskie Foothills:** Wleń 2.5 km N (WIF-25a), Bystrzyca 2 km NW (WIF-15s), Bystrzyca 2 km N (WIF-16a), rez. Wąwóz Myśluborski; X-XI; oak-hornbeam forest, ash tree riparian forest, deciduous forest; litter, wood.

***Coprinellus marculentus* (Britzelm.) Redhead, Vilgalys & Moncalvo**

Kaczawskie Foothills: Bystrzyca 1.6 km N (WIF-16b); VII; forest margin; straw, manure.

***Coprinellus micaceus* (Bull.) Vilgalys, Hopple & Jacq. Johnson (= *Coprinus micaceus* (Bull.) Fr.)**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a), Bystrzyca 1.2 km S (WIF-133a), Bystrzyca 1.3 km SE (WIF-130b), Bystrzyca 1.4 km SE (WIF-130c), Bystrzyca 1.4 km SW (WIF-128b), Rząśnik 1.5 km W (WIF-290m), Tarczyn 1.4 km NE (WIF-133b), Wleń 0.8 km NE (WIF-136a), Wleń 1.3 km E (WIF-136b); IV-X; riparian forest, mixed forest, spruce forest, oak forest; wood. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.5 km NW (WIF-46g), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km NW (WIF-46k), Wleń 1.7 km SW (ŁuF – Gniazdo Mt), Wleń 1.8 km SW (ŁuF – Gniazdo Mt), Wleń 2.1 km NE (WIF-45a), Wleń 2.8 km N (WIF-44j); V-XI; deciduous forest, riparian forest, oak-hornbeam forest; wood. **Kaczawskie Foothills:** Belczyna 0.5 km SW (WIF-125b), Belczyna 0.7 km S (near WIF-123g), Belczyna 1.2 km SE (WIF-123n), Belczyna 2 km SE (WIF-123n), Bystrzyca 1.2 km SE (WIF-126n, 131d), Bystrzyca 1.4 km N (WIF-19b), Bystrzyca 1.8 km N (WIF-13g), Bystrzyca 2 km N (WIF-16a), Bystrzyca 2.2 km N (WIF-19b), Bystrzyca 2.5 km NW (WIF-15h, 22p), Górczyca 1 km E (WIF-12Ao), Górczyca 1.5 km W (WIF-24d), Radomiłowice 0.7 km SW (WIF-11c), Radomiłowice 1 km NW (WIF-12a), Sichów 2 km S (valley of Gajka river), Sobota 1.7 km SW (WIF-24Ab), Sobota 1.8 km SW (WIF-24Ad), Wleń 1.4 km N (WIF-31b), Wleń 1.9 km N (WIF-29d, 30d), Wleń 1.9 km NE (WIF-27d), Wleń 2.5 km N (WIF-25a), Wleń 2.7 km NE (WIF-22/26, WIF-25c), Wleń 3 km NE (WIF-22r); IV-XI; oak-hornbeam forest, riparian forest, spruce forest, pine forest, deciduous forest, mixed forest, meadow at the forest margin, between spruce forest and deciduous forest, oak forest; wood, *Fraxinus* sp. stump, *Populus tremula* trunk, litter, soil.

***Coprinellus pyrrhantes* (Romagn.) Redhead, Vilgalys & Moncalvo**

Kaczawskie Foothills: Leszczyna 1 km N (near Ciche Szczęście mine); IV; oak-hornbeam forest; wood.

***Coprinellus radians* (Desm.) Vilgalys, Hopple & Jacq. Johnson (= *Coprinus radians* (Desmaz.) Fr.)**

Kaczawskie Foothills: Wleń 3 km NE (WIF-22p); IX; deciduous forest; soil.

***Coprinellus sclerocystidiosus* (M. Lange & A.H. Sm.) Vilgalys, Hopple & Jacq. Johnson**

Kaczawskie Foothills: Bystrzyca 0.9 km E (near WIF-126); IX; meadow; soil.

***Coprinellus silvaticus* (Peck) Gminder (= *Coprinus silvaticus* Peck)**

Izerskie Foothills: Wleń; IV-V; garden; buried wood.

***Coprinellus xanthothrix* (Romagn.) Vilgalys, Hopple & Jacq. Johnson (= *Coprinus xanthothrix* Romagn.)**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Grudno, Wleń 1 km E (WIF-137), Płoszczynka 0.5 km W (WIF-250n), Tarczyn 1.2 km E (WIF-133g), Góra Miłek Res., Buki Sudeckie Res.; V-IX; *Cephalanthero-Fagetum*, *Dentario enneaphylli-Fagetum*, beech forest, oak-hornbeam forest, spruce forest, shelterbelt by the road, meadow; litter. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.4 km NW (WIF-47k), Wleń 1.6 km SW (ŁuF-64), Wleń 1.6 km SW, Wleń 2.7 km NW (WIF-44k); VII-IX; deciduous forest, riparian forest; litter, wood. **Kaczawskie Foothills:** Belczyna 0.6 km NW (WIF-13n), Belczyna 1.3 km SE (WIF-126k), Bystrzyca 1.6 km N (WIF-16h), Bystrzyca 2.2 km N (WIF-15c), Bystrzyca 2.5 km NW (WIF-15g, 15h), Bystrzyca 2.6 km NW (WIF-26i), Górczyca 0.8 km E (WIF-12Ao), Górczyca 1.1 km SW (WIF-24g), Górczyca 1.4 km SW (WIF-24h), Radomiłowice 0.5 km N (WIF-11), Radomiłowice 1.5 km N (WIF-14), Wleń 1.6 km NE (WIF-27k), Wleń 1.7 km NE (WIF-28f), Wleń 2.1 km N (WIF-30c), Wleń 2.6 km N (WIF-30b), Wleń 3 km NE (WIF-22p); V-X; deciduous forest, mixed forest, riparian forest, oak-hornbeam forest, roadside; litter, soil.

***Coprinopsis acuminata* (Romagn.) Redhead, Vilgalys & Moncalvo**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Wleń 1.2 km NE (WIF-136b); IX; oak-hornbeam forest, roadside; wood, soil. **Kaczawskie Foothills:** Wleń 2.3 km N (WIF-25a); IX; oak-hornbeam forest; soil.

***Coprinopsis annulopora* (Enderle) P. Specht & H. Schubert**

Izerskie Foothills: Wleń; VI; composter; compost.

***Coprinopsis atramentaria* (Bull.) Redhead, Vilgalys & Moncalvo (= *Coprinus atramentarius* (Bull.) Fr.)**

Kaczawskie Mts.: Rząśnik 2.1 km W (WIF-128d), Wleń 1.1 km E (WIF-136f), Rząśnik 1.4 km SW (WIF-29a); IX-X; forest roadside, deciduous forest; wood, soil. **Izerskie Foothills:** Wleń, Wleń 1.4 km SW (ŁuF-64c); VI-IX; riparian forest, lawn; soil. **Kaczawskie Foothills:** Belczyna 1.2 km E (WIF-122a), Belczyna 2 km SE (WIF-125b), Bystrzyca 1.2 km SE (WIF-126o); X; between spruce forest and birch forest, alder forest; soil, wood.

***Coprinopsis candidata* (Uljé) Noordel.**

Izerskie Foothills: Wleń, 1.6 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest; soil. **Kaczawskie Foothills:** Bystrzyca 1.6 km N (WIF-16b), Przeździec 0.5 km SW (WIF-22f); VII-IX; mixed sapling stand, forest margin; soil.

***Coprinopsis cinerea* (Schaeff.) Redhead, Vilgalys & Moncalvo (= *Coprinus cinereus* (Schaeff.) Gray)**

Kaczawskie Foothills: Sobota 1 km S; VIII; field, mixed sapling stand; manure.

***Coprinopsis cordispora* (T. Gibbs.) Gminder (= *Coprinus cordisporus* Gibbs)**

Karkonosze Mts.: Szklarska Poręba; VIII; mixed forest; *Equus caballus* dung. **Kaczawskie Foothills:** Bystrzyca 1.6 km N (WIF-16b); VII; forest margin; straw, manure.

***Coprinopsis cortinata* (J.E. Lange) Gminder (= *Coprinus cortinatus* J.E. Lange)**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; logging road; soil.

***Coprinopsis ellisii* (P.D. Orton) Redhead, Vilgalys & Moncalvo**

Kaczawskie Foothills: Bystrzyca 2.5 km NW (WIF-15h); VII; deciduous forest; litter.

***Coprinopsis fusispora* L. Nagy, Vágvölgyi & Papp**

Kaczawskie Mts.: Rząśnik 2 km NW (WIF-131f), Buki Sudeckie Res.; X-XI; log landing, *Carici remotae-Fraxinetum*; soil. **Kaczawskie Foothills:** Leszczyna 1 km N (near Ciche Szczęście mine); IV; oak-hornbeam forest; soil.

***Coprinopsis jonesii* (Peck) Redhead, Vilgalys & Moncalvo**

Kaczawskie Foothills: Bystrzyca 2.2 km N (WIF-15c), Leszczyna 1 km N (near Ciche Szczęście mine); IV-IX; oak-hornbeam forest; wood, litter.

***Coprinopsis kubickae* (Pilát & Svrček) Redhead, Vilgalys & Moncalvo**

Izerskie Foothills: Wleń 1.4 km SW (nad Bobrem); VII; field road; decaying grasses.

***Coprinopsis lagopus* (Fr.) Redhead, Vilgalys & Moncalvo var. *lagopus* (= *Coprinus lagopus* (Fr.) Fr.)**

Kaczawskie Mts.: Bystrzyca 1.1 km S (WIF-131g); X; log landing; soil, woodchips. **Izerskie Foothills:** Wleń 1.3 km NW (WIF-47s); IX; log landing; soil, woodchips. **Kaczawskie Foothills:** Bystrzyca 2.2 km N (WIF-15a, 15c), Przeździec 0.8 km SW (WIF-22l), Radomiłowice 1.1 km SW (WIF-11g); VII-IX; oak-hornbeam forest, mixed forest, log landing, logging road; wood, litter, soil.

***Coprinopsis lagopus* (Fr.) Redhead, Vilgalys & Moncalvo var. *vacillans* Uljé**

Izerskie Foothills: Wleń; VIII; lawn; soil.

***Coprinopsis laanii* (Kits van Wav.) Redhead, Vilgalys & Moncalvo**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest; ?coniferous log.

***Coprinopsis macrocephala* (Berk.) Redhead, Vilgalys & Moncalvo (= *Coprinus macrocephalus* (Berk.) Berk.)**

Kaczawskie Mts.: Bystrzyca 0.7 km SE (near WIF-127a); IX; pasture; dung, hay. **Kaczawskie Foothills:** Bystrzyca 1 km NW; XII; field; manure.

***Coprinopsis marcescibilis* (Britzelm.) Örstadius & E. Larss. (= *Psathyrella marcescibilis* (Britzelm) Singer)**

Kaczawskie Mts.: Rząśnik 1.5 km W (near WIF-290); IV; abandoned farm; soil. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest; litter. **Kaczawskie Foothills:** Wleń 1.4 km NW (near WIF-32); IV; wet meadow; soil.

***Coprinopsis picacea* (Bull.) Redhead, Vilgalys & Moncalvo (= *Coprinus picaceus* (Bull.) Gray)**

Kaczawskie Mts.: between Nowe Rochowice and Grudno; XI; beech forest; soil.

***Coprinopsis radiata* (Bolton) Redhead, Vilgalys & Moncalvo (= *Coprinus radiatus* (Bolt.) Pers.)**

Kaczawskie Mts.: Bystrzyca 1.1 km S (WIF-131g); X; pasture; straw. **Izerskie Foothills:** Wleń (Południowe Es.); VI; composter; compost. **Kaczawskie Foothills:** Bystrzyca 1.6 km N (WIF-16b), Przeździec 0.5 km SW (WIF-22f); VII; forest margin, mixed sapling stand; manure, ?*Sus scrofa* dung, straw.

***Coprinopsis romagnesianae* (Singer) Redhead, Vilgalys & Moncalvo**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-59b – Gniazdo Mt); X-XI; riparian forest, forest roadside; soil.

***Coprinopsis saccharomyces* (P.D. Orton) P. Roux & Guy Garcia**

Izera Mts.: Jagnięcy Jar; IX; spruce-larch forest; soil.

***Coprinopsis semitalis* (P.D. Orton) Redhead, Vilgalys & Moncalvo**

Karkonosze Mts.: Szklarska Poręba (Złoty Potok Valley); XI; beech forest (helocrene); soil.

***Coprinus comatus* (O.F. Müll.) Pers.**

Kaczawskie Mts.: Bystrzyca 1.1 km S (WIF-131g), Góra Milek Res.; IX-X; *Cephalanthero-Fagetum*, log landing; soil. **Kaczawskie Foothills:** Radomiłowice 0.7 km SW (WIF-11a); IX; mixed wood margin; soil.

***Coriolopsis gallica* (Fr.) Ryvarden**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.4 km NW (WIF-46a); VIII-XI; deciduous forest, riparian forest; *Fraxinus* sp. branch and log. **Kaczawskie Foothills:** Bystrzyca 2.1 km NW (WIF-26c); XII; deciduous forest; broken *Fraxinus* sp..

***Corticium roseum* Pers.**

Kaczawskie Mts.: Wleń 1.4 km SE (WIF-137h); I; margin of pine sapling stand; *Populus tremula* branch.

***Cortinarius acutus* (Pers.) Fr.**

Izera Mts.: Świeradów-Zdrój (ŚF-429d); wet spruce forest; soil. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Zalesie 0.8 km NE (LuF-199g); VIII-X; wet spruce forest, mire (under birch, pine, spruce and alder); soil, *Sphagnum* sp.

***Cortinarius albovariegatus* (Velen.) Melot**

Izerskie Foothills: Zalesie 0.8 km NE (LuF-200h); VIII; mire (under birch, pine, spruce and alder); *Sphagnum* sp.

***Cortinarius alboviolaceus* (Pers.) Fr.**

Izerskie Foothills: Radomice (near limestone quarry); IX; overgrown spoil tip; soil. **Kaczawskie Foothills:** Przeździec 0.8 km SW (WIF-221); IX; mixed forest; soil.

***Cortinarius alnetorum* (Velen.) M.M. Moser**

Izerskie Foothills: Wleń 2.1 km SW (ŁuF – Gniazdo Mt); X; alder-spruce forest; soil.

***Cortinarius anomalus* (Fr.) Fr.**

Izera Mts.: Świeradów-Zdrój (ŚF-238k); mixed forest; soil. **Karkonosze Mts.:** Szklarska Poręba (SzPF-325g); IX; mixed forest; soil. **Kaczawskie Foothills:** Bełczyna 1 km E (near WIF-122b, 122s); X; mixed tree clump; soil.

***Cortinarius bolaris* (Pers.) Fr.**

Izera Mts.: Pichowice 2 km W; VIII; spruce forest; soil. **Kaczawskie Mts.:** Bystrzyca 1.2 km S (WIF-133c), Bystrzyca 1.3 km SE (WIF-131i); VIII-IX; under beech, beech forest; soil.

***Cortinarius caperatus* (Pers.) Fr. (= *Rozites caperatus* (Pers.) P. Karst.)**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; soil.

***Cortinarius cinnamomeus* (L.) Gray**

Kaczawskie Foothills: Wleń 1.4 km N (WIF-32c); XI; mixed forest; soil.

***Cortinarius conicus* (Velen.) Rob. Henry**

Kaczawskie Foothills: Wleń 2.6 km N (WIF-25a); VIII; deciduous forest; soil.

Cortinarius croceus* (Schaeff.) Gray subsp. *croceus

Kaczawskie Foothills: Bełczyna 0.7 km S (WIF-123h); X; spruce forest; soil.

***Cortinarius delibutus* Fr.**

Karkonosze Mts.: Szklarska Poręba (SzPF-361a); X; spruce forest; soil. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt); X; wet spruce forest; soil.

Cortinarius diasemospermus* Lamoure var. *diasemospermus

Kaczawskie Foothills: Bystrzyca 2 km S (WIF-126w); IX; deciduous forest margin; soil.

***Cortinarius flexipes* (Pers.) Fr. var. *flabellus* (Fr.) H. Lindstr. & Melot**

Izerskie Foothills: Wleń 1.3 km SW (ŁuF – Gniazdo Mt); IX; oak-hornbeam forest; soil.

Cortinarius flexipes* (Pers.) Fr. var. *flexipes

Izerskie Foothills: Wleń 1.5-1.6 km SW (ŁuF – Gniazdo Mt), Zalesie 0.8 km N (LuF-200h), Zalesie 0.8 km NE (LuF-199g); VIII-X; spruce forest, mire (under birch, pine, spruce and alder), dry mire margin; soil, *Sphagnum* sp.

***Cortinarius helvelloides* (Fr.) Fr.**

Kaczawskie Mts.: Bystrzyca 1 km SE (WIF-132a); X; riparian forest; soil. **Izerskie Foothills:** Wleń 2.8 km N (WIF-44j); X; oak-hornbeam forest; soil.

***Cortinarius hinnuleus* Fr. s.str.**

Izerskie Foothills: Gryfów Śląski 5 km NE, X; mixed forest, soil. **Kaczawskie Foothills:** Belczyna 1 km E (near WIF-122a), Bystrzyca 1.5 km NW (WIF-26b); IX-X; mixed forest, deciduous tree clump; soil.

Cortinarius huronensis* Ammirati & A.H. Sm. var. *huronensis

Izerskie Foothills: Zalesie 0.8 km NE (LuF-199g); X; mire (under birch, pine, spruce and alder); *Sphagnum* sp.

***Cortinarius huronensis* (Ammirati) & A.H. Sm. var. *olivaceus* Ammirati & A.H. Sm. (= *C. palustris* (M.M. Moser) M.M. Moser)**

Izerskie Foothills: Zalesie 0.8 km NE (LuF-199g); IX; mire (under birch, pine, spruce and alder); *Sphagnum* sp.

***Cortinarius lebretonii* Quél.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; spruce forest; soil.

***Cortinarius limonius* (Fr.) Fr.**

Izerskie Foothills: Zalesie 0.8 km NE (LuF-199g); VIII; mire (under birch, pine, spruce and alder); *Sphagnum* sp.

***Cortinarius malicorius* Fr.**

Izerskie Foothills: Wleń 1.6-1.7 km SW (ŁuF – Gniazdo Mt); VIII-IX; spruce forest; soil.

***Cortinarius mucifluoides* Rob. Henry**

Kaczawskie Foothills: Bystrzyca 1.2 km N (WIF-17l); VIII; beech forest; soil.

***Cortinarius phoeniceus* (Vent.) Maire**

Kaczawskie Foothills: Bystrzyca 1.9 km N (WIF-14i), Wleń 1.4 km N (WIF-31c); VIII; mixed forest, spruce forest; soil.

***Cortinarius pholideus* (Fr.) Fr.**

Kaczawskie Foothills: Bystrzyca 2.1 km N (WIF-14h); IX; mixed forest; soil.

***Cortinarius sanguineus* (Wulfen) Fr.**

Izerskie Foothills: Wleń 2-2.4 km SW (ŁuF – Gniazdo Mt); X; spruce forest; soil.

***Cortinarius saturninus* (Fr.) Fr.**

Izerskie Foothills: Radomice (limestone quarry); X; overgrown quarry; soil.

***Cortinarius semisanguineus* (Fr.) Fr.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; spruce forest; soil. **Kaczawskie Foothills:** Bystrzyca 1.9 km N (WIF-14i); VIII; spruce forest; soil.

***Cortinarius sodagnitus* Rob. Henry**

Kaczawskie Mts.: Góra Milek Res., Grudno; IX; *Cephalanthero-Fagetum*; soil.

***Cortinarius sordidus* (Velen.) Henry ss. Soop**

Kaczawskie Foothills: Wleń 1.9 km NE (WIF-27d); X; deciduous forest; soil.

***Cortinarius stillatitius* Fr.**

Izerskie Foothills: Zalesie 0.8 km NE (LuF-199g), Zalesie 0.8 km N (LuF-200h); VIII-X; mire (under birch, pine, spruce and alder); *Sphagnum* sp.

***Cortinarius torvus* (Fr.) Fr.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.1 km N (WIF-14); mixed forest; soil.

***Cortinarius trivialis* J.E. Lange s.l.**

Kaczawskie Foothills: Wąwóz Lipy Res.; XI; oak forest; soil.

***Cortinarius tubarius* Ammirati & A.H. Sm.**

Izerskie Foothills: Zalesie 0.8 km NE (LuF-199g); VIII; mire (under birch, pine, spruce and alder); *Sphagnum* sp.

***Cortinarius uraceus* Fr.**

Izera Mts.: at the Izera river (0.4 km S from Torfowiska Doliny Izery Res. towards Jagniątków); X; spruce forest; soil.

***Craterellus cornucopioides* (L.) Pers.**

Kaczawskie Foothills: Bystrzyca 2.1 km N (WIF-14h), Bystrzyca 2.8 km NW (WIF-21c), Wleń 2.2 km NE (WIF-33c), Wleń 2.6 km N (WIF-25a); VII-X; deciduous forest, mixed forest; soil.

Crepidotus cesatii* (Rabenh.) Sacc. var. *cesatii

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-64c – Gniazdo Mt); IX; oak-hornbeam forest; deciduous twigs.

***Crepidotus cesatii* (Rabenh.) Sacc. var. *subsphaerosporus* (J.E. Lange) Senn-Irlet**

Kaczawskie Foothills: Bystrzyca 1.2 km SE (WIF-126n); X; riparian forest; wood.

***Crepidotus epibryus* (Fr.) Quél.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.4 km NW (WIF-46a); XI; riparian forest, forest near a stream, deciduous sapling stand; deciduous twigs, wood, litter. **Kaczawskie Foothills:** Bystrzyca 0.5 km N (WIF-20l), Górczyca 1.1 km SE (WIF-24g); X-XII; mixed forest, overgrown basalt quarry; wood, herbaceous plant debris.

***Crepidotus luteolus* (Lambotte) Sacc.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 2.8 km N (near WIF-44j); X-XI; riparian forest, wastelands at the forest margin; herbaceous plant debris, litter. **Kaczawskie Foothills:** Belczyna 0.8 km S (near WIF-122), Bystrzyca 0.7 km E (WIF-20ax), Bystrzyca 1.3 km N (WIF-16l), Bystrzyca 1.7 km N (WIF-18h), Bystrzyca 2.2 km N (WIF-15a), Bystrzyca 2.5 km NW (WIF-26f), Bystrzyca 2.6 km NW (WIF-26j), Bystrzyca 2.8 km NW (WIF-26i), Przeździec 1.4 km E (WIF-12i), Wleń 1.3 km N (WIF-31); IX-XI; oak-hornbeam forest, coniferous forest, deciduous forest, mixed forest, deciduous forest clearing, ash tree sapling stand; wood, branch, litter, herbaceous plant debris.

Crepidotus mollis* (Schaeff.) Staude var. *mollis

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (WIF-64 – Gniazdo Mt); VII-XI; deciduous forest, riparian forest; wood. **Kaczawskie Foothills:** Belczyna 0.7 km S (WIF-123d), Bystrzyca 1.4 km N (WIF-17c, 19b), Bystrzyca 2 km NW (WIF-15s), Bystrzyca 2.5 km NW (WIF-15h), Przeździec 0.8 km SW (near WIF-22), Wleń 2.3 km N (WIF-25a); VI-XI; mixed forest, riparian forest, oak-hornbeam forest, spruce forest, log landing; wood, *Fraxinus* sp. stump.

***Crepidotus variabilis* (Pers.) P. Kumm.**

Izerskie Foothills: Wleń 1.4 km NW (WIF-46a), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2.3 km NW (WIF-46j); IX-XI; deciduous sapling stand, mixed forest, spruce forest; wood, twigs. **Kaczawskie Foothills:** Belczyna 0.6 km S (WIF-123h), Przeździec 0.7 km S (WIF-22b), Radomiłowice 0.7 km S (near WIF-11), Radomiłowice 1.1 km SW (WIF-11g); VII-IX; coniferous forest, mixed forest, spruce forest margin, under aspen; wood, twigs, litter.

***Crepidotus versutus* (Pers.) Sacc.**

Izerskie Foothills: Wleń 1.5 km SW (ŁuF-65h – Gniazdo Mt); XII; spruce forest; trigs.

***Crinipellis scabella* (Alb. & Schwein.) Murrill**

Kaczawskie Foothills: Bystrzyca 1 km SE (near WIF-126p), Bystrzyca 2.4 km NW (near WIF-21m); VII-X; deciduous forest, forest margin; litter.

***Crucibulum laeve* (Huds.) Kambly**

Kaczawskie Foothills: Bystrzyca 1.5 km NW (WIF-26b), Bystrzyca 2.4 km NW (WIF-21m); VII-IX; oak-hornbeam forest, riparian forest; wood, litter.

***Cyathus olla* (Batsch) Pers.**

Kaczawskie Foothills: Bystrzyca 2.2 km SE (WIF-126o), Przeździec 0.8 km SW (near WIF-21m); VIII-IX; roadside, fallow; soil.

***Cyathus striatus* (Huds.) Willd.**

Kaczawskie Foothills: Bystrzyca 1.4 km N (WIF-17a); VIII; oak-hornbeam forest; stump.

***Cylindrobasidium evolvens* (Fr.) Jülich (= *C. laeve* (Pers.) Chamuris)**

Izerskie Foothills: Maciejowiec 0.5 km S from the palace; II; tree clump; *Malus* sp. branch.

Cystoderma amianthinum* (Scop.) Fayod var. *amianthinum

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-132c); XI; spruce forest; soil. **Kaczawskie Foothills:** Bystrzyca 0.8 km E (near WIF-20w), Radomiłowice 0.8 km W (WIF-12Ar); X; meadow; soil.

Cystoderma carcharias* (Pers.) Fayod var. *carcharias

Kaczawskie Mts.: Bystrzyca 0.9 km S (WIF-132c), Mysłów 1 km E; X-XI; spruce forest, wastelands; soil, rotten wood. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.8 km SW (ŁuF – Gniazdo Mt); IX-X; deciduous forest, spruce forest; litter, soil. **Kaczawskie Foothills:** Belczyna 0.6 km SW (WIF-123h), Belczyna 0.6 km S (WIF-123g), Belczyna 2 km S (WIF-126), Bystrzyca 0.7 km E (WIF-20ax), Bystrzyca 1.8 km S (WIF-126m), Bystrzyca 2 km W (WIF-126n), Wleń 1.7 km NE (WIF-27g); X-XI; coniferous forest, pine forest, spruce forest; soil.

Cystoderma jasonis* (Cooke & Masee) Harmaja var. *jasonis

Izera Mts.: Sępik Mt (tourist trail between Szklarska Poręba and Hala Izerska); X; coniferous forest; soil. **Kaczawskie Mts.:** Rząśnik 1.5 km W (near WIF-290); X; pasture; soil. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.8 km SW (ŁuF – Gniazdo Mt), Zalesie 0.6 km N (LuF); IX-X; mixed forest, spruce-pine forest, spruce forest; soil, rotten stump. **Kaczawskie Foothills:** Belczyna 0.6 km S (near WIF-123g); X; coniferous forest; soil.

***Cystoderma jasonis* (Cooke & Masee) Harmaja var. *lilacipes* (Harmaja) I. Saar.**

Izera Mts.: Sępik Mt (tourist trail between Szklarska Poręba and Hala Izerska); X; coniferous forest; soil.

***Cystoderma jasonis* (Cooke & Masee) Harmaja var. *saarenoksae* (Harmaja) I. Saar**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-132c); X; spruce forest; soil. **Karkonosze Mts.:** Szklarska Poręba; X; spruce forest; soil. **Izerskie Foothills:** Wleń 2 km SW (ŁuF – Gniazdo Mt); IX; beech forest; soil. **Kaczawskie Foothills:** Wleń 1.9 km N (WIF-29g); X; coniferous forest; soil.

***Cystodermella cinnabarina* (Alb. & Schwein.) Harmaja**

Kaczawskie Foothills: Bystrzyca 2.2 km N (WIF-15b); IX; mixed forest; soil.

Cystodermella granulosa* (Batsch) Harmaja var. *granulosa

Kaczawskie Foothills: Wleń 1.7 km NW (WIF-27g); IX; pine forest; soil.

***Cystolepiota bucknallii* (Berk. & Broome) Singer & Clémenton**

Kaczawskie Mts.: Góra Milek Res.; IX; *Cephalanthero-Fagetum*; soil. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); IX; wet spruce forest; soil.

***Cystolepiota seminuda* (Lasch) Bon**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Rzaśnik 1.4 km SW (WIF-290k), Góra Milek Res.; IX-X; *Cephalanthero-Fagetum*, deciduous forest, oak-hornbeam forest; soil. **Izerskie Foothills:** Wleń 1.6 km SW; VII-VIII; deciduous forest; soil. **Kaczawskie Foothills:** Belczyna 0.5 km SW (WIF-123g), Belczyna 0.7 km S (WIF-123b), Bystrzyca 2.1 km N (WIF-14h), Przeździec 1 km SE (WIF-21k), Wleń 2 km N (WIF-30f); VII-XI; mixed forest, deciduous forest, spruce forest, under lime tree; soil.

***Dacrymyces lacrymalis* (Pers.) Sommerf.**

Kaczawskie Foothills: Wleń 1.3 km N (WIF-31h); I; bird cherry shrubs; ?*Padus avium* wood.

***Dacrymyces stillatus* Nees**

Kaczawskie Mts.: Bystrzyca 1.5 km S (WIF-132d); XII; spruce forest; deciduous log. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); V; deciduous forest; *Picea abies* log. **Kaczawskie Foothills:** Bystrzyca 2.4 km NW (WIF-22); XI; mixed forest; *Crataegus* sp. wood.

***Daedalea quercina* (L.) Pers.**

Kaczawskie Foothills: Bystrzyca 1.4 km N (WIF-17a); VII; oak-hornbeam forest; *Quercus* sp. stump.

***Daedaleopsis confragosa* (Bolton) Pers.**

Kaczawskie Mts.: Bystrzyca 0.9 km S (near WIF-127a); riparian forest; *Corylus avellana* branch. **Izerskie Foothills:** Wleń 1.8 km SW (ŁuF – Gniazdo Mt); I; willow shrubs; branch. **Kaczawskie Foothills:** Radomilowice 0.8 km S (near WIF-11); XII; mixed forest; *Salix* sp. wood.

***Datronia mollis* (Sommerf.) Donk**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 2 km SW (ŁuF – Gniazdo Mt); X; mixed forest, spruce forest; deciduous log, *Alnus incana* branch. **Kaczawskie Foothills:** Bystrzyca 1.5 km E (WIF-126d), Bystrzyca 1.8 km N (WIF-19b); IV, VII; deciduous forest, mixed forest; ?*Fagus sylvatica* wood, *Fraxinus* sp. branch.

Deconica montana* (Pers.) P. Kumm. var. *montana

Kaczawskie Mts.: Rząśnik 3 km SW (near WIF-135), Tarczyn 0.6 km E (near WIF-134, Tarczynka Mt); X; slate quarry; soil.

***Delicatula integrella* (Pers.) Fayod.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-132c); VII-VIII; riparian forest, spruce forest; wood. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.7 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest, mixed forest, riparian forest; wood. **Kaczawskie Foothills:** Bełczyna 0.6 km S (WIF-123d), Radomiłowice 0.8 km S (WIF-11i); VIII-X; riparian forest, spruce forest; wood.

***Dendrothele acerina* (Pers.) P.A. Lemke**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); II; mixed forest; *Ulmus* sp. trunk.

***Dichostereum durum* (Bourdot & Galzin) Pilát**

Izerskie Foothills: Wleń 2.1 km SW (ŁuF – Gniazdo Mt); I; mixed forest; ?*Fraxinus* sp. log.

***Echinoderma pseudoasperula* (Knudsen) Bon**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); VIII; under spruce; soil.

***Entoloma araneosum* (Quél.) M.M. Moser**

Izerskie Foothills: Wleń 1.6 km NW (WIF-46), Wleń 1.7 km NW (WIF-46a); VIII; oak-hornbeam forest; soil. **Kaczawskie Foothills:** Bystrzyca 1.4 km N (WIF-17c); X; riparian forest; soil.

Entoloma caccabus* (Kühner) Noordel. var. *caccabus

Kaczawskie Foothills: Bystrzyca 1.2 km SE (WIF-126n); X; riparian forest; soil.

***Entoloma cetratum* (Fr.) M.M. Moser**

Izerskie Foothills: Wleń 1.5 km SW (ŁuF – Gniazdo Mt); XI; spruce forest; soil. **Kaczawskie Mts.:** Rząśnik 2.5 km W (WIF-129d); XI; spruce forest; soil.

Entoloma chalybeum* (Fr.) Noordel. var. *chalybeum

Kaczawskie Mts.: Nowe Rochowice; VIII; calcareous grassland; soil.

***Entoloma chlorophyllum* Noordel.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); IX; oak-hornbeam forest; soil.

Entoloma conferendum* (Britzelm.) Noordel. var. *conferendum

Kaczawskie Mts.: Rząśnik 1.5 km W (near WIF-290); X; pasture; soil. **Karkonosze Mts.:** Karkonosze NP (Czeska Ścieżka), Szklarska Poręba (SzPF-299a, 355g); VIII-IX; spruce forest; soil.

***Entoloma conferendum* (Britzelm.) Noordel. var. *incrustans* (Largent & Thiers) Noordel. & Hauskn.**

Kaczawskie Foothills: Bełczyna 1.8 km SE (WIF-124a); IX; mixed forest; soil.

***Entoloma conferendum* (Britzelm.) Noordel. var. *pusillum* (Velen.) Noordel.**

Izera Mts.: 0,2 km E from “Orle” tourist hostel; IX; meadow; soil. **Kaczawskie Foothills:** Bełczyna 1.1 km SE (WIF-123m); X; spruce clump in oak-hornbeam forest; soil.

***Entoloma conocybecystis* Noordel. & Liiv**

Kaczawskie Foothills: Radomiłowice 0.8 km W (near WIF-12Ar); IX; wet meadow; soil.

***Entoloma euchroum* (Pers.) Donk**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2.1 km SW (ŁuF – Gniazdo Mt); VIII-X; mixed forest near a stream, deciduous forest; wood. **Kaczawskie Foothills:** Bystrzyca 1.3 km SE (WIF-126k), Bystrzyca 2.1 km N (WIF-13g), Bystrzyca 2.3 km N (WIF-13f), Bystrzyca 2.5 km NW (WIF-15h), Bystrzyca 2.8 km NW (WIF-22o); IX-X; deciduous forest, oak-hornbeam forest, forest margin; deciduous wood, deciduous stump.

***Entoloma exiguum* Esteve-Rav. & M. de la Cruz**

Kaczawskie Foothills: Bystrzyca 2.6 km NW (WIF-22k); VI; riparian forest; soil.

***Entoloma fuscotomentosum* F.H. Møller**

Karkonosze Mts.: Karkonosze NP (Równia Pod Śnieżką, 0.45 km W from “Dom Śląski” tourist hostel); IX; gravel excavation on alpine grassland; soil.

***Entoloma griseopruinatum* Noordel. & Cheype**

Kaczawskie Foothills: Wleń 1.6 km NW (WIF-28f); X; deciduous forest; soil.

***Entoloma incanum* (Fr.) Hesler**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry), Nowe Rochowice, Góra Miłek Res.; VIII-X; calcareous grassland; soil.

***Entoloma incarnatofuscescens* (Britzelm.) Noordel.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VII; deciduous forest; soil.

***Entoloma inusitatum* Noordel., Enderle & Lammers**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); IX; oak-hornbeam forest; soil. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest (logging road); soil.

***Entoloma jubatum* (Fr.) P. Karst.**

Izerskie Foothills: Maciejowiec; X; lawn; soil.

Entoloma longistriatum* (Peck) Noordel. var. *longistriatum

Izerskie Foothills: Radomice; IX; limestone quarry (calcareous grassland); soil.

***Entoloma nitens* (Velen.) Noordel.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-64c – Gniazdo Mt); VI; fresh deciduous forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.2 km N (WIF-15c); IX; mixed forest; soil. **Sudetes Foreland:** Szklary 0.5 km SW; IX; overgrown excavation; soil.

***Entoloma noordeloosi* Hauskn.**

Kaczawskie Foothills: Bystrzyca 2.1 km E (near WIF-125); IX; deciduous tree clump; soil.

***Entoloma occultripigmentatum* Arnolds & Noordel. var. *cystidiatum* Hauskn. & Noordel.**

Kaczawskie Foothills: Belczyna 1.1 km SW (WIF-123m); X; deciduous forest; soil. **Kaczawskie Mts.:** Tarczyn 0.6 km E (Tarczynka Mt); IX; tree clump; soil.

Entoloma occultripigmentatum* Arnolds & Noordel. var. *occultripigmentatum

Kaczawskie Foothills: Belczyna 1 km E (WIF-122k); IX; mixed tree clump; soil.

***Entoloma pleopodium* (Bull.) Noordel.**

Kaczawskie Mts.: Rzaśnik 2.3 km W (WIF-128c); X; deciduous forest; soil. **Kaczawskie Foothills:** Belczyna 1.8 km SE (WIF-126a), Przeździeca 1.3 km W (WIF-24r); IX-X; alder sapling stand, mixed forest; soil.

Entoloma poliopus* (Romagn.) Noordel. var. *poliopus

Kaczawskie Foothills: Nad Groblą Res.; IX; riparian forest; soil.

***Entoloma politum* (Pers.) Donk**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; soil. **Izerskie Foothills:** Wleń 2.1 km SW (ŁuF – Gniazdo Mt); IX; wet mixed forest; soil.

***Entoloma porphyrophaeum* (Fr.) P. Karst.**

Izera Mts.: Świeradów-Zdrój (ŚF-291f); X; grassy forest margin; soil. **Kaczawskie Foothills:** Belczyna 1.1 km SE (near WIF-123m), Bystrzyca 1.1 km E (near WIF-126k); X; meadow; soil.

***Entoloma rhodocalix* (Lasch) M.M. Moser**

Izerskie Foothills: Zalesie 0.8 km N (LuF); IX; boarder between mire and pine forest; soil.

Entoloma rhodopolium* (Fr.) P. Kumm. var. *rhodopolium

Izerskie Foothills: Wleń 2.8 km N (WIF-44j); X; riparian forest; soil. **Kaczawskie Foothills:** Bełczyna 1.1 km E (WIF-122a), Bystrzyca 1.2 km SE (WIF-126o); VIII-X; deciduous tree clump, deciduous forest; soil.

***Entoloma rhodopolium* (Fr.) P. Kumm. var. *pseudopolitum* Noordel.**

Kaczawskie Mts.: Rząśnik 2.3 km W (WIF-128g); X; forest near a stream; soil.

***Entoloma sepium* (Noulet & Dass.) Richon & Roze (= *E. saepium* (Noulet-Dasier) Richon & Roze)**

Izerskie Foothills: Wleń; V; lawn (under plum-tree); soil.

***Entoloma sericeum* Qué. var. *cinereoopacum* Noordel.**

Kaczawskie Mts.: Tarczyn 0.6 km E (Traczynka Mt, quarry); IX; grassland; soil. **Kaczawskie Foothills:** Sobota 1.5 km S; IV-XI; pasture; soil.

Entoloma sericeum* Qué. var. *sericeum

Kaczawskie Mts.: Rząśnik 1.5 km W (WIF-290a); IX; meadow; soil. **Kaczawskie Foothills:** Wleń 2 km N (WIF-29d); X; coniferous forest; soil.

***Entoloma solstitiale* (Fr.) Noordel.**

Kaczawskie Foothills: Bełczyna 1.1 km SE (near WIF-123m); X; meadow; soil.

***Entoloma tjallingiorum* Noordel.**

Izerskie Foothills: Wleń 1.7 km SW (ŁuF – Gniazdo Mt); VIII; spruce forest; soil.

***Entoloma undatum* (Fr.) M.M. Moser**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; riparian forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.8 km NW (WIF-22o), Wleń 2.3 km N (WIF-25a); VII-IX; oak-hornbeam forest, deciduous forest; soil.

Entoloma vinaceum* (Scop.) Arnolds & Noordel. var. *vinaceum

Izera Mts.: Sępik Mt (tourist trail between Szklarska Poręba and Hala Izerska); X; coniferous forest; soil. **Karkonosze Mts.:** Karkonosze NP (Mumławski Wierch Mt); X; spruce forest; soil.

***Entoloma xanthochroum* (P.D. Orton) Noordel.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); IX; riparian forest; soil.

***Eocronartium muscicola* (Pers.) Fitzp.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VII; deciduous forest; mosses.

***Exidia plana* Donk**

Kaczawskie Foothills: Bystrzyca 2.1 km NW (WIF-26c), Bystrzyca 2.2 km N (WIF-15b), Przeździec 1.7 km E (WIF-12l); XII-II; deciduous forest, mixed forest; *Corylus avellana* branch, *Crataegus* sp. branch, *Picea abies* log.

***Exidia recisa* (Ditmar) Fr.**

Izerskie Foothills: Wleń 2.8 km N (near WIF-44); I; wastelands near forest; *Salix* sp. branch.

***Exidia saccharina* (Alb. & Schwein.) Fr.**

Izerskie Foothills: Wleń 2.3 km SW (ŁuF – Gniazdo Mt); III-V; spruce-pine forest; *Pinus sylvestris* log.

***Exidia truncata* Fr. (= *E. glandulosa* (Bull.) Fr.)**

Kaczawskie Mts.: Bystrzyca 1.1 km SE (WIF-131), Rząśnik 1.5 km W (WIF-290m); I-III; oak forest, overgrown pond; *Quercus* sp. branch. **Izerskie Foothills:** Wleń 1.4 km NW (WIF-46g); VIII; mixed forest; *Quercus* sp. branch. **Kaczawskie Foothills:** Górczyca 1.4 km W (WIF-24f), Przeździec 0.9 km S (WIF-22k), Radomilowice 1.1 km SW (WIF-11g), Wleń 1.5 km NE (WIF-28l); I-IV; mixed forest, riparian forest; *Quercus* sp. branch, *Fraxinus* sp. branch. **Sudetes Foreland:** Szklary 0.3 km S; I; tree clump; *Salix* sp. branch.

***Exidia villosa* Neuhoﬀ**

Kaczawskie Mts.: near Bystrzyca (WIF-134dx), Rząśnik 1.4 km SW (WIF-290k), Wleń 1.4 km SE (WIF-137h), Wleń 1.7 km SE (WIF-137o); I-III; deciduous forest, lime forest, pine sapling stand; *Tilia* sp. branch. **Izerskie Foothills:** Wleń 1.6 km NW (WIF-47f); V; deciduous forest; *Tilia* sp. branch. **Kaczawskie Foothills:** Bystrzyca 1.4 km N (WIF-16k), Przeździec 1.2 km SW (WIF-23d), Tarczyn 0.6 km E (WIF-134), Wleń 2 km N (WIF-30f); I-VII; oak-hornbeam forest, deciduous forest, mixed forest, roadside shelterbelt, under lime tree; *Tilia* sp. branch.

***Fistulina hepatica* (Schaeff.) With.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Grudno (quarry); VII, IX; beech forest, oak-hornbeam forest; *Quercus* sp. trunk. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); VIII; mixed forest; *Quercus* sp. stump. **Kaczawskie Foothills:** Bystrzyca 1.2 km SE (WIF-126n), Górczyca 1.5 km W (WIF-24a), Radomilowice 1.4 km W (WIF-12Ah, 12Ai), Wleń 0.9 km NW (WIF-32b), Wleń 1.4 km N (WIF-32c), Wleń 2.1 km N (WIF-30b), Nad Groblą Res.; VIII-IX; oak forest, lime forest, lime-oak forest, mixed forest, riparian forest; *Quercus* sp. trunk and stump, ?*Tilia* sp. stump. **Sudetes Foreland:** Szklary (near estate no. 11); VIII; *Quercus* sp. stump.

***Flammula alnicola* (Fr.) P. Kumm. (= *Pholiota alnicola* (Fr.) Singer)**

Izera Mts.: Świeradów-Zdrój (ŚF-238b); X; spruce forest; wood. **Kaczawskie Mts.:** Rząśnik 2.3 km W (WIF-128g); X; mixed forest; wood. **Kaczawskie Foothills:** Sobota 1.7 km SW (WIF-24Ab); IX; deciduous forest; deciduous trunk base.

***Flammulaster rhombisporus* (G.F. Atk.) Watling**

Kaczawskie Foothills: Bystrzyca 2 km N (WIF-16a), Bystrzyca 2.5 km N (WIF-15h); X; deciduous forest; *Tilia* sp. log, wood.

***Flammulina elastica* (Lasch) Redhead & R.H. Petersen**

Kaczawskie Foothills: Wleń 2.1 km N (WIF-30c); I; sapling stand margin near a stream; *Salix* sp./*Alnus* sp. trunk.

Flammulina velutipes* (Curtis) P. Karst. var. *velutipes

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); XI-XII; deciduous forest, riparian forest; *Fraxinus* sp. wood, wood. **Kaczawskie Foothills:** Bystrzyca 2 km NE (WIF-15s), Wleń 2.3 km SW (WIF-28d), Wleń 2.6 km N (WIF-25a), Wleń 2.1 km N (WIF-30c); XI-I; deciduous forest, log landing near deciduous forest, sapling stand margin near a stream; wood, litter, branch.

***Fomes fomentarius* (L.) Fr.**

Kaczawskie Foothills: Radomiłowice 0.7 km SW (WIF-12a); IV; mixed forest; *Betula* sp. trunk.

***Galerina ampullaceocystis* P.D. Orton**

Izerskie Foothills: Wleń 2.2 km NW (WIF-46c); V; pine forest; wood.

***Galerina atkinsoniana* A.H. Sm.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-59b – Gniazdo Mt); X; riparian forest; soil. **Kaczawskie Foothills:** Belcyna 1.4 km SE (near WIF-123s); X; meadow boarder (under oak); litter.

***Galerina clavata* (Velen.) Kühner.**

Kaczawskie Mts.: Tarczyn 1.8 km E (near WIF-133); IX-X; riparian forest, meadow; soil. **Kaczawskie Foothills:** Radomiłowice 0.8 km W, Sobota 1.5 km S; X-XI; swamp at the forest margin, pasture; soil. **Sudetes Foreland:** Szklary 0.5 km W; X; overgrown spoil tip; mosses.

***Galerina graminea* (Velen.) Kühner (= *G. laevis* (Pers.) Singer)**

Izerskie Foothills: Radomice (limestone quarry); XII; overgrown calcareous grassland; soil. **Kaczawskie Foothills:** Przeździec 1 km SW (near WIF-23), Sobota 1.5 km S, Wleń 2.8 km N (near WIF-25b); X-XI; meadow, overgrown pasture; soil.

***Galerina heimansii* Reijnders**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt, Białe Skały); II; spruce-pine forest; litter.

***Galerina hybrida* Kühner**

Izerskie Foothills: Zalesie 0.6 km N (LuF); VIII-IX; mire; *Sphagnum* sp.

***Galerina hypnorum* (Schränk) Kühner s Horak**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-132c); X; spruce forest; mosses. **Izerskie Foothills:** Radomice (limestone quarry), Wleń 1.1 km SW (ŁuF-58m), Wleń 1.4 km SW (ŁuF-59b – Gniazdo Mt), Wleń 1.9 km SW (ŁuF – Gniazdo Mt); VIII-X; calcareous grassland, mixed forest, riparian forest; mosses, mossy wood. **Kaczawskie Foothills:** Bystrzyca 2.2 km NE (WIF-15d), Wleń 1.4 km N (WIF-32c), Wleń 2.4 km NE (WIF-26m); VII-XI; oak-hornbeam forest, deciduous forest, mixed forest; mossy wood, mosses.

***Galerina karstenii* A.H. Sm. & Singer**

Izera Mts.: 0,2 km E from “Orle” tourist hostel; IX; meadow; soil.

***Galerina marginata* (Batsch) Kühner s.l.**

Izera Mts.: Świeradów-Zdrój (ŚF-452b); X; spruce forest; wood. **Kaczawskie Mts.:** Bystrzyca 1 km S (WIF-133a); X; riparian forest; litter. **Karkonosze Mts.:** Szklarska Poręba (SzPF-350b); X; spruce forest; wood. **Izerskie Foothills:** Radomice (limestone quarry), Wleń 2.3 km SW (ŁuF – Gniazdo Mt); X; mixed forest, calcareous grassland; wood, ?soil. **Kaczawskie Foothills:** Belczyna 0.6-0.7 km S (WIF-123g), Przeździec 0.8 km SW (WIF-221), Radomiłowice 1.2 km S (WIF-14a); IX-XI; mixed forest, coniferous forest, spruce forest; wood, litter.

***Galerina mniophila* (Lasch) Kühner**

Izera Mts.: Sępik Mt (tourist trail between Szklarska Poręba and Hala Szrenicka); X; forest roadside; mosses.

***Galerina pallida* (Pilát) E. Horak & M.M. Moser**

Kaczawskie Mts.: Nielestno 1 km E (CzF-150), Wleń 1.7 km SE (WIF-137o); I; lime forest, spruce forest (logging road); wood.

***Galerina paludosa* (Fr.) Kühner**

Izera Mts.: at the Izera river (0.4 km S from Torfowiska Doliny Izery Res. towards Jagniątków); VI; spruce forest; mosses. **Izerskie Foothills:** Zalesie 0.8 km N (LuF); VII; mire; *Sphagnum* sp.

Galerina pumila* (Pers.) M. Lange var. *pumila

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1 km SE (WIF-132c), Rząśnik 1.5 km W (WIF-290m), Tarczyn 0.9 km NE (WIF-134c); IX-XI; oak forest, spruce forest; soil, mosses. **Kaczawskie Foothills:** Belczyna 0.7 km S (WIF-123g), Wąwóz Lipy Res.; X-XI; oak forest, spruce forest; mosses.

***Galerina sideroides* (Bull.) Kühner**

Kaczawskie Foothills: Rzaśnik 1.3 km NW (WIF-125g); X; mixed forest; wood.

***Galerina triscopa* (Fr.) Kühner**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); X; riparian forest; mossy log.

***Galerina uncialis* (Britzelm.) Kühner**

Izera Mts.: Sępik Mt (tourist trail between Szklarska Poręba and Hala Szrenicka); X; forest roadside; wood.

Kaczawskie Mts.: Wleń 1.5 km E (WIF-137h); X; pine sapling stand; wood. **Izerskie Foothills:** Wleń 2.8 km N (WIF-44j); X; oak-hornbeam forest; soil. **Sudetes Foreland:** Szklary 0.5 km W; X; overgrown spoil tip; wood.

***Galerina vittiformis* (Fr.) Singer var. *vittiformis* f. *bispora* A.H. Sm. & Singer**

Kaczawskie Foothills: Wleń 2.3 km NE (WIF-25c); VIII; spruce forest; soil.

***Ganoderma lipsiense* (Batsch) G.F. Atk. (= *G. applanatum* (Pers.) Pat.)**

Izerskie Foothills: Wleń 1.4 km NW (WIF-46a); XI; deciduous sapling stand; deciduous log. **Kaczawskie Foothills:** Bystrzyca 2.1 km NW (WIF-26c), Wleń 1.9 km NE (WIF-28d); VIII-IX; deciduous forest, mixed forest; trunk base, stump.

***Geastrum fimbriatum* Fr.**

Kaczawskie Mts.: Góra Miłek Res.; IX; *Cephalanthero-Fagetum*; soil. **Izerskie Foothills:** near Pławna Górna (limestone quarry), Wleń 1.4 km SW (ŁuF – Gniazdo Mt); IX-X; oak-hornbeam forest, overgrown quarry; soil.

Kaczawskie Foothills: near Nad Groblą Res.; VIII; mixed forest; soil.

***Geastrum rufescens* Pers.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); IX; under spruce; soil.

***Gerronema xanthophyllum* (Bres.) Norvell, Redhead & Ammirati (= *Chrysomphalina strombodes* (Berk. & Mont.) Clémenton)**

Kaczawskie Foothills: Wleń 1.4 km N (WIF-31c); VIII; mixed forest; wood.

***Gloeophyllum abietinum* (Bull.) P. Karst.**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt, Białe Skąły); IX-X; pine forest; coniferous log, *Pinus sylvestris* log. **Kaczawskie Foothills:** Bystrzyca 1.4 km N (WIF-19b), Wleń 2.3 km NE (WIF-25c); II, X; mixed forest, spruce forest; *Alnus* sp. log, *Picea abies* log.

***Gloeophyllum odoratum* (Wulfen) Imazeki**

Kaczawskie Mts.: Bystrzyca 1.4 km SE (WIF-130f); II; spruce forest; *Picea abies* stump. **Kaczawskie Foothills:** Bystrzyca 1.4 km N (WIF-17a), Skąła (Palace Park); I; oak-hornbeam forest, park; *Pinus sylvestris* stump, *Picea abies* stump.

***Gloeophyllum sepiarium* (Wulfen) P. Karst.**

Kaczawskie Foothills: Bystrzyca 2.5 km NW (WIF-26f); IV; coniferous forest; *Picea abies* log.

***Gloeoporus dichrous* (Fr.) Bres.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); XI; oak-hornbeam forest; *Alnus* sp. log. **Izerskie Foothills:** Wleń 2.1 km SW (ŁuF – Gniazdo Mt); XII; mixed forest; wood. **Kaczawskie Foothills:** Wleń 2.3 km NE (WIF-25c), Bystrzyca 2.6 km NW (WIF-21h); II, X; spruce forest, mixed forest; wood.

***Gomphidius glutinosus* (Schaeff.) Fr.**

Kaczawskie Mts.: Tarczyn 0.9 km NE (WIF-134c); X; spruce forest; soil. **Karkonosze Mts.:** Szklarska Poręba (Złoty Potok Valley); XI; spruce forest; soil. **Izerskie Foothills:** Zalesie 0.6 km N (LuF); IX; spruce-pine forest; soil.

***Gomphus clavatus* (Pers.) Gray**

Kaczawskie Mts.: Góra Milek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Grifola frondosa* (Dicks.) Gray**

Kaczawskie Mts.: Buki Sudeckie Res.; X; *Dentario enneaphylli-Fagetum*; *Fagus sylvatica* trunk base.

***Gymnopilus penetrans* (Fr.) Murrill**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1.2 km S (WIF-133b), between Nowe Rochowice and Grudno, Wleń 1.2 km E (WIF-136f); VIII-X; oak-hornbeam forest, mixed forest; wood. **Izerskie Foothills:** Wleń 1.8 km SW (ŁuF – Gniazdo Mt), Zalesie 0.8 km NE (LuF-200h); X-XI; oak forest, dry mire margin; wood, litter. **Kaczawskie Foothills:** Bystrzyca 1.4 km N (WIF-19b), Wleń 1.7 km NE (WIF-27g), Wleń 1.9 km N (WIF-29g), Wleń 2.3 km NE (WIF-25c); IX-X; pine forest, spruce clump in oak-hornbeam forest, spruce forest; wood, *Picea abies* stump.

***Gymnopilus picreus* (Pers.) P. Karst.**

Izerskie Foothills: Zalesie 0.6 km N (LuF); IX; pine forest; wood.

***Gymnopus acervatus* (Fr.) Murrill**

Kaczawskie Foothills: Bystrzyca 1.5 km E (near WIF-126d); IX; deciduous forest; wood.

***Gymnopus androsaceus* (L.) J.L. Mata & R.H. Petersen. (= *Setulipes androsaceus* (L.) Antonín)**

Kaczawskie Mts.: Rząśnik 2.9 km W (WIF-135j); XII; young spruce forest; litter. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2.1 km SW (ŁuF – Gniazdo Mt); V-VII; spruce forest; litter. **Kaczawskie Foothills:** near Nad Groblą Res.; VIII; spruce forest; litter.

***Gymnopus aquosus* (Bull.) Antonín & Noordel.**

Kaczawskie Foothills: Wleń 1.5 km NW (near WIF-31a); V; forest margin; litter.

***Gymnopus confluens* (Pers.) Antonín, Halling & Noordel.**

Kaczawskie Mts.: Bystrzyca 1 km SE (WIF-131a); VII; riparian forest; litter. **Kaczawskie Foothills:** Bystrzyca 1.2 km SE (WIF-126o), Bystrzyca 2.1 km N (WIF-14h), Leszczyna 1 km N (near Ciche Szczyście mine), Wleń 2.6 km N (WIF-25a); IV-X; oak-hornbeam forest, deciduous forest, mixed forest, black alder forest; litter.

***Gymnopus dryophilus* (Bull.) Murrill**

Izerskie Foothills: Wleń 2.3 km NW (near WIF-46c); VII; field at the forest margin; litter.

***Gymnopus hariolorum* (Bull.) Antonín, Halling & Noordel.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 2 km SW (ŁuF – Gniazdo Mt); V-VI; mixed forest; litter.

***Gymnopus ocior* (Pers.) Antonín & Noordel.**

Kaczawskie Foothills: Bystrzyca 2.1 km N (WIF-14h); IX; mixed forest; litter.

***Gymnopus peronatus* (Bolton) Gray**

Kaczawskie Foothills: Bystrzyca 1.4 km N (WIF-19g); X; mixed forest; soil.

***Gyroporus castaneus* (Bull.) Quél.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; soil. **Kaczawskie Foothills:** Górczyca 1.4 km W (WIF-24h); VII; mixed forest; soil.

***Hapalopilus nidulans* (Fr.) P. Karst.**

Kaczawskie Mts.: Wleń 2.1 km E (near WIF-136a); IV; oak-hornbeam forest; *Corylus avellana* branch.

Kaczawskie Foothills: Bystrzyca 1.3 km SE (WIF-126k); VII, XI; deciduous forest; deciduous wood, *Sorbus aucuparia* branch.

***Hebeloma collariatum* Bruchet**

Izerskie Foothills: Radomice (limestone quarry); X; overgrown quarry; soil.

***Hebeloma crustuliniforme* (Bull.) Quél. s.str.**

Kaczawskie Foothills: Przeździec 1.4 km E (WIF-12i); X; forest; soil. **Sudetes Foreland:** Szklary 0.5 km W; IX; overgrown spoil tip; soil.

***Hebeloma mesophaeum* (Pers.) Quél.**

Karkonosze Mts.: Szklarska Poręba (SzPF-347c); IX; spruce forest; soil. **Izerskie Foothills:** Radomice (limestone quarry); X; overgrown quarry; soil.

***Hebeloma theobrominum* Quad. (= *H. truncatum* (Schaeff.) P. Kumm.)**

Kaczawskie Foothills: Wąwóz Lipy Res.; XI; oak forest; soil.

***Hebeloma velutipes* Bruchet**

Izera Mts.: Świeradów-Zdrój (ŚF-238k); X; mixed forest; soil. **Izerskie Foothills:** Gryfów Śląski 5 km NE, Zalesie 0.7 km N (LuF); IX-X; mixed forest, mire; soil, *Sphagnum* sp.

***Hemimycena candida* (Bres.) Singer**

Izerskie Foothills: Wleń 1.3 km NW (WIF-47s); IX; log landing; comfrey roots.

Hemimycena delectabilis* (Peck) Singer s. auct. Eur. var. *delectabilis

Kaczawskie Foothills: Bystrzyca 1.9 km N (WIF-18d); XI; oak-hornbeam forest; deciduous wood.

***Hemimycena gracilis* (Quél.) Singer**

Izerskie Foothills: Wleń 1.5 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VII-IX; spruce forest; litter. **Kaczawskie Foothills:** Radomiłowice 1.2 km S (WIF-14b); VIII; mixed forest; *Picea abies* litter.

Hemimycena lactea* (Pers.) Singer var. *lactea

Izerskie Foothills: Maciejowiec (Palace Park), Radomice (limestone quarry); IX-X; park, overgrown quarry; litter.

***Hemimycena pseudocrispata* (Valla) Maas Geest.**

Kaczawskie Foothills: Bełczyna 0.8 km S (WIF-123j); VIII; spruce forest; bark.

***Hemipholiota populnea* (Pers.) Bon (= *Pholiota populnea* (Pers.) Kuyper & Tjall.)**

Kaczawskie Foothills: Przeździec 0.2 km (by the road to Wleń); XI; roadside; *Populus* sp. trunk.

***Henningsomyces candidus* (Pers.) Kuntze**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); XI; oak-hornbeam forest; wood.

***Hericium cirrhatum* (Pers.) Nikol. (= *Creolophus cirrhatus* (Pers.) P. Karst.)**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); IX; riparian forest; *Acer pseudoplatanus* trunk.

***Heterobasidion annosum* (Fr.) Bref.**

Kaczawskie Foothills: near Wąwóz Myśliborski Res. towards Myślibórz; XI; riparian forest; *Picea abies* log.

***Heterobasidion parviporum* Niemelä & Korhonen**

Kaczawskie Mts.: Tarczyn 0.9 km NE (WIF-134c); XI; young spruce forest; coniferous log.

***Hohenbuehelia mastrucata* (Fr.) Singer**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest; deciduous twig. **Kaczawskie Foothills:** Wleń 2.7 km NE (WIF-26d); IX; riparian forest; wood.

***Hydnellum peckii* Banker**

Izerskie Foothills: Wleń 1.8 km SW (ŁuF – Gniazdo Mt); VIII; wet spruce forest; soil.

***Hydnum repandum* L.**

Kaczawskie Foothills: Wąwóz Lipy Res.; XI; oak forest; soil.

***Hydnum rufescens* Pers.**

Karkonosze Mts.: Szklarska Poręba (Złoty Potok Valley); XI; spruce forest; soil.

***Hydropus scabripes* (Murrill) Singer var. *quadrisporus* Bas**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-59b – Gniazdo Mt); VI; riparian forest; soil. **Kaczawskie Foothills:** Rząśnik 1.3 km NW (WIF-125g); X; mixed forest; *Fagus sylvatica*/*Fraxinus* sp. wood.

***Hygrocybe acutoconica* (Clem.) Singer var. *acutoconica* (= *H. persistens* (Britzelm.) Singer)**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry); VIII-X; calcareous grassland; soil. **Izerskie Foothills:** Radomice (limestone quarry); IX; calcareous grassland; soil.

***Hygrocybe calciphila* Arnolds**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry), Góra Milek Res.; VIII-IX; calcareous grassland; soil.

***Hygrocybe cantharellus* (Schwein.) Murrill (= *H. lepida* Arnolds)**

Karkonosze Mts.: Karkonosze NP (Hala pod Łabskim Szczytem); IX; grassland, soil. **Izerskie Foothills:** Zalesie 0.7 km N (LuF); VIII; mire; *Sphagnum* sp.

***Hygrocybe ceracea* (Wulfen) P. Kumm.**

Kaczawskie Mts.: Buki Sudeckie Res.; IX; *Dentario enneaphylli-Fagetum*; soil. **Kaczawskie Foothills:** Belczyna 0.6 km E (near WIF-122h), Belczyna 1 km SE (near WIF-123), Bystrzyca 0.8 km E (near WIF-20w, 20y, 20bx), Bystrzyca 1.5 km E (near WIF-126), Bystrzyca 1.5 km NE (near WIF-131), Radomiłowice 0.8 km W (near WIF-12Ar), Sobota 1.5 km S; VII-XI; meadow, pasture, overgrown pasture; soil.

***Hygrocybe chlorophana* (Fr.) Wünsche**

Kaczawskie Mts.: Milek Mt; VIII; calcareous grassland; soil.

***Hygrocybe coccinea* (Schaeff.) P. Kumm.**

Kaczawskie Foothills: Radomiłowice 0.8 km W (near WIF-12Ar); X; meadow; soil.

Hygrocybe conica* (Schaeff.) P. Kumm. var. *conica

Kaczawskie Mts.: Mysłów, Nowe Rochowice, Buki Sudeckie Res., Miłek Mt; VIII-IX; *Dentario enneaphylli-Fagetum*, calcareous grassland; soil. **Kaczawskie Foothills:** Bystrzyca 0.8 km E (near WIF-20w), Bystrzyca 1.2 km E (near WIF-126), Bystrzyca 1.5 km NE (near WIF-131), Wleń 1.9 km NE (WIF-27d); IX-XI; meadow, mixed forest; soil.

***Hygrocybe conica* (Schaeff.) P. Kumm. var. *conicopalustris* Arnolds**

Izerskie Foothills: Radomice (limestone quarry); IX; calcareous grassland; soil.

Hygrocybe glutinipes* (J.E. Lange) R. Haller Aar. var. *glutinipes

Kaczawskie Foothills: Belczyna 1.8 km SE (WIF-124d), Wleń 1.9 km NE (WIF-27d); VIII-IX; deciduous forest, riparian forest; soil.

***Hygrocybe insipida* (J.E. Lange) M.M. Moser**

Kaczawskie Foothills: Belczyna 1 km SE (near WIF-123), Belczyna 1.1 km SE (near WIF-123m), Bystrzyca 0.8 km E (near WIF-20w), Bystrzyca 1.5 km NE (near WIF-131); X-XI; meadow; soil.

***Hygrocybe irrigata* (Pers.) Bon (= *H. unguinosa* (Fr.) P. Karst.)**

Kaczawskie Foothills: Belczyna 1.1 km SE (near WIF-123m), Bystrzyca 0.8 km E (near WIF-20w); VIII-X; meadow; soil.

***Hygrocybe lacmus* (Schumach.) P.D. Orton & Watling (= *Camarophyllus lacmus* (Schum.) J.E. Lange)**

Kaczawskie Foothills: Radomiłowice 0.8 km W (near WIF-12Ar); X; meadow; soil.

Hygrocybe miniata* (Fr.) P. Kumm. var. *miniata

Izera Mts.: Świeradów-Zdrój (ŚF-291f), Jelenia Łąka, 0.2 km E from “Orle” tourist hostel; IX-X; meadow, roadside; soil. **Kaczawskie Mts.:** Bystrzyca 1.3 km SE (WIF-130b), Miłek Mt; VIII; deciduous forest, calcareous grassland; soil. **Karkonosze Mts.:** Jakuszyce 1.5 km SE (mire near Mumłowski Wierch Mt); X; mire; soil.

***Hygrocybe phaeococcinea* (Arnolds) Bon**

Izera Mts.: tourist trail between Jakuszyce and Orle; IX; mossy roadside; soil.

***Hygrocybe pratensis* (Pers.) Murrill var. *pallida* (Cooke) Arnolds**

Kaczawskie Foothills: Belczyna 1 km S (near WIF-123n), Belczyna 1.4 km SE (near WIF-123s); X; meadow; soil.

***Hygrocybe pratensis* (Pers.) Murrill var. *pratensis* (= *Camarophyllus pratensis* (Pers.) Kumm.)**

Kaczawskie Foothills: Bystrzyca 0.7 km E (near WIF-20w), Radomiłowice 0.8 km W (near WIF-12Ar), Sobota 1.5 km S; IX-XI; pasture, meadow; soil.

Hygrocybe psittacina* (Schaeff.) P. Kumm. var. *psittacina

Kaczawskie Foothills: Bystrzyca 0.8 km E (near WIF-20bx), Bystrzyca 1.5 km NE (near WIF-131), Radomiłowice 0.8 km W (near WIF-12Ar); X-XI; meadow; soil.

***Hygrocybe splendidissima* (P.D. Orton) M.M. Moser**

Kaczawskie Foothills: Radomiłowice 0.8 km W (near WIF-12Ar); X; meadow; soil.

***Hygrocybe virginea* (Wulfen) P.D. Orton & Watling var. *virginea* (= *Camarophyllus virgineus* (Wulf.) P. Kumm.)**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry), Miłek Mt; X; calcareous grassland; soil. **Kaczawskie Foothills:** Bełczyna 1.1 km SE (WIF-123m), Bełczyna 1.2 km S (near WIF-123m), Bystrzyca 1.5 km E (near WIF-126i), Sobota 1.5 km S; X-XI; meadow, pasture, deciduous forest margin; soil. **Sudetes Foreland:** Szklary 0.5 km W; X; overgrown spoil tip; soil.

***Hygrophoropsis aurantiaca* (Wulfen.) Maire**

Kaczawskie Mts.: Bystrzyca 1.4 km SE (WIF-130f), Rząśnik 2.2 km W (near WIF-128); VIII-IX; deciduous forest, spruce forest; *Picea abies* stump, litter. **Kaczawskie Foothills:** Bystrzyca 2.2 km N (WIF-13d); XI; deciduous forest; litter.

***Hygrophorus chrysodon* (Batsch) Fr.**

Kaczawskie Mts.: Góra Miłek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Hygrophorus eburneus* (Bull.) Fr.**

Kaczawskie Mts.: Góra Miłek Res.; X; *Cephalanthero-Fagetum*; soil. **Izerskie Foothills:** Wleń 2 km SW (ŁuF – Gniazdo Mt); X; beech forest; soil.

Hygrophorus hypothejus* (Fr.) Fr. var. *hypothejus

Kaczawskie Mts.: Tarczyn 0.6 km E (near WIF-134 – Tarczynka Mt); XII-I; mixed tree clump; soil. **Kaczawskie Foothills:** Bystrzyca 1.8 km E (WIF-126m), Bystrzyca 2.6 km NW (WIF-26j); X-XII; pine forest, mixed forest; soil.

Hygrophorus lucorum* Kalchbr. var. *lucorum

Izera Mts.: Kobylina Mt (N slope); XI; pine-larch sapling stand; soil. **Izerskie Foothills:** Łupki (near tourist hostel); XI; lawn (under larch); soil. **Kaczawskie Foothills:** Bystrzyca 0.7 km E (WIF-20ax), Bystrzyca 1.7 km N (WIF-18h), Bystrzyca 2.5 km NW (WIF-26k), Bystrzyca 2.6 km NW (WIF-21k, 26j), Przeździec 0.9 km S (near WIF-22a); X-XII; coniferous forest, mixed forest, mixed sapling stand; soil.

***Hygrophorus nemoreus* (Pers.) Fr.**

Izerskie Foothills: Wleń 1.4 km NW (WIF-46g); IX; mixed forest; soil. **Kaczawskie Foothills:** Bełczyna 1.3 km E (WIF-122d), Bełczyna 2 km SE (WIF-125b), Górczyca 1.5 km W (WIF-24a), Wleń 0.9 km NW (WIF-32b), Wleń 2.6 km N (WIF-25a), Nad Groblą Res.; VII-X; oak forest, oak-hornbeam forest, deciduous forest, mixed forest; soil.

***Hygrophorus olivaceoalbus* (Fr.) Fr.**

Karkonosze Mts.: Szklarska Poręba (SzPF-350g); VIII; mixed forest; soil. **Izera Mts.:** 0.4 km from Torfowiska Doliny Izery Res. (at the Izera river); IX; spruce forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.1 km N (WIF-14h); IX; mixed forest; soil.

***Hygrophorus penarius* Fr.**

Karkonosze Mts.: Szklarska Poręba (Złoty Potok Valley); XI; beech forest; soil.

***Hygrophorus piceae* Kühner**

Kaczawskie Mts.: Bystrzyca 1.3 km SE (WIF-130c); X; mixed forest; soil. **Kaczawskie Foothills:** Bełczyna 0.7 km S (WIF-123g); young spruce forest; soil.

***Hygrophorus pustulatus* (Pers.) Fr.**

Izera Mts.: Świeradów-Zdrój (ŚF-308f); X; spruce forest; soil. **Kaczawskie Mts.:** Tarczyn 0.9 km NE (WIF-134c); X; spruce forest; soil. **Izerskie Foothills:** Wleń 2.8 km N (WIF-44j); X; oak-hornbeam forest; soil. **Kaczawskie Foothills:** Bystrzyca 1.3 km N (WIF-16l); XII; mixed forest; soil.

***Hygrophorus tephroleucus* (Pers.) Fr.**

Kaczawskie Mts.: Bystrzyca 1.1 km S (WIF-132c); riparian forest; soil. **Kaczawskie Foothills:** Bystrzyca 1.7 km N (WIF-18h), Rzaśnik 1.3 km NW (WIF-125g); X; mixed forest, spruce forest margin; soil.

***Hymenochaete cinnamomea* (Pers.) Bres.**

Izerskie Foothills: Wleń 1.6 km SE (ŁuF – Gniazdo Mt); XI; deciduous forest; *Padus serotina* branch.

***Hymenopellis radicata* (Relhan) R.H. Petersen f. *radicata* (= *Xerula radicata* (Relh.) Dörfelt)**

Kaczawskie Mts.: Buki Sudeckie Res.; IX; *Dentario enneaphylli-Fagetum*; soil. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2 km SW (ŁuF – Gniazdo Mt); IX-XI; beech forest, mixed forest; wood. **Kaczawskie Foothills:** Wleń 2.6 km N (WIF-25a); VIII; deciduous forest; wood.

***Hymenopellis radicata* (Relhan) R.H. Petersen f. *marginata* (Konrad & Maubl.) R.H. Petersen**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); IX; riparian forest; wood. **Kaczawskie Foothills:** Bystrzyca 1.4 km N (WIF-19); VIII; mixed forest; wood.

***Hyphoderma setigerum* (Fr.) Donk**

Kaczawskie Mts.: Wleń 1.8 km SE (WIF-137k); I; deciduous forest; *Tilia* sp. branch. **Kaczawskie Foothills:** Przeździec 1.7 km E (near WIF-12); X; tree clump near a stream; *Corylus avellana* branch.

***Hyphodontia sambuci* (Pers.) J. Erikss.**

Kaczawskie Mts.: Rząśnik 2.2 km W (WIF-128c); X; deciduous forest; *Sambucus nigra* branch. **Kaczawskie Foothills:** Bystrzyca 1.2 km SE (WIF-126o), Przeździec 1.1 km NW (WIF-24l), Przeździec 1.4 km E (WIF-12d); I, X; deciduous forest, riparian forest; *Sambucus nigra* branch.

***Hypholoma capnoides* (Fr.) P. Kumm. (= *Psilocybe capnoides* (Fr.) Noordel.)**

Izera Mts.: Świeradów-Zdrój (ŚF-452b); X; spruce forest; wood. **Kaczawskie Mts.:** Rząśnik 2.2 km W (WIF-128k), Wleń 1.2 km E (WIF-136c); X; deciduous forest, mixed forest; coniferous wood, ?*Picea abies* stump. **Izerskie Foothills:** Wleń 2 km SW (ŁuF – Gniazdo Mt), Wleń 2.2 km N (ŁuF – Gniazdo Mt), Wleń 2.4 km W (ŁuF-61i), Wleń 2.8 km N (WIF-44j); X-II; oak-hornbeam forest, spruce-beech forest, spruce forest, spruce-pine forest; coniferous wood, coniferous stump. **Kaczawskie Foothills:** Wąwóz Lipy Res.; XI; oak forest; wood.

***Hypholoma elongatum* (Pers.) Ricken (= *Psilocybe elongata* (Pers.) J.E. Lange)**

Izera Mts.: Świeradów-Zdrój (ŚF-361a), Sępik Mt (tourist trail between Szklarska Poręba and Hala Izerska), 0.4 km from Torfowiska Doliny Izery Res. (at the Izera river); IX-X; forest roadside, mossy place; mosses, soil. **Karkonosze Mts.:** Jakuszyce 0.9 km SE; X; roadside; *Sphagnum* sp. **Izerskie Foothills:** Zalesie 0.7 km N (LuF), Zalesie 0.8 km NE (LuF-199f); IX-X; mire; *Sphagnum* sp.

***Hypholoma ericaeoides* P.D. Orton (= *Psilocybe ericaeoides* (Orton) Noordel.)**

Kaczawskie Foothills: Bełczyna 1.1 km SE (WIF-123m); X; deciduous forest; soil.

***Hypholoma fasciculare* (Huds.) P. Kumm. var. *fasciculare* (= *Psilocybe fascicularis* (Huds.) Noordel.)**

Izerskie Foothills: Wleń 1.5 km SW (ŁuF – Gniazdo Mt); VI; wet spruce forest; wood. **Kaczawskie Foothills:** Wleń 1.8 km NE (WIF-28f), Wleń 2.3 km NE (WIF-25a); IV-V; oak-hornbeam forest, mixed forest; wood.

***Hypholoma fasciculare* (Huds.) P. Kumm. var. *subviride* (Berk. & M.A. Curtis) Krieglst. (= *Psilocybe subviridis* (Berk. & M.A. Curtis) Sacc.)**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); VII; riparian forest; wood. **Izerskie Foothills:** Wleń 1.5 km SW (ŁuF – Gniazdo Mt); IX; mixed forest; wood. **Kaczawskie Foothills:** Leszczyna 1 km N (near Ciche Szczęście mine), Radomiłowice 1 km SW (WIF-11b); IV-VI; oak-hornbeam forest, mixed forest; deciduous stump.

***Hypholoma lateritium* (Schaeff.) P. Kumm. (= *Psilocybe lateritia* (Schaeff.) Noordel.)**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Rząśnik 1.4 km SW (WIF-290k); IX-X; oak-hornbeam forest, deciduous forest; wood, deciduous stump. **Izerskie Foothills:** Wleń 1.4 km NW (WIF-46a); XI; deciduous sapling stand; wood.

***Hypholoma polytrichi* (Fr.) Ricken (= *Psilocybe polytrichi* (Fr.) Pears. & Dennis)**

Izerskie Foothills: Wleń 2.8 km N (WIF-44j), Zalesie 0.6 km N (LuF); IX-X; oak-hornbeam forest near a stream, coniferous forest; mosses, soil.

***Infundibulicybe gibba* (Pers.) Harmaja (= *Clitocybe gibba* (Pers.) P. Kumm.)**

Kaczawskie Foothills: Bystrzyca 1.6 km N (WIF-16h); VII; oak-hornbeam forest; litter.

***Inocutis rheades* (Pers.) Fiasson & Niemelä (= *Inonotus rheades* (Pers.) P. Karst.)**

Kaczawskie Mts.: Wleń 1.2 km SE (WIF-137d); IX; alder forest; *Alnus* sp. stump. **Izerskie Foothills:** Wleń 1.5 km SW (ŁuF – Gniazdo Mt); VIII; spruce forest; *Alnus* sp. log.

***Inocybe agardhii* (Lund) P.D. Orton**

Sudetes Foreland: Szklary 0.5 km W; IX; overgrown excavation; soil.

***Inocybe assimilata* Britzelm. (= *I. umbrina* Bres.)**

Kaczawskie Mts.: Bystrzyca 1.1 km S (WIF-133c); XI; beech forest; soil. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF-59d – Gniazdo Mt), Wleń 2 km SW (ŁuF – Gniazdo Mt); VIII-IX; spruce-beech forest, mixed forest; soil.

Kaczawskie Foothills: Bystrzyca 2.3 km NW (WIF-15p); XI; spruce forest; soil.

***Inocybe asterospora* Quél.**

Kaczawskie Foothills: Bystrzyca 1.2 km SE (WIF-126n); X; deciduous forest; soil.

***Inocybe brunneotomentosa* Huijsman**

Kaczawskie Mts.: Grudno (near quarry); IX; beech forest; soil.

***Inocybe calamistrata* (Fr.) Quél.**

Karkonosze Mts.: Karkonosze NP (Czeska Ścieżka); VIII; spruce forest; soil.

***Inocybe castanea* Peck**

Kaczawskie Foothills: Wąwóz Lipy Res.; XI; oak forest; soil.

***Inocybe cervicolor* (Pers.) Quél.**

Kaczawskie Foothills: Wleń 1.7 km NE (WIF-27h); VIII; coniferous forest; soil.

Inocybe cincinnata* (Fr.) Quél. var. *cincinnata

Izerskie Foothills: Łupki (near tourist hostel), Wleń 1.6 km SW (ŁuF-64g – Gniazdo Mt); IX-X; deciduous forest, lawn (under spruce and larch); soil. **Kaczawskie Foothills:** Bełczyna 1.8 km S (WIF-126j), Wleń 1.6 km NW (WIF-28f); IX-X; deciduous forest; soil.

***Inocybe cincinnata* (Fr.) Quél. var. *major* (S. Petersen) Kuyper**

Izera Mts.: Świeradów-Zdrój (ŚF-238i); X; spruce forest; soil. **Izerskie Foothills:** Wleń; IX-X; lawn (under fir); soil.

***Inocybe cookei* Bres.**

Kaczawskie Mts.: Stara Kraśnica 1.5 km SWW; X; deciduous forest; soil. **Kaczawskie Foothills:** Bełczyna 1.3 km SE (WIF-122d); VII; oak-hornbeam forest; soil.

***Inocybe curvipes* P. Karst.**

Izerskie Foothills: Łupki (near tourist hostel), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; oak-hornbeam forest; lawn (under spruce and larch), soil. **Sudetes Foreland:** Szklary 0.5 km SW; IX; overgrown excavation; soil.

Inocybe geophylla* (Fr.) P. Kumm. var. *geophylla

Kaczawskie Mts.: Mysłów 1 km E; X; spruce forest; soil. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.7 km SW (ŁuF – Gniazdo Mt), Wleń 2 km SW (ŁuF – Gniazdo Mt); VIII-XI; riparian forest, deciduous forest, mixed forest; soil. **Kaczawskie Foothills:** Bystrzyca 1.7 km N (WIF-18h); X; mixed forest; soil.

Inocybe lacera* (Fr.) P. Kumm. var. *lacera

Kaczawskie Mts.: Rząśnik 3 km SW (near WIF-135); VI; slate excavation; soil. **Karkonosze Mts.:** Szklarska Poręba (SzPF-347c), Karkonosze NP (Czeska Ścieżka); VIII-IX; spruce forest; soil. **Izerskie Foothills:** Radomice (limestone quarry); X; overgrown calcareous grassland; soil. **Kaczawskie Foothills:** Wleń 2.6 km N (WIF-25a); VIII; deciduous forest; soil.

***Inocybe leptophylla* G.F. Atk.**

Izerskie Foothills: Wleń 1.3 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest; soil.

***Inocybe leucoblema* Kühner**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry); VIII; overgrown calcareous grassland; soil.

***Inocybe lilacina* (Peck) Kauffman (= *I. geophylla* (Fr.) P. Kumm. var. *lilacina* (Peck) Gillet)**

Kaczawskie Foothills: Bełczyna 1.3 km SE (WIF-123t), Nad Groblą Res., Wąwóz Siedmicki Res.; IX-X; riparian forest, alder forest; soil.

***Inocybe maculata* Boud.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (near WIF-127a); IX; oak-hornbeam forest; soil. **Kaczawskie Foothills:** Bełczyna 1.7 km SE (WIF-124a); IX; deciduous forest; soil.

***Inocybe mixtilis* (Britzelm.) Sacc.**

Karkonosze Mts.: Szklarska Poręba (Turystyczna St.); X; lawn (under spruce), soil. **Izerskie Foothills:** Łupki (near tourist hostel); XI; lawn (under spruce and larch); soil. **Kaczawskie Foothills:** Bełczyna 0.7 km S (WIF-123g); X; young spruce forest; soil.

***Inocybe napipes* J.E. Lange**

Izera Mts.: Świeradów-Zdrój (ŚF-308f); X; spruce forest; soil. **Izerskie Foothills:** Wleń 2.3 km NW (WIF-46i); IX; mixed forest; soil.

***Inocybe petiginosa* (Fr.) Gillet**

Kaczawskie Mts.: Stara Kraśnica 1.5 km SWW; X; deciduous forest; soil. **Karkonosze Mts.:** Szklarska Poręba (Złoty Potok Valley); XI; beech forest (helocrene); soil.

***Inocybe proximella* P. Karst.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.9 km SW (ŁuF – Gniazdo Mt); VIII-IX; deciduous forest, mixed forest; soil.

***Inocybe pruinosa* R. Heim (= *I. halophila* Heim)**

Izerskie Foothills: Radomice (limestone quarry); V-VIII; overgrown calcareous grassland; soil.

***Inocybe quietiodor* Bon**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-59c – Gniazdo Mt); IX; mixed forest; soil.

***Inocybe relicina* (Fr.) Quél.**

Karkonosze Mts.: Karkonosze NP (Zielony Klin Mt); X; spruce forest; soil.

***Inocybe rimosa* (Bull.) P. Kumm. (= *I. fastigiata* (Schaeff.) Quél.)**

Izerskie Foothills: Wleń 1.3 km NW (WIF-47w); VIII; deciduous forest; soil.

***Inocybe sindonia* (Fr.) P. Karst.**

Izera Mts.: Świeradów-Zdrój (ŚF-238k); X; spruce forest; soil. **Karkonosze Mts.:** Szklarska Poręba (Turystyczna Mt); X; lawn (under spruce and larch); soil. **Kaczawskie Foothills:** Przeździec 0.8 km SW (WIF-221); IX; deciduous forest; soil.

***Inocybe soluta* Velen.**

Karkonosze Mts.: Szklarska Poręba (Złoty Potok Valley); XI; spruce forest, beech forest; soil. **Izerskie Foothills:** Wleń 2 km SW (ŁuF – Gniazdo Mt); VIII; mixed forest; soil.

***Inocybe terrigena* (Fr.) Kühner**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry); VIII-X; overgrown calcareous grassland; soil.

***Inonotus hispidus* (Bull.) P. Karst.**

Kaczawskie Foothills: Rzańnik (side towards Bełczyna); X; roadside; *Fraxinus* sp. trunk.

***Inonotus nodulosus* (Fr.) P. Karst.**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt); X; mixed forest; *Fagus sylvatica* log.

***Inonotus obliquus* (Ach. ex Pers.) Pilát**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF- 127a), Tarczyn 0.6 km E (near WIF-134h), Wleń 1.1 km E (near WIF-136 & 137), Wleń 1.2 km E (WIF-136i), Wleń 1.3 km NE (WIF-136b), Wleń 1.8 km SE (WIF-137l); I-XI; oak-hornbeam forest, forest, deciduous forest, mixed forest, roadside; *Betula* sp. trunk, *Alnus incana* trunk.

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2.1 km N (WIF-44b); VIII-XI; deciduous forest; *Betula* sp. log and trunk. **Kaczawskie Foothills:** Bełczyna 0.8 km N (WIF-122l), Bełczyna 1.8 km SE (WIF-124d), Bystrzyca 1.2 km SE (WIF-126n), Bystrzyca 1.4 km N (WIF-19b), Bystrzyca 2.4 km NW (WIF-22/26), Bystrzyca 2.5 km NW (WIF-26f), Bystrzyca 2.8 km NW (WIF-22p), Przeździec 0.8 km SE (WIF-21n), Przeździec 1.4 km E (WIF-12l, 12m), Sobota 1.6 km SW (WIF-24Aa), Sobota 1.7 km SW (WIF-24Ab & 24Ag), Sobota 2.4 km S (WIF-24Aj), Wleń 1.4 km N (WIF-32c); I-XII; coniferous forest, deciduous forest, mixed forest, riparian forest, mixed wood margin; *Betula* sp. log and trunk, *Alnus incana* log and trunk, *Alnus* sp. trunk.

***Inonotus radiatus* (Sowerby) P. Karst.**

Kaczawskie Mts.: Buki Sudeckie Res., Bystrzyca 1 km S (WIF-133a); X-XI; *Carici remotae-Fraxinetum*, riparian forest; *Alnus* sp. trunk and stump.

***Irpex lacteus* (Fr.) Fr.**

Kaczawskie Foothills: Przeździec 1.7 km E (near WIF-12); X; tree clump near a stream; deciduous branch.

***Ischnoderma benzoinum* (Wahlenb.) P. Karst.**

Izerskie Foothills: Wleń 1.5 km SW (ŁuF – Gniazdo Mt), Zalesie 0.6 km N (LuF); IX-X; coniferous forest, mixed forest; spruce stump, *Picea abies* log. **Kaczawskie Foothills:** Bełczyna 1.8 km SE (WIF-124d); II; mixed forest; coniferous log.

***Ischnoderma resinosum* (Schr.) P. Karst.**

Izerskie Foothills: Lubomierz 2 km W (LuF), Wleń 2.1 km SW (ŁuF – Gniazdo Mt); X-XI; riparian forest, tree clump near a pond; *Fagus sylvatica* trunk, *Alnus incana* trunk. **Kaczawskie Foothills:** Bystrzyca 2.5 km NE (WIF-15g, 15h), Ostrzyca Proboszczowiecka Res.; X; deciduous forest; deciduous log, ?*Tilia* sp. log, ?*Betula* sp. stump.

***Junghuhnia luteoalba* (P. Karst.) Ryvarden (= *Irpex luteoalbus* (P. Karst.) Kotiranta & Saarenoksa)**

Kaczawskie Foothills: Wleń 1.1 km N (WIF-31k); I; deciduous forest; *Sorbus aucuparia* branch.

***Junghuhnia nitida* (Pers.) Ryvarden (= *Irpex nitidus* (Pers.) Saarenoksa & Kotiranta)**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest; deciduous branch.

***Kuehneromyces mutabilis* (Schaeff.) Singer & A.H. Sm. (= *Pholiota mutabilis* (Scop.) P. Kumm.)**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt), Wleń 2.1 km SW (ŁuF – Gniazdo Mt); VII-XI; spruce-beech forest, mixed forest, alder forest; wood. **Kaczawskie Foothills:** Wleń 1.9 km NE (WIF-27d); VIII; deciduous forest; wood.

***Laccaria amethystina* Cooke**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); oak-hornbeam forest; soil. **Karkonosze Mts.:** Szklarska Poręba (Złoty Potok Valley); XI; beech forest (helocrene); soil. **Kaczawskie Foothills:** Bystrzyca 2.1 km N (WIF-14h); VIII; mixed forest; soil.

***Laccaria laccata* (Scop.) Cooke var. *pallidifolia* (Peck) Peck**

Karkonosze Mts.: Szklarska Poręba (Złoty Potok Valley); XI; spruce forest (helocrene); soil, *Sphagnum* sp.

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt – Białe Skąły), Wleń 2.4 km NW (ŁuF-48a), Zalesie 0.7 km N (LuF); IX; near a stream, mire, forest margin (under aspen); soil, *Sphagnum* sp. **Kaczawskie Foothills:** Wleń 2.3 km NE (WIF-25c); IX; mixed forest; soil.

***Laccaria proxima* (Boud.) Pat.**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt); IX; spruce forest near a stream; soil. **Kaczawskie Mts.:** Tarczyn 0.6 km E (Tarczynka Mt); X; overgrown slate excavation; soil.

***Laccaria pumila* Fayod**

Kaczawskie Foothills: Bełczyna 1.8 km SE (WIF-124d), Bystrzyca 0.5 km N (WIF-20I); IX-X; riparian forest, basalt quarry; soil.

***Laccaria purpureobadia* D.A. Reid**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.7 km SW (ŁuF – Gniazdo Mt), Zalesie 0.8 km NE (LuF-199f); IX-I; spruce forest, mire; soil, *Sphagnum* sp.

***Laccaria tortilis* (Bolton) Cooke**

Kaczawskie Mts.: Wleń 1.1 km E (WIF-136g); IX; forest; soil.

***Lacrymaria lacrymabunda* (Bull.) Pat.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-59b – Gniazdo Mt); X; riparian forest; soil. **Kaczawskie Foothills:** Wleń 2.6 km N (WIF-25a); IX-X; log landing; litter. **Sudetes Foreland:** Szklary 0.3 km S; IX; floodplain (under alder); wood.

***Lacrymaria pyrotricha* (Holmsk.) Konrad & Maubl.**

Kaczawskie Foothills: Bystrzyca 1.4 km N (WIF-17c); X; riparian forest; soil.

***Lactarius aurantiofulvus* J. Blum ex Bon**

Kaczawskie Foothills: Bystrzyca 2.2 km E (WIF-126n); X; spruce forest; soil.

Lactarius azonites* (Bull.) Fr. f. *azonites

Kaczawskie Foothills: Wleń 2 km N (WIF-30f); IX; under lime tree; soil.

***Lactarius blennius* (Fr.) Fr.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; beech forest; soil.

***Lactarius camphoratus* (Bull.) Fr.**

Kaczawskie Mts.: Stara Kraśnica 1.5 km SWW; X; deciduous forest; soil. **Kaczawskie Foothills:** Wleń 2 km N (WIF-29g); VIII; spruce forest; soil.

***Lactarius chrysorrheus* Fr.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); IX; oak-hornbeam forest; soil. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); VIII; mixed forest; soil.

***Lactarius controversus* Pers.**

Kaczawskie Foothills: Belczyna 0.8 km S (WIF-123k); IX; spruce forest (under poplar); soil.

***Lactarius cyathuliformis* Bon**

Kaczawskie Foothills: Sobota 1.6 km SW (WIF-24Aa), Nad Groblą Res.; VIII; riparian forest, alder sapling stand; soil.

***Lactarius deterrimus* Gröger**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; spruce forest; soil. **Kaczawskie Foothills:** Belczyna 0.7 km S (WIF-123g), Belczyna 0.8 km S (near WIF-123); IX-X; spruce forest, overgrown pasture; soil.

***Lactarius fulvissimus* Romagn. (= *L. ichoratus* (Batsch) Fr.)**

Kaczawskie Foothills: Belczyna 0.7 km S (WIF-123g); XI; spruce forest; soil.

***Lactarius glyciosmus* (Fr.) Fr.**

Kaczawskie Mts.: Tarczyn 0.6 km E (Tarczynka Mt), Tarczyn 1.8 km E (near WIF-133), Stara Kraśnica 2 km S; X; meadow, meadow with birch-poplar brushwoods, overgrown slate excavation; soil. **Izerskie Foothills:** Zalesie 0.8 km NE (LuF-199g); IX-X; mire; soil, *Sphagnum* sp.

***Lactarius helvus* (Fr.) Fr.**

Izerskie Foothills: Zalesie 0.8 km N (LuF); IX; mire; soil.

***Lactarius leonis* Kytöv.**

Izerskie Foothills: Wleń 1.6 km SW (LuF – Gniazdo Mt); IX; wet spruce forest; soil.

***Lactarius lignyotus* Fr.**

Kaczawskie Mts.: Bystrzyca 1.5 km S (WIF-133c); VIII; spruce forest; soil. **Karkonosze Mts.:** Karkonosze NP (Łabski Kocioł); IX; spruce forest, soil.

***Lactarius necator* (Bull.) Pers.**

Kaczawskie Mts.: Tarczyn 1.2 km E (WIF-133f); X; coniferous forest; soil.

***Lactarius obscuratus* (Lasch) Fr.**

Izerskie Foothills: Wleń 1.4 km NW (WIF-46a); IX; wet deciduous sapling stand; soil.

***Lactarius pubescens* (Schräd.) Fr.**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry); IX; overgrown limestone quarry (under birch); soil.
Sudetes Foreland: Szklary 0.5 km W; IX; overgrown spoil tip; soil.

***Lactarius pyrogalus* (Bull.) Fr.**

Kaczawskie Foothills: Bystrzyca 1.3 km SE (WIF-126k); IX; deciduous forest; soil.

***Lactarius scrobiculatus* (Scop.) Fr.**

Izerskie Foothills: Wleń 1.6 km SW (LuF – Gniazdo Mt), Wleń 1.7 km SW (LuF – Gniazdo Mt); IX-XI; spruce forest; soil.

***Lactarius serifluus* (DC.) Fr.**

Kaczawskie Foothills: Wleń 2 km N (WIF-30f); VIII; under lime tree; soil.

***Lactarius spinulosus* Qué. & Le Bret.**

Kaczawskie Foothills: Bełczyna 0.7 km S (WIF-123g), Bełczyna 1 km E (near WIF-122); X-XI; spruce forest, deciduous tree clump; soil.

***Lactarius tabidus* Fr. (= *L. thejogalus* (Bull.) Gray ss. Neuhoff)**

Karkonosze Mts.: Karkonosze NP (Czeska Ścieżka); VIII; spruce forest; soil. **Izerskie Foothills:** Wleń 1.7 km SW (LuF – Gniazdo Mt), Zalesie 0.8 km N (LuF); VIII-IX; mire margin, spruce forest; soil.

***Lactarius torminosus* (Schaeff.) Pers.**

Kaczawskie Foothills: near Nad Groblą Res.; IX; near a stream (under pine and birch); soil.

***Lactarius trivialis* (Fr.) Fr.**

Izerskie Foothills: Zalesie 0.6 km N (LuF); IX; spruce-pine forest; soil.

***Lactarius vietus* (Fr.) Fr.**

Izerskie Foothills: Zalesie 0.8 km N (LuF); IX; mire margin; soil.

***Lactarius volemus* (Fr.) Fr.**

Kaczawskie Foothills: Belczyna 0.7 km SE (WIF-122g), Wleń 2 km N (WIF-30f); VII; oak-hornbeam forest, under lime tree and oak; soil.

***Laetiporus sulphureus* (Bull.) Murrill**

Kaczawskie Mts.: Bystrzyca 0.9 km S (WIF-127a); VIII; riparian forest; *Quercus* sp. stump. **Kaczawskie Foothills:** Belczyna 0.6 km S (WIF-123f), Bystrzyca 2.5 km NE (WIF-15h), Radomiłowice 1 km SW (WIF-11c); V-VIII; deciduous forest, mixed forest; *Quercus* sp. wood, deciduous stump, *Prunus avium* trunk.

***Langermannia gigantea* (Batsch) Rostk.**

Izerskie Foothills: Wleń (Południowe Es.), Wleń (ŁuF-58m); VIII; garden, forest margin; soil. **Kaczawskie Foothills:** Belczyna (E part of the village), Wleń (Kasprowicza St.); III-VIII; tree clump; soil. **Sudetes Foreland:** Szklary (S from the village); III-IV; aspen forest; soil.

***Laxitextum bicolor* (Pers.) Lentz**

Kaczawskie Mts.: Wleń 1.7 km SE (WIF-137o); I; lime forest; *Tilia* sp. branch. **Izerskie Foothills:** Wleń 2 km SW (ŁuF – Gniazdo Mt); X; mixed forest; *Fagus sylvatica* branch. **Kaczawskie Foothills:** Belczyna 0.8 km E (near WIF-122), Bystrzyca 2.2 km N (WIF-15d), Wleń 1.9 km NE (WIF-27d), Wleń 2.2 km N (WIF-29c); VIII-IX; deciduous forest, mixed forest, riparian forest, spruce forest; *Alnus incana* branch, *Fagus sylvatica* branch, *Salix* sp. log, wood.

***Leccinum aurantiacum* (Bull.) Gray**

Izerskie Foothills: Wleń 1.6 km NW (WIF-46f); VIII; spruce forest (under aspen); soil.

***Leccinum niveum* (Fr.) Rauschert**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Zalesie 0.8 km NE (LuF-199f, 199g); VII, IX; spruce forest (under birch), overgrown mire; soil.

***Leccinum pseudoscabrum* (Kallenb.) Šutara**

Kaczawskie Mts.: Bystrzyca (WIF-134x); VII; mixed forest; soil. **Kaczawskie Foothills:** Przeździec 1.3 km E (near WIF-12); VII; deciduous forest; soil.

***Leccinum quercinum* (Pilát) E.E. Green & Watling**

Kaczawskie Foothills: Belczyna 1.8 km SE (WIF-124a); IX; *Quercus* sp. forest; soil.

***Leccinum scabrum* (Bull.) Gray**

Izerskie Foothills: Wleń 1.5 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF-59c – Gniazdo Mt); IX-X; spruce forest; soil. **Kaczawskie Foothills:** Radomiłowice 0.9 km W (WIF-12Ac), N from Nad Groblą Res.; IX; mixed forest; soil.

***Leccinum schistophilum* Bon**

Izerskie Foothills: Zalesie 0.8 km N (LuF-200h); IX; overgrown mire; soil.

***Leccinum variicolor* Watling**

Izerskie Foothills: Zalesie 0.6 km N (LuF); IX; pine forest; soil. **Kaczawskie Foothills:** N from Nad Groblą Res.; IX; mixed forest; soil.

Lentinellus cochleatus* (Pers.) P. Karst. var. *cochleatus

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; spruce forest; *Betula* sp. stump. **Kaczawskie Foothills:** Bystrzyca 1.3 km SE (WIF-126k), Bystrzyca 1.5 km E (WIF-126d), Bystrzyca 2.2 km N (WIF-14b); IX; deciduous forest, mixed forest; wood, *Alnus* sp. stump.

***Lentinellus cochleatus* (Pers.) P. Karst. var. *inolens* (Konrad & Maubl.) Kühner & Romagn.**

Kaczawskie Mts.: Wleń 1.3 km NE (WIF-136b); VIII; mixed forest; stump.

***Lentinellus flabelliformis* (Bolton) S. Ito s.l.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); X; deciduous forest; *Salix* sp. wood. **Kaczawskie Foothills:** Bystrzyca 1.6 km N (WIF-19d), Bystrzyca 1.9 km N (WIF-18c); IX-X; deciduous sapling stand, oak-hornbeam forest; wood, ?*Fraxinus* sp. twigs.

***Lenzites betulina* (L.) Fr.**

Kaczawskie Mts.: Wleń 1.9 km E (WIF-136a); IV; spruce forest margin; *Betula* sp. log. **Kaczawskie Foothills:** Belczyna 1 km SE (WIF-122/124); Przeździec 1.7 km E (WIF-12n), Radomiłowice 1 km SW (WIF-11b), Wleń 1.4 km NW (WIF-31b), Wleń 1.5 km NW (near WIF-31); I, IX-XII; riverside vegetation, mixed forest, overgrown wastelands; *Betula* sp. trunk and branch, deciduous log, *Salix* sp. branch, ?*Quercus* sp. stump.

***Lepiota apatelia* Vellinga & Huijser**

Kaczawskie Foothills: Wleń 1.5 km NW (near WIF-31), Wleń 2.3 km NE (WIF-26m); VII-VIII; forest margin, mixed tree clump near a stream; soil.

***Lepiota boudierii* Bres. (= *L. fulvella* Rea)**

Kaczawskie Mts.: Góra Milek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Lepiota castanea* Quél.**

Kaczawskie Mts.: Mysłów 2 km E, Góra Miłek Res.; IX-X; *Cephalanthero-Fagetum*, deciduous forest; soil.

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); IX; oak-hornbeam forest; soil. **Kaczawskie Foothills:** Belczyna 1.8 km S (WIF-126j), Bystrzyca 1.2 km SE (WIF-126n), Wleń 1.5 km N (WIF-30f); VIII-X; deciduous forest, mixed forest, riparian forest; soil.

***Lepiota clypeolaria* (Bull.) Kumm.**

Kaczawskie Mts.: Góra Miłek Res.; VIII; *Cephalanthero-Fagetum*; soil. **Izerskie Foothills:** Wleń 1.3 km NW (WIF-47a), Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.7 km SW (ŁuF – Gniazdo Mt), Wleń 1.7 km NW (WIF-46j), Wleń 1.9 km NW (WIF-46d), Wleń 2 km SW (ŁuF – Gniazdo Mt), Wleń 2.4 km NW (near WIF-48a), Wleń 2.8 km N (WIF-44j); VIII-X; oak-hornbeam forest, spruce forest, under spruce near a stream, deciduous forest, mixed forest; soil. **Kaczawskie Foothills:** Belczyna 1 km SE (WIF-122f), Belczyna 1.2 km SE (WIF-122i), Belczyna 1.8 km SE (WIF-124a), Bystrzyca 1.5 km E (WIF-126d), Bystrzyca 1.5 km N (WIF-16k), Bystrzyca 2.2 km N (WIF-13d, 15c), Bystrzyca 2.5 km NW (WIF-15h), Górczyca 1.4 km SW (WIF-24h), Radzimowice 1.5 km S (WIF-13), Wleń 1.5 km N (WIF-30f), Wleń 2.3 km NE (WIF-25d), Wleń 2.5 km N (WIF-25a), Wleń 3 km NE (WIF-22p); VIII-XI; birch forest, oak-hornbeam forest, between beech forest and riparian forest, deciduous forest, mixed forest; soil.

***Lepiota cristata* (Bolton) P. Kumm.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1 km S (WIF-133a), Nowe Rochowice, Góra Miłek Res., Miłek Mt; VIII-X; oak-hornbeam forest, riparian forest, mixed forest, *Cephalanthero-Fagetum*, overgrown calcareous grassland; soil. **Izerskie Foothills:** Wleń, Wleń 1.4 km SW (ŁuF-59b – Gniazdo Mt); IX-X; riparian forest, lawn; soil. **Kaczawskie Foothills:** Wleń 2 km N (WIF-30b), Wleń 2.1 km N (WIF-30c), Wleń 2.6 km N (WIF-25b), Wleń 2.5 km N, N od Nad Groblą Res.; VIII-XI; deciduous forest, nitrophilous vegetation, oak forest, riparian forest; soil. **Sudetes Foreland:** Szklary 0.5 km SW; IX; between alder forest and oak forest; soil.

***Lepiota echinella* (Quél.) G.E. Bernard var. *echinella* (= *L. setulosa* J.E. Lange)**

Kaczawskie Foothills: Bystrzyca 1.4 km N (WIF-17h); IX; oak-hornbeam forest; soil.

***Lepiota felina* (Pers.) P. Karst.**

Kaczawskie Mts.: Tarczyn 0.9 km NE (WIF-134c); X; spruce forest; soil. **Kaczawskie Foothills:** Belczyna 0.6-0.7 km S (WIF-123g); IX-X; spruce forest, coniferous forest; soil.

***Lepiota magnispora* Murrill (= *L. ventriosospora* Reid)**

Izerskie Foothills: Wleń 1.1 km SW (ŁuF-58m), Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2 km SW (ŁuF – Gniazdo Mt), Wleń 2.3 km NW (WIF-45i); IX-X; beech forest, mixed forest; soil. **Kaczawskie Foothills:** Belczyna 0.5 km SW (WIF-123g), Belczyna 0.7 km S (WIF-123g), Belczyna 1 km SE (WIF-122f), Bystrzyca 2.2 km N (WIF-14h, 15b), Wleń 1.4 km S (WIF-31c); VIII-X; mixed forest, pine forest, spruce forest; soil. **Sudetes Foreland:** Szklary 0.5 km SW; IX; overgrown excavation; soil.

***Lepiota oreadiformis* Velen.**

Kaczawskie Foothills: Radomiłowice 0.8 km W (near WIF-12Ar); IX; meadow; soil.

***Lepiota pseudolilacea* Huijsman (= *L. pseudohelveola* Kühner ex Hora)**

Sudetes Foreland: Szklary 0.5 km SW; IX; overgrown excavation; soil.

***Lepiota subalba* P.D. Orton**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); IX; oak-hornbeam forest; soil.

***Lepista nuda* (Bull.) Cooke**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); X; deciduous forest; soil. **Kaczawskie Foothills:** Bystrzyca 1.5 km E (WIF-126i), Przeździec 1.4 km E (WIF-12i); X; meadow near a forest, forest; soil.

***Lepista saeva* (Fr.) P.D. Orton (= *L. personata* (Fr.) Cooke)**

Kaczawskie Foothills: Wleń 1.7 km NE (WIF-28d); X; mixed forest (roadside); soil.

***Lepista sordida* (Schumach.) Singer**

Kaczawskie Foothills: Sobota 1.5 km S, Sobota 2.6 km S; X-XI; pasture; *Equus caballus* manure, soil.

***Leptosporomyces mutabilis* (Bres.) Krieglst. (= *Fibulomyces mutabilis* (Bres.) Jülich)**

Izerskie Foothills: Wleń 1.5 km SW (ŁuF – Gniazdo Mt); XI; spruce forest; litter, *Larix* sp. tree base.

***Leratiomyces squamosus* (Pers.) Bridge & Spooner var. *squamosus* (= *Psilocybe squamosa* (Pers.) P.D. Orton var. *squamosa*)**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a), Wleń 1.3 km E (WIF-136b); X; oak forest, riparian forest; litter. **Izerskie Foothills:** Wleń 2.8 km N (WIF-44j); X; oak-hornbeam forest; litter. **Kaczawskie Foothills:** Belczyna 1 km E (WIF-126b, 126s), Bystrzyca 2.2 km N (WIF-13d), Przeździec 0.8 km SW (WIF-22l); IX-X; deciduous forest, mixed forest; litter, soil.

***Leratiomyces squamosus* (Pers.) Bridge & Spooner var. *thraustus* (Kalchbr.) Bridge & Spooner (= *Psilocybe squamosa* (Pers.) P.D. Orton var. *thrausta* (S. Schulz.) Guzmán)**

Kaczawskie Mts.: Wleń 1.3 km NE (WIF-136c); IX-X; spruce forest; litter. **Kaczawskie Foothills:** Bystrzyca 2 km W (WIF-126w), Wleń 2.5 km N (WIF-25a); IX-X; beech forest, oak-hornbeam forest; litter.

***Leucoagaricus leucothites* (Vittad.) Wasser var. *leucothites* (= *L. cretaceus* (Bull.) M.M. Moser)**

Izerskie Foothills: Łupki (near tourist hostel); IX; lawn (near spruce and larch); soil.

***Leucoagaricus nymphaeum* (Kalchbr.) Bon (= *Macrolepiota nymphaeum* (Kalchbr.) Wasser)**

Izerskie Foothills: Wleń 2.1 km SW (ŁuF – Gniazdo Mt); IX; spruce forest; soil. **Kaczawskie Foothills:** Wleń 2.1 km N (WIF-22l), Wleń 3 km NE (WIF-22r); VIII; mixed forest; soil.

Leucoagaricus sericifer* (Locq.) Vellinga f. *sericifer

Kaczawskie Foothills: Wleń 2.1 km N (WIF-30c); IX; deciduous sapling stand; soil.

***Lindtneria panphyliensis* Bernicchia & M.J. Larsen**

Kaczawskie Foothills: Przeździec 0.5 km NE (WIF-24z); XII; oak-hornbeam forest; ?*Tilia* sp. log.

***Lopharia spadicea* (Pers.) Boidin (= *Porostereum spadiceum* (Pers.) Hjortstam & Ryvarden)**

Kaczawskie Mts.: Tarczyn 0.6 km E (Tarczynka Mt); I; tree clump; *Populus tremula* log.

***Lycoperdon echinatum* Pers.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Grudno (Bukowa G.); IX; *Cephalanthero-Fagetum*, forest; soil. **Kaczawskie Foothills:** Belczyna 1 km SE (WIF-122f), Bystrzyca 1.3 km SE (WIF-126h), Bystrzyca 1.4 km N (WIF-17a), Bystrzyca 1.5 km E (WIF-126d), Bystrzyca 1.8 km N (WIF-16a), Bystrzyca 1.9 km N (WIF-13g); VIII-X; oak-hornbeam forest, deciduous forest, beech forest; *Betula* sp. stump, soil.

***Lycoperdon molle* Pers.**

Kaczawskie Foothills: Sobota 1.5 km S; X; pasture; soil.

***Lycoperdon nigrescens* Pers.**

Kaczawskie Mts.: Tarczyn 0.6 km E (Tarczynka Mt, slate quarry); X; overgrown quarry; soil. **Kaczawskie Foothills:** Wleń 2 km NE (WIF-27b); X; pine-larch forest; soil.

***Lycoperdon perlatum* Pers.**

Kaczawskie Foothills: Belczyna 1.1 km SE (WIF-123m); X; deciduous forest; soil.

***Lycoperdon pratense* Pers. (= *Vascellum pratense* (Pers.) Kreisel)**

Kaczawskie Mts.: Rząśnik 3 km SW (slate quarry); X; overgrown quarry; soil. **Kaczawskie Foothills:** Górczyca 1.5 km W (by the Bóbr river near WIF-24); X; meadow; soil.

***Lycoperdon utriforme* Bull. (= *Calvatia utriformis* (Bull.) Jaap)**

Kaczawskie Mts.: Bystrzyca 0.7 km S (near WIF-127a); VIII; meadow; soil. **Kaczawskie Foothills:** Belczyna 1 km S (near WIF-123n); X; meadow; soil.

***Lyophyllum anthracophilum* (Lasch) M. Lange & Sivertsen**

Kaczawskie Foothills: Belczyna 0.7 km S (WIF-123g), Wleń 1.7 km NE (WIF-27h); X-I; spruce-pine forest (old campfire), spruce forest (old campfire); charcoal.

***Lyophyllum palustre* (Peck) Singer**

Izerskie Foothills: Zalesie 0.6-0.8 km N (LuF), Zalesie 0.8 km NE (LuF-199g, 199j); VII-IX; mire, overgrown mire (mixed forest); *Sphagnum* sp.

***Lyophyllum rancidum* (Fr.) Singer**

Kaczawskie Foothills: Bystrzyca 0.7 km E (WIF-20ax), Bystrzyca 2.1 km E (WIF-125f); X; deciduous tree clump, mixed forest; soil.

Macrocystidia cucumis* (Pers.) Joss. var. *cucumis

Kaczawskie Foothills: Bystrzyca 1.4 km N (WIF-17a), Bystrzyca 1.4 km N (WIF-17c), Bystrzyca 2 km NE (WIF-15a); X-XI; oak-hornbeam forest, riparian forest; soil.

***Macrolepiota excoriata* (Schaeff.) Wasser**

Kaczawskie Foothills: Bystrzyca 2.4 km NW (WIF-22); IX; mixed forest; soil.

***Macrolepiota fuliginosa* (Barla) Bon**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; mixed forest; soil. **Kaczawskie Foothills:** Bystrzyca 0.5 km N (near WIF-20j); X; forest; soil.

***Macrolepiota mastoidea* (Fries) Singer s.l.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; oak-hornbeam forest, mixed forest; soil.

Kaczawskie Foothills: Bystrzyca (boarder of WIF-20y), Górczyca 1.4 km W (WIF-24d); VIII-X; mixed forest, meadow; soil.

***Macrotyphula filiformis* (Bull.) Rauschert (= *Clavariadelphus junceus* Alb. & Schwein.) Corner**

Kaczawskie Mts.: Rząśnik 2.3 km W (WIF-128c), Wleń 0.8 km NE (WIF-136f); X; deciduous forest; litter.

Kaczawskie Foothills: Bełczyna 0.7 km SW (WIF-123r), Bystrzyca 2.5 km NW (WIF-15h), Przeździec (near WIF-24n), Radomiłowice 0.9 km SW (WIF-11b), Wleń 2.2 km N (WIF-25c); X; oak forest, oak-hornbeam forest, deciduous forest, mixed forest; litter.

***Macrotyphula fistulosa* (Holmsk.) R.H. Petersen var. *fistulosa* (= *Clavariadelphus fistulosus* (Holmsk.) Corner var. *fistulosus*)**

Kaczawskie Mts.: Bystrzyca 0.5 km E (near WIF-134j), Bystrzyca 0.5 km SE (WIF-134w); XI; deciduous forest; litter. **Kaczawskie Foothills:** Bełczyna 0.7 km SW (WIF-123r), Bystrzyca 0.7 km E (WIF-20ax), Bystrzyca 1.2 km SE (WIF-126n), Bystrzyca 1.9 km W (WIF-126n), Bystrzyca 2.5 km NW (WIF-15h, 26k), Górczyca 0.9 km E (WIF-12Ao), Radomiłowice 1.2 km W (WIF-12Ac); X-XI; coniferous forest, deciduous forest, mixed forest, riparian forest; litter, *Corylus avellana* branch.

***Marasmiellus perforans* (Hoffm.) Antonín, Halling & Noordel.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; spruce forest; litter.

Marasmiellus ramealis* (Bull.) Singer var. *ramealis

Izerskie Foothills: Wleń 2.2 km NW (WIF-46c), Wleń 2.8 km N (WIF-44j); VII, X; oak-hornbeam forest, pine forest; twigs, *Picea abies* wood. **Kaczawskie Foothills:** Nad Groblą Res.; VIII; oak forest; wood.

Marasmiellus tricolor* (Alb. & Schwein.) Singer var. *tricolor

Kaczawskie Foothills: Wleń 2.8 km N (near WIF-25b); VII; meadow; litter.

***Marasmiellus vaillantii* (Pers.) Singer**

Kaczawskie Foothills: Przeździec 0.8 km SE (near WIF-21n); VIII; deciduous sapling stand; litter.

***Marasmius bulliardii* Quél.**

Kaczawskie Foothills: Ostrzyca Proboszczowiecka Res.; X; deciduous forest; decaying leaves. **Sudetes**

Foreland: Szklary 0.3 km S; IX; alder forest; decaying leaves.

***Marasmius cohaerens* (Pers.) Cooke & Quél.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1.2 km SE (WIF-131d); VIII; between riparian forest and oak-hornbeam forest, mixed forest; litter. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII-IX; riparian forest, deciduous forest; litter. **Kaczawskie Foothills:** near Nad Groblą Res.; IX; riparian forest; litter.

***Marasmius curreyi* Berk. & Broome**

Izerskie Foothills: Wleń 2.3 km NW (WIF-46c); VII; field near a forest; decaying grasses. **Kaczawskie**

Foothills: Radomiłowice 1.2 km S (WIF-14b), Wleń 2.6 km N (WIF-25a); VIII; forest margin; decaying grasses.

***Marasmius epiphyllus* (Pers.) Fr.**

Kaczawskie Foothills: Belczyna 1.1 km SE (WIF-123m), Bystrzyca 1.2 km SE (WIF-126n), Bystrzyca 1.9 km N (WIF-18a), Radomiłowice 0.9 km W (WIF-12Ac); X; deciduous forest, mixed forest; decaying leaves, decaying *Quercus* sp. leaves.

***Marasmius oreades* (Bolton) Fr.**

Kaczawskie Foothills: Sobota 1.5 km S; X; pasture; soil.

***Marasmius rotula* (Scop.) Fr.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; twigs. **Izerskie Foothills:** Wleń 2.7 km NW (WIF-44k); VI; riparian forest; twigs. **Kaczawskie Foothills:** Bystrzyca 1.2 km SE (WIF-126r), near Wąwóz Lipy Res.; VII; riparian forest, alder forest; deciduous twigs.

***Marasmius setosus* (Sowerby) Noordel.**

Kaczawskie Foothills: Belczyna 1 km SE (WIF-122f), Bystrzyca 2.5 km NW (WIF-15k); X; beech forest, deciduous sapling stand; decaying *Fagus sylvatica* leaves.

***Marasmius torquescens* Quél.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.5 km SW (ŁuF – Gniazdo Mt); VIII; mixed forest; litter. **Kaczawskie Foothills:** Bystrzyca 1.4 km N (WIF-17f), Nad Groblą Res.; VIII-X; mixed forest, riparian forest; litter.

***Marasmius wettsteinii* Sacc. & P. Syd.**

Kaczawskie Foothills: Belczyna 0.7 km S (WIF-123h), Bystrzyca 1.7 km N (WIF-16g); VII; spruce forest; litter.

***Marasmius wynneae* Berk. & Broome**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt); X; beech forest; litter. **Kaczawskie Foothills:** Nad Groblą Res.; X; riparian forest; litter.

***Melanoleuca friesii* (Bres.) Bon**

Izerskie Foothills: Wleń (Południowe Es.); VII; lawn; soil.

***Melanoleuca subpulverulenta* (Pers.) Métrod**

Izerskie Foothills: Wleń 2.7 km NW (WIF-44k); X; riparian forest; soil.

***Melanophyllum haematospermum* (Bull.) Kreisel**

Kaczawskie Foothills: Wleń 2.1 km N (WIF-30c); IX; deciduous sapling stand; soil.

***Meripilus giganteus* (Pers.) P. Karst.**

Kaczawskie Mts.: Bystrzyca 1.2 km S (WIF-133b), Wleń 1.1 km E, Wleń 1.3 km NE (WIF-136b, 136c), Wleń 1.1 km E (by the road to Tarczyn); VII-IX; mixed forest, roadside, spruce forest; *Fagus sylvatica* trunk base, deciduous stump, ?*Quercus* sp. stump. **Kaczawskie Foothills:** Bystrzyca 1.2 km SE (WIF-126r), Bystrzyca 1.4 km N (WIF-17a), Bystrzyca 1.4 km N (WIF-17a), Bystrzyca 1.7 km N (WIF-13i), Bystrzyca 2 km S (WIF-126w), Rząśnik 1.7 km towards Belczyny, między Rząśnikiem a Belczyną (near WIF-125), Radomiłowice 1.1 km SW (WIF-11g); VIII- X; oak-hornbeam forest, riparian forest, deciduous forest, mixed forest, roadside; deciduous wood, deciduous stump, *Fraxinus* sp. stump.

***Merismodes anomala* (Pers.) Singer**

Kaczawskie Foothills: Wleń 2.1 km N (WIF-30c); I; deciduous sapling stand; wood.

***Merismodes confusa* (Bres.) D.A. Reid**

Izerskie Foothills: Radomice (limestone quarry), Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.9 km SW; XI-II; riparian forest, overgrown quarry, tree clump; *Corylus avellana* branch, *Salix* sp. branch. **Kaczawskie Foothills:** Belczyna 0.9 km E (near WIF-122), Bystrzyca 2.6 km NW (near WIF-21), Górczyca 1.5 km W (WIF-24b), Wleń 1.5 km N (WIF-31i); XI-III; forest, mixed forest, tree clump; deciduous branch, *Corylus avellana* branch, *Salix* sp. branch.

***Merismodes connivens* (P. Karst.) Knudsen**

Kaczawskie Foothills: Bystrzyca 1.5 km N (WIF-17d), Bystrzyca 2.1 km N (WIF-13g); II-III; riparian forest; *Tilia* sp. wood, *Fraxinus* sp. stump.

***Mucidula mucida* (Schrad.) Pat. (= *Oudemansiella mucida* (Schrad.) Höhn.)**

Kaczawskie Mts.: Góra Milek Res.; X; *Cephalanthero-Fagetum*; *Fagus sylvatica* branch and log.

***Muscinupta laevis* (Fr.) Redhead, Lücking & Lawrey (= *Cyphellostereum laeve* (Fr.) D.A. Reid**

Kaczawskie Mts.: Bystrzyca (WIF-134dx); X-XII; roadside slope, near a stream; mosses.

***Mutinus caninus* (Huds.) Fr.**

Kaczawskie Mts.: Bystrzyca 0,8-0.9 km S (WIF-127a), Bystrzyca 1 km S (WIF-132c); VIII-IX; oak-hornbeam forest, riparian forest, under spruce; soil. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; spruce forest; soil. **Kaczawskie Foothills:** Belczyna 1.1 km E (WIF-122), Belczyna 0.7 km S (WIF-123g), Bystrzyca 1.4 km N (WIF-19b), Bystrzyca 1.2 km SE (WIF-126o), Bystrzyca 2.1 km N (WIF-13g), Górczyca 1.1 km W (WIF-24g), Przeździec 0.8 km SW (WIF-22), Wleń 1.9 km N (WIF-30d), Wleń 2 km N (WIF-29h, 30f); VII-X; mixed forest, pine forest, alder forest, under lime tree, spruce forest; soil.

***Mycena abramsii* (Murrill) Murrill**

Izerskie Foothills: Wleń 2.7 km NW (WIF-44k); VI; riparian forest; deciduous twigs. **Kaczawskie Foothills:** Przeździec 0.9 km S (WIF-22k); VI; spruce forest on the habitat of riparian forest; deciduous branch.

***Mycena acicula* (Schaeff.) P. Kumm.**

Izerskie Foothills: Radomice (limestone quarry), Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VI-VIII; deciduous forest, overgrown quarry; litter. **Kaczawskie Foothills:** Belczyna 0.7 km S (WIF-123g), Bystrzyca 0.5 km N (WIF-20l), Bystrzyca 2.4 km NW (near WIF-15/21); VII-X; young spruce forest, overgrown wastelands, basalt quarry; wood, litter, soil.

***Mycena aetites* (Fr.) Quél.**

Kaczawskie Mts.: Rząśnik 2.3 km W (near WIF-128); X; pasture; soil. **Izerskie Foothills:** Wleń 2.1 km SW (ŁuF – Gniazdo Mt), Wleń (at the Bóbr river near water intake); X-XI; mixed forest, wastelands; soil. **Kaczawskie Foothills:** Bystrzyca 0.8 km S (near WIF-20w), Bystrzyca 2.6 km NW (WIF-21f), Przeździec 1 km SW (near WIF-23), Sobota 1.5 km S; X-XI; meadow, pasture, spruce forest; litter, soil.

***Mycena amicta* (Fr.) Quél.**

Kaczawskie Mts.: Góra Milek Res.; X; deciduous forest; litter. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 2.3 km SW (ŁuF-66c – Gniazdo Mt); V-VI; riparian forest, spruce forest; wood, *Picea abies* log. **Kaczawskie Foothills:** Belczyna 1.8 km S (WIF-126j), Bystrzyca 0.6 km N (WIF-20k), Bystrzyca 1.4 km N (WIF-17h), Wleń 2.3 km NE (WIF-26m); VI-X; oak-hornbeam forest, mixed forest; litter.

***Mycena atropapillata* Kühner & Maire**

Izerskie Foothills: Radomice (limestone quarry); XI; overgrown quarry; soil.

***Mycena aurantiomarginata* (Fr.) Quél.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2.1 km N (WIF-44d); XI; mixed forest; litter.

Kaczawskie Foothills: Bełczyna 2 km SE (WIF-125b), Bystrzyca 1.8 km E (WIF-126m), Bystrzyca 2.1 km N (WIF-14h), Bystrzyca 2.5 km NE (WIF-15h), Górczyca 1.1 km W (WIF-24g); IX-XI; deciduous forest, mixed forest (under spruce), pine forest, spruce forest; litter.

***Mycena capillaris* (Schumach.) P. Kumm.**

Kaczawskie Mts.: Bystrzyca 1.2 km S (WIF-133); XI; mixed forest; decaying *Quercus* sp. leaves, decaying *Fagus sylvatica* leaves. **Kaczawskie Foothills:** Bystrzyca 1.8 km N (WIF-18b), Sobota 1.5 km S; X-XI; mixed sapling stand, pasture; decaying *Quercus* sp. leaves, decaying *Fagus sylvatica* leaves.

***Mycena cinerella* (P. Karst.) P. Karst.**

Izera Mts.: Szklarska Poręba district (0.3 km E from the tourist hostel “Orle”); X; spruce forest; litter.

***Mycena diosma* Krieglst. & Schwöbel**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Tarczyn 1 km NE (WIF-133h); VIII-X; spruce forest, hornbeam-hazel shrubs; litter, soil. **Izerskie Foothills:** Wleń 1.7 km SW (ŁuF – Gniazdo Mt); VIII-XI; spruce forest; litter, soil.

Mycena epipterygia* (Scop.) Gray var. *epipterygia

Kaczawskie Mts.: Tarczyn 0.9 km NE (WIF-134c), Tarczyn 1.1 km E (WIF-134j); XI-XII; mixed forest, spruce forest; wood, litter. **Izerskie Foothills:** Wleń 1.5 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF-59c – Gniazdo Mt); X; spruce forest; litter. **Kaczawskie Foothills:** Bystrzyca 2.5 km NW (WIF-15k); X; deciduous sapling stand; litter.

***Mycena epipterygia* (Scop.) Gray var. *pelliculosa* (Fr.) Maas Geest**

Izerskie Foothills: Zalesie 0.8 km N (LuF-200h); X; mixed sapling stand (mire margin); litter. **Kaczawskie Foothills:** Wleń 1.4 km NW (WIF-31); XI; mixed forest; soil.

***Mycena filopes* (Bull.) P. Kumm.**

Izera Mts.: Szklarska Poręba district (Sępik Mt N slope); X; spruce forest; wood, bark. **Kaczawskie Mts.:** Wleń 2.5 km E (near WIF-134z); XI; overgrown wastelands; soil. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest; soil. **Kaczawskie Foothills:** Bełczyna 1.2 km SE (WIF-126n), Bełczyna 2 km SE (WIF-125b); VII, XI; riparian forest, spruce forest; litter.

***Mycena flavescens* Velen.**

Izera Mts.: Szklarska Poręba district (0,2 km E from the tourist hostel “Orle”); XI; spruce forest margin; soil.

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); XI; riparian forest; deciduous log. **Izerskie Foothills:** Wleń 1.2 km SW (ŁuF-58m); IX; deciduous forest; litter.

***Mycena flavoalba* (Fr.) Quél.**

Kaczawskie Mts.: Tarczyn 0.6 km E (Traczynka Mt); XI; overgrown pasture; soil. **Izerskie Foothills:** Wleń (near Bóbr river by the water intake); XI; wastelands; soil. **Kaczawskie Foothills:** Bełczyna 1 km SE (WIF-122f), Sobota 1.5 km S; X-XI; beech forest, pasture; litter, soil.

***Mycena galericulata* (Scop.) Gray**

Kaczawskie Mts.: Tarczyn 0.9 km NE (WIF-134c), Buki Sudeckie Res.; IX-X; riparian forest (helocrene), spruce forest; deciduous wood, stump. **Izerskie Foothills:** Wleń 1.5 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.7 km SW (ŁuF – Gniazdo Mt), Zalesie 0.8 km N (ŁuF-200h); VIII-X; spruce forest, mixed sapling stand (mire), deciduous forest, mixed forest; wood. **Kaczawskie Foothills:** Bystrzyca 1.2 km SE (WIF-126n), Bystrzyca 2 km NE (WIF-15s), Bystrzyca 2.5 km NW (WIF-26f), Bystrzyca 2.7 km NW (near WIF-26f), Wleń 1.3 km N (WIF-31h), Wleń 1.5 km NW (near WIF-33d), Wleń 2.1 km N (WIF-30c), Wleń 2.6 km N (WIF-25a); IX-I, V; riparian forest, deciduous sapling stand, coniferous forest, deciduous forest, black cherry shrubs (holocene in spruce forest); wood, *Salix* sp. wood, deciduous stump, soil.

***Mycena galopus* (Pers.) P. Kumm. var. *candida* J.E. Lange**

Kaczawskie Mts.: Wleń 1.3 km NE (WIF-136b); X; mixed forest; litter.

Mycena galopus* (Pers.) P. Kumm. var. *galopus

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); X; oak-hornbeam forest; wood. **Izerskie Foothills:** Wleń 1.2 km SW (ŁuF-58m), Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF-59 – Gniazdo Mt), Zalesie 0.8 km NE (ŁuF-199g); V, VIII-XI; deciduous forest, mixed forest (mire), under pine; wood, litter, *Sphagnum* sp. **Kaczawskie Foothills:** Bełczyna 2 km SE (WIF-125b), Bystrzyca 2.5 km NW (WIF-26f), Wleń 2.3 km NE (WIF-26m); VIII-XI; coniferous forest, spruce forest, mixed tree clump near a stream; litter.

***Mycena galopus* (Pers.) P. Kumm. var. *leucogala* (Cooke) J.E. Lange**

Izera Mts.: at the bank of Izera river (0.4 km S from Torfowiska Rzeki Izery Res.); X; spruce forest; litter.

Karkonosze Mts.: Karkonosze NP (Śnieżne Kotły); VIII; spruce forest; litter. **Izerskie Foothills:** Wleń (Południowe Es.); XI; garden; *Astilbe* sp. clump.

***Mycena haematopus* (Pers.) P. Kumm.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a), Góra Milek Res.; VI, VIII; beech forest, riparian forest; log, *Acer pseudoplatanus* wood.

***Mycena hiemalis* (Osbeck) Quél.**

Kaczawskie Foothills: Wleń 2.1 km N (WIF-30c); XII; deciduous sapling stand; *Salix* sp. trunk.

***Mycena inclinata* (Fr.) Quél.**

Kaczawskie Mts.: Bystrzyca 1.2 km S (WIF-133b), Rząśnik 1.9 km W (WIF-290g), Rząśnik 2.7 km W (WIF-128b), Wleń 1.3 km NE (WIF-136b); VIII-X; deciduous forest, mixed forest; wood, ?*Quercus* sp. stump, stump.

Izerskie Foothills: Wleń 1.2 km SW (ŁuF-58m), Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2.8 km N (WIF-44j); VIII-X; oak-hornbeam forest, deciduous forest; deciduous log, ?*Quercus* sp. stump, stump. **Kaczawskie Foothills:** Bystrzyca 1.2 km SE (WIF-126n), Bystrzyca 2.1 km N (WIF-13g); VIII-IX; oak-hornbeam forest, riparian forest; stump.

***Mycena latifolia* (Peck) A.H. Sm.**

Kaczawskie Foothills: Bystrzyca 1.7 km N (WIF-18h); XI; young spruce forest; litter.

***Mycena leptcephala* (Pers.) Gillet**

Kaczawskie Mts.: Bystrzyca 1.1 km S (WIF-133c), Tarczyn 1.2 km NE (WIF-133f), Wleń 1.9 km E (WIF-136a); IV, X-XI; beech forest, mixed forest; litter, soil. **Izerskie Foothills:** Łupki (near tourist hostel); X-XI; lawn; soil. **Kaczawskie Foothills:** Bełczyna 1.4 km SE (near WIF-123s), Wleń 2.6 km N (WIF-25a); X-XI; deciduous forest, meadow margin; soil. **Sudetes Foreland:** Szklary 0.5 km SW; IX; overgrown outcrop; soil.

***Mycena maculata* P. Karst.**

Karkonosze Mts.: Szklarska Poręba district (Mumławski Wierch Mt); X; spruce forest; *Picea abies* log. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF-59b – Gniazdo Mt); X; riparian forest; *Acer pseudoplatanus* stump.

***Mycena metata* (Fr.) P. Kumm.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a), Rząśnik 1.5 km W; X; riparian forest; log, litter. **Izerskie Foothills:** Łupki (near tourist hostel), Wleń 1.5 km SW (ŁuF – Gniazdo Mt), Wleń 1.7 km SW (ŁuF – Gniazdo Mt); XI; lawn, spruce forest; litter, soil. **Kaczawskie Foothills:** Wleń 1.8 km NE (WIF-27c); XI; spruce forest; litter.

***Mycena mirata* (Peck) Sacc.**

Izerskie Foothills: between Barcinek and Wrzeszczyn (near estuary of Kamienna river to Bóbr river); XII; spruce forest; stump.

***Mycena mitis* Maas Geest.**

Kaczawskie Foothills: Bystrzyca 2.5 km NW (WIF-15k); X; deciduous sapling stand; litter.

***Mycena niveipes* (Murrill) Murrill**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-64a – Gniazdo Mt); X; riparian forest; litter.

***Mycena olida* Bres. (= *M. minutula* (Peck) Sacc.)**

Kaczawskie Foothills: Rząśnik 1.3 km NW (WIF-125g); X; mixed forest; deciduous wood.

***Mycena olivaceomarginata* (Massee) Massee**

Kaczawskie Mts.: Tarczyn 1.8 km E (near WIF-133); X; overgrown meadow; soil. **Izerskie Foothills:** Łupki (near tourist hostel); XI; lawn; soil.

***Mycena pearsoniana* Singer s. auct. Eur.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); XI; riparian forest; litter.

***Mycena plumipes* (Kalchbr.) P.A. Moreau (= *M. strobilicola* J. Favre & Kühner)**

Kaczawskie Foothills: Bystrzyca 1.7 km N (WIF-18h); III; spruce forest; *Picea abies* cone, litter.

***Mycena polyadelpha* (Lasch) Kühner**

Izerskie Foothills: Wleń 2.1 km SW (ŁuF – Gniazdo Mt); XI; mixed forest; decaying *Quercus* sp. leaves, decaying *Populus tremula* leaves. **Kaczawskie Foothills:** Górczyca 1.1 km W (WIF-24d), Przeździec 0.5 km NE (WIF-24z); XI-XII; oak-hornbeam forest, mixed forest; decaying *Quercus* sp. leaves.

***Mycena polygramma* (Bull.) Gray**

Kaczawskie Mts.: Tarczyn 0.8 km N (WIF-134ax), Wleń 1.4 km SE (WIF-137i), Góra Milek Res.; X-XI; oak forest, deciduous forest, mixed forest; *Carpinus betulus* log, stump, litter. **Izerskie Foothills:** Wleń 1.6 km NW (WIF-46f); X; spruce forest; litter. **Kaczawskie Foothills:** Belczyna 0.6 km S (near WIF-123f), Belczyna 0.8 km S (WIF-123j), Bystrzyca 2.5 km NE (WIF-15g), Bystrzyca 2.5 km NW (WIF-15k), Radomiłowice 1.1 km SW (WIF-11g), Radomiłowice 1.4 km SW (WIF-11l), Rząśnik 1.3 km NW (WIF-125g), Rząśnik 1.8 km NW (WIF-125), Nad Groblą Res., Wąwóz Siedmicki Res.; IX-XII; deciduous forest, mixed forest, deciduous sapling stand, spruce forest; wood, stump, litter, soil.

***Mycena pseudocorticola* Kühner**

Izerskie Foothills: Wleń 1.4 km NW (WIF-47p); XI; mixed sapling stand; *Salix* sp. branch.

***Mycena pseudopicta* (J.E. Lange) Kühner**

Kaczawskie Mts.: Tarczyn 0.6 km E (Tarczynka Mt); XI; overgrown slate excavation; soil.

***Mycena pura* (Pers.) P. Kumm. f. *alba* (Gillet) Arnolds**

Kaczawskie Foothills: Przeździec 1.7 km E (WIF-12s); X; mixed forest; soil.

Mycena pura* (Pers.) P. Kumm. f. *pura

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1 km S (WIF-133a), Bystrzyca 1.5 km S (WIF-133d); VII-XI; oak-hornbeam forest, riparian forest, spruce forest; litter, soil. **Izerskie Foothills:** Wleń 1.2 km SW (ŁuF-58m), Wleń 1.4 km NW (WIF-46g), Wleń 1.6 km SW (ŁuF-64g – Gniazdo Mt); VII-X; deciduous forest, mixed forest; litter, soil. **Kaczawskie Foothills:** Przeździec 1.3 km W (WIF-24k); X; riparian forest; litter.

***Mycena purpureofusca* (Peck) Sacc.**

Kaczawskie Mts.: Bystrzyca 1.1 km S (WIF-134a); VII; riparian forest; *Acer pseudoplatanus* wood.

***Mycena rosea* (Bull.) Gramberg**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2.8 km N (WIF-44j), Wleń 1.6 km NW (WIF-46f); X-XI; oak-hornbeam forest, mixed forest, spruce forest; litter, soil. **Kaczawskie Foothills:** Wleń 1.3 km N (WIF-31h); VIII; mixed forest (holocene); soil. **Sudetes Foreland:** Szklary 0.5 km SW; IX; overgrown spoil tip; litter.

***Mycena rosella* (Fr.) P. Kumm.**

Izerskie Foothills: Wleń 1.7 km SW (ŁuF – Gniazdo Mt); XI; spruce forest; litter.

***Mycena rubromarginata* (Fr.) P. Kumm.**

Kaczawskie Foothills: Wleń 2.3 km NE (WIF-25c); X; spruce forest; *Picea abies* wood.

***Mycena sanguinolenta* (Alb. & Schwein.) P. Kumm.**

Kaczawskie Mts.: Rząśnik 1.1 km SW (WIF-290l); VII; oak-hornbeam forest; litter. **Izerskie Foothills:** Wleń 1.2 km SW (ŁuF-58m), Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Zalesie 0.8 km N (ŁuF-200h), Zalesie 0.8 km NE (ŁuF-199g); VIII; deciduous forest, mixed forest, mixed forest (mire); ?*Quercus* sp. bark, stump, *Sphagnum* sp. **Kaczawskie Foothills:** Bystrzyca 0.6 km N (WIF-20k), Bystrzyca 1.8 km E (WIF-126n), Bystrzyca 2.1 km N (WIF-14h), Bystrzyca 2.8 km NW (WIF-22o), Wleń 1.6 km NE (WIF-27j), Wleń 2 km N (WIF-30f); VI-VIII; coniferous forest, deciduous forest, mixed forest, under lime tree, under spruce; litter, *Picea abies* twigs.

***Mycena silvae-nigrae* Maas Geest. & Schwöbel**

Kaczawskie Foothills: Wleń 1.5 km NE (near WIF-33d), Wleń 2.7 km NE (WIF-26d); XII, V; deciduous forest, riparian forest; wood.

Mycena speirea* (Fr.) Gillet f. *speirea

Izerskie Foothills: Wleń 1.4 km NW (WIF-47p), Wleń 2.1 km SW (ŁuF-65h – Gniazdo Mt); VI, XI; mixed forest, spruce forest; ?*Alnus incana* branch, ?*Salix* sp. bark. **Kaczawskie Foothills:** Bystrzyca 0.6 km N (WIF-20l), Bystrzyca 1.4 km N (WIF-17f), Bystrzyca 1.6 km N (WIF-17b), Przeździec 1.3 km S (near WIF-23.25 – quarry); VI, VIII, XII; deciduous forest, mixed forest, basalt quarry, melaphyre quarry; ?*Fraxinus* sp. log, litter, bark, wood, deciduous twigs.

***Mycena speirea* (Fr.) Gillet f. *camptophylla* (Berk.) Kühner**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF-64 – Gniazdo Mt); VI; deciduous forest; litter.

***Mycena stipata* Maas Geest. & Schwöbel**

Izerskie Mts.: Sepik Mt (N slope); X-XI; spruce forest; *Picea abies* stump.

***Mycena stylobates* (Pers.) P. Kumm.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; mixed forest; litter. **Kaczawskie Foothills:** Bystrzyca 1.7 km N (WIF-16g), Bystrzyca 1.8 km SE (WIF-124a), Radomiłowice 0.7 km SW (near WIF-11c); VII-VIII; mixed forest, log landing, spruce forest; litter.

***Mycena tintinabulum* (Fr.) Quél.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Rząśnik 1.1 km SW (WIF-290l); I-II; oak-hornbeam forest; stump. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); II-III; deciduous forest, riparian forest; ?*Alnus incana* log, stump. **Kaczawskie Foothills:** Bystrzyca 1.2 km SE (WIF-126n), Bystrzyca 1.3 km SE (WIF-126k), Przeździec 0.4 km NE (near WIF-24z), Przeździec 1.7 km E (near WIF-12n), Wleń 1.5 km NE (near WIF-28); XII-III, X; deciduous forest, riparian forest, tree clump, shrubs near a stream; *Alnus incana* wood, deciduous stump.

***Mycena viridimarginata* P. Karst.**

Kaczawskie Foothills: Bystrzyca 2.8 km NW (WIF-22o, near WIF-22); VI-VII; deciduous forest; deciduous wood, log.

***Mycena vitilis* (Fr.) Quél.**

Kaczawskie Mts.: Bystrzyca, Tarczyn 1.2 km E (WIF-133g), Wleń 1.3 km NE (WIF-136b); X-XII; coniferous sapling stand, deciduous forest, mixed forest; *Quercus* sp. branch, litter. **Izerskie Foothills:** Wleń (Południowe Es.); X; lawn; soil. **Kaczawskie Foothills:** Belczyna 0.8 km S (WIF-123i), Belczyna 1 km W (WIF-13l), Bystrzyca 2.5 km NE (WIF-15h), Przeździec 1.1 km W (WIF-24n), Radomiłowice 0.9 km W (WIF-12Ac), near Wąwóz Lipy Res.; VII, X-XII; oak forest, deciduous forest, mixed forest, deciduous sapling stand; litter, soil.

***Mycena vulgaris* (Pers.) P. Kumm.**

Izerskie Foothills: Łupki (near tourist hostel), Wleń 1.4 km SW (ŁuF – Gniazdo Mt); X-XI; mixed forest, lawn; litter, soil. **Kaczawskie Foothills:** Belczyna 0.7 km S (WIF-123g), Bystrzyca 1.7 km W (near WIF-26); X-XI; meadow, spruce sapling stand; litter, soil.

***Mycena zephrus* (Fr.) P. Kumm.**

Kaczawskie Foothills: Bystrzyca 2.5 km NE (WIF-15i); X; spruce forest; litter.

***Mycenella salicina* (Velen.) Singer**

Izerskie Foothills: Radomice (limestone quarry); XI; overgrown quarry; soil.

***Mycetinis querceus* (Britzelm.) Antonín & Noordel. (= *Marasmius quercinus* Britzelm.)**

Kaczawskie Foothills: Bełczyna 0.7 km S (WIF-123h); XI; mixed sapling stand; litter.

***Mycetinis scorodonius* (Fr.) A. Wilson & Desjardin (= *Marasmius scorodonius* (Fr.) Fr.)**

Kaczawskie Mts.: Tarczyn, 1.2 km NE (WIF-133f); VII; mixed forest; litter.

***Myxarium hialinum* (Pers.) Donk (= *Exidia nucleata* (Schwein.) Burt)**

Kaczawskie Mts.: Wleń 1.7 km SE (WIF-137j); I; deciduous forest; *Tilia* sp. branch. **Kaczawskie Foothills:** Górczyca 1.4 km W (WIF-24h); II; mixed forest; *Tilia* sp. branch.

***Naematelia encephala* (Pers.) Fr. (= *Tremella encephala* Pers.)**

Kaczawskie Foothills: Bełczyna 1.8 km SE (WIF-124d); II; mixed forest; *Pinus sylvestris* branch with *Stereum sanguinolentum* basidiocarps.

***Naucoria escharioides* (Fr.) P. Kumm.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-59b, 64c – Gniazdo Mt); IX-X; deciduous forest, riparian forest; soil.

Kaczawskie Foothills: Bełczyna 1.2 km E (WIF-122a), Przeździec 0.8 km SW (WIF-22m); IX-X; alder forest; soil. **Kaczawskie Mts.:** Bystrzyca 1 km S (WIF-133a), Bystrzyca 1.2 km S (WIF-132c), Buki Sudeckie Res.; X; mixed forest, *Carici remotae-Fraxinetum*; soil.

***Naucoria luteolofibrillosa* (Kühner) Pilát**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a), Rzęśnik 1.6 km SW (WIF-290l); IX-X; riparian forest, alder forest; soil. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); X; alder forest; soil. **Kaczawskie Foothills:** near Wąwóz Lipy Res.; XI; riparian forest; soil.

***Naucoria scolecina* (Fr.) Quél.**

Kaczawskie Mts.: Rzęśnik 2.1 km W (WIF-128d), Rzęśnik 2.3 km W (WIF-128g), Buki Sudeckie Res.; X; deciduous forest, mixed forest, *Carici remotae-Fraxinetum*; soil. **Izerskie Foothills:** Zalesie 0.8 km N (ŁuF-200h); X; mixed sapling stand; soil. **Kaczawskie Foothills:** Bełczyna 1.1 km E (near WIF-122), Sobota 1.6 km SW (WIF-24Aa); VIII-XI; forest near a stream, alder forest; soil.

***Naucoria striatula* P.D. Orton**

Kaczawskie Foothills: Sobota 1.6 km SW (WIF-24Aa); XI; deciduous sapling stand near a stream; soil.

***Naucoria subconsersa* P.D. Orton**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); X; riparian forest; soil. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF-59b – Gniazdo Mt); X; riparian forest; soil. **Kaczawskie Foothills:** Przeździec 0.8 km SW (WIF-22m), Sobota 1.6 km SW (WIF-24Aa); VIII-IX; alder forest; soil.

***Neolentinus lepideus* (Fr.) Redhead & Ginns (= *Lentinus lepideus* (Fr.) Fr.)**

Izerskie Foothills: Zalesie 0.8 km N (LuF); VII-IX; spruce-pine forest, mire (under spruce and pine); coniferous wood, *Pinus sylvestris* stump.

***Omphalina pyxidata* (Bull.) Quél.**

Izera Mts.: Sępik Mt (by the tourist trail between Szklarska Poręba and Hala Izerska); X; forest roadside; soil. **Kaczawskie Mts.:** Mysłów (Mysłów-Sobocin quarry); IX; overgrown limestone quarry; soil. **Karkonosze Mts.:** Jakuszyce 0.9 km SE; X; roadside; soil.

***Oxyporus populinus* (Schumacher) Donk**

Kaczawskie Foothills: Wąwóz Siedmicki Res.; X; deciduous forest; deciduous trunk.

***Oxyporus ravidus* (Fr.) Bondartsev & Singer**

Kaczawskie Foothills: Bystrzyca 2.1 km N (WIF-14h); VIII; between spruce forest and deciduous forest; ?hornbeam stump.

***Panaeolina foenisecii* (Pers.) Maire var. *foenisecii* (= *Panaeolus foenisecii* (Pers.) Kühner)**

Izerskie Foothills: Wleń (Południowe Es.); VI; lawn; soil.

Panaeolus acuminatus* (Schaeff.) Quél. var. *acuminatus

Karkonosze Mts.: Szklarska Poręba; VIII; mixed forest; *Equus caballus* dung. **Kaczawskie Foothills:** Belczyna 1.4 km SE (near WIF-123s), Bystrzyca 2.4 km NW (near WIF-22), E part of Bystrzyca village (WIF-20o, 20y), Sobota 1.5 km S; V-XI; meadow, pasture, field, overgrown wastelands; soil, *Equus caballus* dung.

***Panaeolus fimicola* (Pers.) Gillet**

Kaczawskie Foothills: Sobota 1.5 km S; IV; pasture; soil.

***Panaeolus foenisecii* (Pers.) J. Schröt. var. *intermedius* E. Ludw.**

Izerskie Foothills: Wleń (Południowe Es.); VI; lawn; soil.

Panaeolus papilionaceus* (Bull.) Quél. var. *papilionaceus

Izera Mts.: Sępik Mt (by the tourist trail between Szklarska Poręba and Hala Izerska); X; forest roadside; dung. **Kaczawskie Mts.:** Rząśnik 1.5 km W (near WIF-290); VII-IX; pasture; soil. **Kaczawskie Foothills:** Wleń 2.6 km N (WIF-25a); log landing; soil, woodchips.

***Panaeolus papilionaceus* (Bull.) Quél. var. *parvisporus* Ew. Gerhardt**

Kaczawskie Foothills: Bystrzyca 1.5 km NE (WIF-31c); X; meadow; soil.

***Panellus mitis* (Pers.) Singer**

Kaczawskie Mts.: Bystrzyca 1.3 km SE (WIF-130h), Rząśnik 2.5 km W (WIF-128m); XI; coniferous forest, mixed forest; coniferous wood. **Kaczawskie Foothills:** Bystrzyca 2.6 km NW (WIF-26f); I; spruce forest; coniferous wood.

***Panellus stipticus* (Bull.) P. Karst.**

Kaczawskie Foothills: Bełczyna 0.7 km S (WIF-123g), Wleń 1.4 km NW (WIF-31b); X; mixed forest, spruce forest; wood.

***Panus conchatus* (Bull.) Fr. (= *Lentinus torulosus* (Pers.) Lloyd)**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VII; deciduous forest; deciduous log. **Kaczawskie Foothills:** Bystrzyca 2.5 km NW (WIF-26f), Bystrzyca 2.8 km NW (WIF-22o, 22r), Górczyca 1 km E (WIF-12Ao), Wleń 1.8 km NE (WIF-33d); VIII-X; birch forest, coniferous forest, deciduous forest, mixed forest; wood, *Populus* sp./*Quercus* sp. log, *Betula* sp. stump, *Populus tremula* stump.

***Parasola auricoma* (Pat.) Redhead, Vilgalys & Hopple (= *Coprinus auricomus* Pat.)**

Izerskie Foothills: Wleń, Wleń 1.5 km N; VII-VIII; deciduous forest, roadside; soil.

***Parasola conopilea* (Fr.) Örstadius & E. Larss. (= *Psathyrella conopilus* (Fr.) A. Pearson & Dennis)**

Izerskie Foothills: Wleń 1.3 km NW (WIF-47o), Wleń 1.4 km SW (at the Bóbr river); IX-XI; mixed forest, riverside tree clump; litter. **Kaczawskie Foothills:** Bystrzyca 1.2 km N (WIF-16m); XI; ash forest; litter.

***Parasola galericuliformis* (Watling) Redhead, Vilgalys & Hopple**

Kaczawskie Mts.: Grudno; IX; limestone quarry; soil.

***Parasola kuehnerii* (Uljé & Bas) Redhead, Vilgalys & Hopple (= *Coprinus kuehnerii* Uljé & Bas)**

Kaczawskie Mts.: Wleń 2 km E (WIF-134gx); VII; lawn, shrubs; soil. **Izerskie Foothills:** Wleń; IX; lawn; soil.

***Parasola leioccephala* (P.D. Orton) Redhead, Vilgalys & Hopple (= *Coprinus leioccephalus* P.D. Orton)**

Kaczawskie Mts.: Wleń 1.1 km E (WIF-136g, 136h); IX; forest roadside; soil. **Izerskie Foothills:** Wleń, Wleń 1.6 km SW (ŁuF-64 – Gniazdo Mt); VI-VIII; lawn, deciduous forest; soil. **Kaczawskie Foothills:** Bełczyna 1.8 km SE (WIF-124a), Bystrzyca 0.5 km N (WIF-20l), Leszczyna 1 km N (near Ciche Szczyście mine), Radomiłowice 0.9 km S (WIF-11c), Wleń 2.3 km N (WIF-29a); IV-IX; roadside, log landing, between spruce forest and riparian forest, oak-hornbeam forest margin, oak-hornbeam forest; soil.

***Parasola lilatincta* (Bender & Uljé) Redhead, Vilgalys & Hopple**

Kaczawskie Foothills: Bystrzyca 1.7 km N (between WIF-14m & WIF-18h); VIII; forest road; soil.

***Parasola megasperma* (P.D. Orton) Redhead, Vilgalys & Hopple**

Kaczawskie Mts.: Rząśnik 2.5 km W (WIF-129c); VIII; spruce forest; soil.

***Parasola plicatilis* (Curtis) Redhead, Vilgalys & Hopple (= *Coprinus plicatilis* (M.A. Curtis) Fr.)**

Izerskie Foothills: Wleń, Wleń 1.4 km SW (ŁuF-64c – Gniazdo Mt); VIII-IX; lawn, oak-hornbeam forest; soil.

Kaczawskie Foothills: Bełczyna 1.8 km SE (WIF-124a), Sobota 1.5 km SW, Wleń 2.4 km N (WIF-29b); IX-XI; meadow, log landing, mixed forest; soil.

***Parasola schroeteri* (P. Karst.) Redhead, Vilgalys & Hopple (= *Coprinus schroeteri* P. Karst.)**

Kaczawskie Foothills: Sobota 1.5 km S, Wleń 2.1 km N (WIF-30c); IV-IX; deciduous sapling stand, pasture; soil.

***Paxillus rubicundulus* P.D. Orton s.l.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a), VIII; riparian forest (under black alder); soil. **Kaczawskie Foothills:** Wleń 2.7 km NW (near WIF-26); IX; poplar-birch shrubs; soil.

***Peniophora cinerea* (Pers.) Cooke**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt, Białe Skały); XI; mixed forest; *Tilia* sp. branch.

***Peniophora incarnata* (Pers.) P. Karst.**

Kaczawskie Foothills: Bystrzyca 1.8 km N (WIF-16a); II; oak-hornbeam forest; *Tilia* sp. branch.

***Peniophora limitata* (Chaillet ex Fr.) Cooke**

Kaczawskie Foothills: Bystrzyca 1.4 km N (WIF-17c); riparian forest; *Fraxinus* sp. branch.

***Peniophora polygonia* (Pers.) Bourdot & Galzin**

Kaczawskie Mts.: Wleń 1.4 km SE (WIF-137h); I; margin of pine sapling stand; *Populus tremula* branch.

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); XI; riparian forest; *Populus tremula* branch.

***Peniophora quercina* (Pers.) Cooke**

Kaczawskie Foothills: Bystrzyca 2.6 km SW (WIF-26j), Górczyca 1.1 km W (WIF-24d); IV, X; mixed forest; *Quercus* sp. branch.

***Peniophora rufomarginata* (Pers.) Bourdot & Galzin**

Kaczawskie Mts.: Tarczyn 0.6 km E (near WIF-134 – Tarczynka Mt); VII; overgrown pasture; ?*Populus tremula* branch.

***Perenniporia medulla-panis* (Jacq.) Donk**

Izerskie Foothills: Pokrzywnik 1.2 km S (WIF-176d – near Pilchowice Lake), between Łupki and Klecza (near Piaskowcowe Porwaki); I, III; mixed forest, spruce forest; *Quercus* sp. log, *Acer pseudoplatanus* stump.

Kaczawskie Foothills: Przeździec 0.7 km NE (near WIF-24bx); XII; roadside tree clump; stump.

***Phaeolus schweinitzii* (Fr.) Pat.**

Kaczawskie Mts.: Bystrzyca 1.1 km S (WIF-132b), Bystrzyca 1.2 km S (WIF-133b), Bystrzyca 1.3 km SE (WIF-130h), Bystrzyca 1.6 km SE (WIF-130i, 131i); VII-IX; mixed forest, spruce forest; wood, *Picea abies* wood, *Larix* sp. wood, *Pinus sylvestris* trunk, roots. **Kaczawskie Foothills:** Bystrzyca 2.2 km N (WIF-15d), Bystrzyca 2.3 km NW (WIF-15p); VII, IX; mixed sapling stand, spruce forest; wood, *Larix* sp. trunk base.

***Phaeonematoloma myosotis* (Fr.) Bon (= *Pholiota myosotis* (Fr.) Singer)**

Izerskie Foothills: Zalesie 0.8 km NE (LuF-199g); X; mire; *Sphagnum* sp.

***Phaeotremella pseudofoliacea* Rea (= *Tremella foliacea* Pers.)**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1 km S (WIF-133a), Rząśnik 2.7 km W (WIF-135c), Rząśnik 2.8 km W (WIF-135h), Tarczyn 1.2 km NE (WIF-133f); IX-II; riparian forest, oak-hornbeam forest, mixed forest, mixed forest roadside, spruce forest; *Betula* sp. wood, *Quercus* sp. wood, *Corylus avellana* wood, deciduous wood, *Quercus* sp. roots. **Izerskie Foothills:** Wleń 1.4 km NW (WIF-47k), Wleń 1.6 km SW (LuF – Gniazdo Mt); XII-IV; deciduous forest; *Corylus avellana* wood, *Tilia* sp. wood. **Kaczawskie Foothills:** Bystrzyca 2.5 km NW (WIF-22p), Górczyca 1.4 km W (WIF-24h), Przeździec 1.7 km E (WIF-12m), Radomilowice 1.1 km SW (WIF-11g), Wleń 0.9 km NW (WIF-32b); VIII-II; oak forest, mixed forest, deciduous forest; *Quercus* sp. wood, *Corylus avellana* wood, ?*Alnus* sp. wood, deciduous wood.

Phallus impudicus* L. var. *impudicus

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 0.9 km S (WIF-127a); VIII; oak-hornbeam forest, riparian forest; soil. **Izerskie Foothills:** Wleń 1.4 km NW (WIF-46a); VIII; deciduous sapling stand; soil.

Kaczawskie Foothills: Bystrzyca 1.4 km N (WIF-19b), Bystrzyca 1.8 km N (WIF-16a), Bystrzyca 2.1 km N (WIF-14h), Wleń 1.8 km NE (WIF-33d), Wleń 2.6 km N (WIF-25a); VII-X; oak-hornbeam forest, deciduous forest, mixed forest; soil.

***Phellinus alni* (Bondartsev) Parmasto**

Kaczawskie Mts.: Bystrzyca (WIF-134x); X; mixed forest; deciduous stump (?*Malus* sp., ?*Alnus* sp.).

***Phellinus contiguus* (Pers.) Pat.**

Kaczawskie Mts.: Wleń 1.1 km SE (near WIF-137i); I; deciduous forest; *Corylus avellana* branch. **Izerskie Foothills:** Wleń 1.6 km NW (WIF-46l); III; deciduous forest; *Corylus avellana* branch. **Kaczawskie Foothills:** Bystrzyca 1.4 km N (WIF-19c); IV; riparian forest; *Fraxinus* sp. branch.

***Phellinus ferruginosus* (Schrad.) Pat.**

Kaczawskie Foothills: Wleń 1.5 km N (WIF-31i); X; mixed forest; *Prunus avium* log.

***Phellinus igniarius* (L.) Quél. s.str.**

Kaczawskie Foothills: near Wąwóz Lipy Res.; XI; riparian forest; *Quercus* sp. log.

***Phellinus lundellii* Niemelä**

Kaczawskie Foothills: Bystrzyca 2.4 km NW (WIF-22/26), Bystrzyca 2.5 km NW (WIF-22p); III, XII; deciduous forest, mixed forest; ?*Fraxinus* sp. log, *Corylus avellana* branch.

***Phellinus nigricans* (Fr.) P. Karst.**

Kaczawskie Foothills: Wilcza Góra Res. (top of the Wilcza Mt); IV; deciduous forest; *Prunus avium* trunk.

***Phellinus punctatus* (P. Karst.) Pilát**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); II; riparian forest; *Salix caprea* trunk. **Kaczawskie Foothills:** Belczyna 1.8 km SE (WIF-124d), Bystrzyca 2.1 km NW (near WIF-26f), Wleń 2.7 km NE (WIF-26d); I-II, XI; deciduous forest, mixed forest; *Corylus avellana* branch, *Alnus* sp. trunk, *Salix caprea* trunk.

***Phellinus ribis* (Schumach.) Quél. (= *Phylloporia ribis* (Schumach.) Ryvarden)**

Kaczawskie Foothills: Górczyca 1.6 km W, Wleń 1.5 km NW (near WIF-31), Wleń 2.6 km N (WIF-25a); I-II, X; riverbank tree clump, deciduous forest; spindle tree trunk.

***Phellinus tuberosus* (Baumg.) Niemelä (= *P. pomaceus* (Fr.) Pilát)**

Kaczawskie Mts.: Bystrzyca 0.6 km SE; I; roadside tree clump; *Prunus cerasifera* stump.

***Phellinus viticola* (Schwein.) Donk**

Karkonosze Mts.: Mumławski Wierch Mt (ca. 1200 m a.s.l.); X; spruce forest; *Picea abies* wood.

***Phellodon niger* (Fr.) P. Karst.**

Kaczawskie Mts.: Góra Milek Res.; IX; *Dentario enneaphylli*-*Fagetum*; soil.

***Phlebia centrifuga* P. Karst.**

Kaczawskie Foothills: Bystrzyca 1.8 km N (WIF-16a); II; oak-hornbeam forest; ?*Quercus* sp. log.

***Phlebia radiata* Fr.**

Kaczawskie Mts.: Wleń 1.8 km SE (near WIF-137k); I; deciduous forest; *Quercus* sp. branch. **Kaczawskie Foothills:** Wleń 1.9 km NE (WIF-27f); XI; mixed forest; *Betula* sp. log.

***Phlebia tremellosa* (Schr.) Nakasone & Burds.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); XI; spruce forest; deciduous wood. **Kaczawskie Foothills:** Bystrzyca 1.5 km N (WIF-17d), Bystrzyca 2.5 km NE (WIF-15h), Wleń 1.5 km NW (near WIF-31 at the Bóbr river), Wleń 2.1 km NE (WIF-27a), Wleń 2.6 km N (WIF-25a); IX-I; oak-hornbeam forest, spruce forest near a stream, riverside tree clump, deciduous forest; deciduous wood, *Betula* sp. wood, ?*Tilia* sp. wood.

***Phlebiopsis ravenelii* (Cooke) Hjortstam (= *Phanerochaete ravenelii* (Cooke) Burds.)**

Kaczawskie Foothills: Przeździec 0.8 km SW (WIF-22l); IX; mixed forest; *Betula* sp. stump.

***Pholiota adiposa* (Batsch) P. Kumm. s.str.**

Izerskie Foothills: Wleń 2.1 km SW (ŁuF – Gniazdo Mt); X; beech forest; ?*Fagus sylvatica* trunk.

***Pholiota flammans* (Batsch) P. Kumm.**

Kaczawskie Foothills: Bystrzyca 1.2 km SE (WIF-126n), Bystrzyca 1.8 km S (WIF-126m), Wleń 2 km N (WIF-29d); VIII-X; coniferous forest, mixed forest, riparian forest; wood, *Alnus* sp. wood, stump.

Pholiota gummosa* (Lasch) Singer var. *gummosa

Kaczawskie Mts.: Bystrzyca 1.1 km SE (WIF-131g); IX; log landing; wood.

***Pholiota highlandensis* (Peck) Quadr.**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt); VIII; pine forest (old campfire); charcoal. **Kaczawskie Foothills:** Radomilowice 1.1 km SW (WIF-11g); VII; mixed forest (old campfire); charcoal.

***Pholiota lenta* (Pers.) Singer**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); IX; oak-hornbeam forest; litter. **Izerskie Foothills:** Wleń 1.5 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); X-XI; between spruce forest and riparian forest, deciduous forest; wood, litter. **Kaczawskie Foothills:** Belczyna 1.8 km SE (WIF-124a), Bystrzyca 2.5 km NW (WIF-15h), Górczyca 1.1 km SE (WIF-24), Belczyna 0.7 km SE (WIF-123); X-XII; deciduous forest, mixed forest, spruce forest; litter, wood.

***Pholiota limonella* (Peck) Sacc.**

Kaczawskie Mts.: Tarczyn 0.6 km E; X; roadside; *Malus* sp. trunk. **Izerskie Foothills:** Wleń 1.8 km SW (ŁuF – Gniazdo Mt); IX; spruce forest; stump. **Kaczawskie Foothills:** Bystrzyca 2.2 km NW (WIF-26i); IX; deciduous forest; *Alnus* sp. log.

***Pholiota squarrosa* (Weigel) P. Kumm.**

Kaczawskie Mts.: Bystrzyca 0.9 km SE (near WIF-131g); X; tree clump; *Salix* sp. trunk.

***Pholiotina aporos* (Kits van Wav.) Clémenton var. *aporos* (= *Conocybe aporus* Kits van Wav.)**

Izera Mts.: Świeradów-Zdrój (ŚF-268d); V; grassy forest margin; soil. **Kaczawskie Foothills:** Leszczyna 1 km N (near Ciche Szczęście mine); IV; oak-hornbeam forest; soil.

***Pholiotina arrhenii* (Fr.) Singer (= *Conocybe arrhenii* (Fr.) Kits van Wav.)**

Kaczawskie Mts.: Bystrzyca 1.1 km SE (WIF-131g); X; log landing; soil, woodchips. **Kaczawskie Foothills:** Bystrzyca 0.5 km N (WIF-20l); IX; overgrown basalt quarry; soil.

***Pholiotina brunnea* (Watling) Singer (= *Conocybe brunnea* (J.E. Lange & Kühner) ex Watling)**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); IX; riparian forest; soil. **Izerskie Foothills:** Wleń 1.4 km NW (WIF-46a), Wleń 1.6 km NW (WIF-46f), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VII-IX; deciduous forest, spruce forest; litter, soil. **Kaczawskie Foothills:** Bystrzyca 2.2 km N (WIF-15c); X; oak-hornbeam forest; soil.

***Pholiotina dasypus* (Romagn.) P.-A. Moreau**

Izerskie Foothills: Wleń 2.7 km NW (WIF-44k); VI; riparian forest; soil.

***Pholiotina filipes* (G.F. Atk.) Singer (= *Conocybe sulcatipes* (Peck) Kühner)**

Kaczawskie Foothills: Bełczyna 1.2 km SE (WIF-122k); VII; spruce forest; litter.

***Pholiotina vexans* (P.D. Orton) Bon (= *Conocybe blattaria* (Fr.) Kühner ss. Watling)**

Kaczawskie Foothills: Wleń 2.2 km N (WIF-25c); X; oak-hornbeam forest; rooten wood.

***Physisporinus vitreus* (Pers.) Donk**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); X; riparian forest; *Fraxinus* sp. stump. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); X; riparian forest; deciduous wood. **Kaczawskie Foothills:** Przeździec 0.9 km S (WIF-22b), Bystrzyca 2.8 km NW (WIF-22o); VII, IX; beech sapling stand, mixed forest; *Picea abies* stump, litter.

***Piptoporus betulinus* (Bull.) P. Karst.**

Kaczawskie Foothills: Bystrzyca 2.5 km NW (WIF-26f); IX; coniferous forest; *Betula* sp. wood.

***Pleurocybella porrigens* (Pers.) Singer**

Izera Mts.: 0.5 km W from tourist hostel “Chatka Górzystów” (near yellow tourist trail), Kobyla Mt (N slope); X-XI; coniferous forest; wood.

***Pleurotus dryinus* (Pers.) P. Kumm.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133), Tarczyn 0.8 km NE (WIF-134d); IX-X; riparian forest, spruce forest; *Fraxinus* sp. trunk, *Picea abies* trunk. **Kaczawskie Foothills:** Wleń 1.4 km NW (WIF-31b); X; mixed forest; *Quercus* sp. trunk base.

***Pleurotus ostreatus* (Jacq.) P. Kumm.**

Kaczawskie Mts.: Wleń 1.1 km SE (WIF-137i); I; deciduous forest; deciduous wood. **Izerskie Foothills:** Wleń (Południowe Es.), Wleń 1.4 km SW (ŁuF-59b); X-XI; deciduous forest, riparian forest, lawn; wood, *Fraxinus* sp. wood, *Picea abies* wood. **Kaczawskie Foothills:** Przeździec 0.9 km S (WIF-22k), Wleń 1 km NE (near WIF-27, 28); IX, XI; field road, riparian forest; deciduous wood.

***Pleurotus pulmonarius* (Fr.) Quél.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; *Quercus* sp. log. **Izerskie Foothills:** Wleń 2 km SW (ŁuF – Gniazdo Mt); VIII; mixed forest; *Sorbus aucuparia* wood. **Kaczawskie Foothills:** Wleń 1.3 km N (WIF-31h), Wleń 1.6 km N (WIF-31g), Wleń 2.6 km N (WIF-25a); VII-VIII; deciduous forest, mixed forest; branch, *Quercus* sp. branch, deciduous wood.

***Plicaturopsis crispa* (Pers.) D.A. Reid (= *Plicatura crispa* (Pers.) Rea)**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Tarczyn 0.9 km N (WIF-134fx), Wleń 1.1 km SE (WIF-137i), Wleń 1.3 km SE (near WIF-137c); I, VIII; mixed forest, deciduous forest, oak-hornbeam forest, spruce forest; *Corylus avellana* branch, *Alnus incana* branch. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.4 km NW (WIF-46a); I-IV, X; deciduous forest, mixed forest; *Alnus incana* wood, *Corylus avellana* branch, *Fagus sylvatica* log, deciduous log. **Kaczawskie Foothills:** Belczyna 0.7 km S (WIF-123g), Bystrzyca 1.5 km E (WIF-126d), Bystrzyca 1.8 km E (WIF-126m), Bystrzyca 2.6 km NW (WIF-26j), Radomiłowice 1.1 km SW (WIF-11g), Sobota 1.6 km SW (WIF-24Aa), Sobota 2 km S (WIF-24g); I-IV, X-XI; deciduous forest, mixed forest, pine forest, spruce forest; wood, *Betula* sp. wood, *Corylus avellana* branch, *Padus avium* branch, *Prunus avium* branch, *Fagus sylvatica* log, deciduous log.

***Pluteus brunneoradiatus* Bonnard**

Izerskie Foothills: Wleń 1.9 km SW (ŁuF – Gniazdo Mt); VIII; mixed forest; wood. **Kaczawskie Foothills:** Belczyna 1.1 km E (near WIF-122a), Bystrzyca 1.5 km E (WIF-126d), Wleń 1.7 km SE (WIF-27g), Wleń 1.7 km NE (WIF-27h); IX-X; deciduous tree clump, mixed forest, spruce-pine forest, pine sapling stand; wood.

***Pluteus cervinus* (Schaeff.) P. Kumm. (= *P. atricapillus* (Batsch) Fayod)**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a), Bystrzyca 1.3 km SW (WIF-130h), Rząśnik 1.4 km SW (WIF-290k), Wleń 1.5 km E (WIF-137h); IX-X; deciduous forest, riparian forest, pine sapling stand; wood. **Izerskie Foothills:** Wleń 1.3 km SW (ŁuF – Gniazdo Mt), Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.4 km NW (WIF-46g), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.7 km SW (ŁuF – Gniazdo Mt), Wleń 1.8 km SW (ŁuF – Gniazdo Mt), Wleń 2 km SW (ŁuF – Gniazdo Mt), Wleń 2.8 km N (WIF-44j); VIII-X; oak forest, oak-hornbeam forest, riparian forest, wet spruce forest, deciduous forest, mixed forest; wood. **Kaczawskie Foothills:** Bystrzyca 1.4 km N (WIF-14d), Bystrzyca 1.5 km E (WIF-126d), Bystrzyca 2.2 km N (WIF-15c), Bystrzyca 2.4 km NW (WIF-21l), Sichów 2 km S (Gajka river valley), Wleń 1.4 km N (WIF-31b), Wleń 1.5 km N (WIF-30f), Wleń 2.3 km NE (WIF-25c); IV-X; oak-hornbeam forest, riparian forest, spruce forest, deciduous forest, mixed forest; litter, trunk base, wood.

***Pluteus cinereofuscus* J.E. Lange**

Kaczawskie Mts.: Wleń 1.1 km E (WIF-136f); IX; forest roadside; soil. **Izerskie Foothills:** Wleń 1.3 km NW (WIF-47s), Wleń 1.6 km NW (WIF-46f); VIII-IX; log landing, spruce forest; wood.

***Pluteus cyanopus* Quél.**

Kaczawskie Mts.: limestone quarry "Grudno"; IX; overgrown quarry; soil.

***Pluteus exiguus* (Pat.) Sacc.**

Kaczawskie Foothills: Przeździec 1.3 km W (WIF-24k); X; riparian forest; wood.

***Pluteus leoninus* (Schaeff.) P. Kumm.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; deciduous forest; wood. **Kaczawskie Foothills:** Bełczyna 1.1 km SE (WIF-122l), Radomiłowice 0.5 km SW (WIF-12a), Wleń 1.6 km NE (WIF-28f); VII-IX; oak forest, mixed forest; wood.

Pluteus nanus* (Pers.) P. Kumm. var. *nanus

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); IX; riparian forest; soil. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF-59b, 59d – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII-X; deciduous forest, mixed forest near a stream, riparian forest; wood.

***Pluteus pallescens* P.D. Orton**

Kaczawskie Mts.: Bystrzyca 1.2 km S (WIF-133a); IX; ash tree riparian forest; log. **Kaczawskie Foothills:** Bystrzyca 1.9 km N (WIF-13h); VII; under lime tree; litter.

***Pluteus petasatus* (Fr.) Gillet**

Kaczawskie Foothills: Wleń 1.6 km NE (WIF-28j); VIII; riparian forest; wood.

***Pluteus phlebophorus* (Ditmar) P. Kumm.**

Kaczawskie Mts.: Rzaśnik 1.3 km W (WIF-290a); X; oak forest; *Quercus* sp. wood. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX-X; deciduous forest, riparian forest; wood, litter.

***Pluteus plautus* (Weinm.) Gillet**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; deciduous forest; wood.

***Pluteus podospileus* Sacc. & Cub. f. *minutissimus* Wasser**

Izerskie Foothills: Wleń; IX; garden; soil. **Kaczawskie Foothills:** Przeździec 0.9 km S (WIF-22k); IX; riparian forest; wood.

Pluteus pouzarianus* Singer var. *pouzarianus

Kaczawskie Mts.: Bystrzyca 1.1 km S (LWl-132c), Rzaśnik 1.3 km W (WIF-290a); X; oak forest, mixed forest near a stream; wood, *Picea abies* log. **Kaczawskie Foothills:** Belczyna 0.5 km SW (WIF-123h), Górczyca 1.5 km W (WIF-24i), Wleń 2.4 km NE (WIF-26m), Wleń 2.8 km N (WIF-25d); VIII-X; spruce forest, mixed forest, willow forest; *Picea abies* stump and log, coniferous log, litter.

***Pluteus romellii* (Britzelm.) Sacc.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; litter. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF-59b), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF-64f); VI-X; deciduous forest, riparian forest; deciduous wood. **Kaczawskie Foothills:** Wleń 2.1 km N (WIF-30c); IX; sapling stand margin near a stream; ?*Salix* sp. trunk.

***Pluteus salicinus* (Pers.) P. Kumm.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a), Rzaśnik 2.3 km W (WIF-128g); IX-X; riparian forest, deciduous forest; deciduous wood. **Izerskie Foothills:** Wleń 1.9 km SW (ŁuF – Gniazdo Mt); VIII; mixed forest; *Betula* sp. log. **Kaczawskie Foothills:** Bystrzyca 1.8 km S (WIF-126b); X; between pine forest and oak forest; *Salix* sp. trunk.

***Pluteus semibulbosus* (Lasch) Quél.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-132c); VII; mixed sapling stand; *Alnus incana* wood. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2.8 km N (WIF-44j); VII-IX; oak-hornbeam forest, mixed forest; wood, *Fagus sylvatica* log.

Pluteus umbrosus* (Pers.) P. Kumm. var. *umbrosus

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; deciduous forest (logging road), riparian forest; wood. **Kaczawskie Foothills:** Przeździec 0.9 km S (WIF-22k); IX; riparian forest; wood.

***Polyporus badius* (Pers.) Schwein.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; spruce forest; *Sorbus aucuparia* branch. **Kaczawskie Foothills:** Bystrzyca 1.5 km N (WIF-18g), Bystrzyca 1.7 km N (WIF-16g), Bystrzyca 2.5 km NW (WIF-22o); VIII-X; deciduous forest, mixed forest, log landing; wood, deciduous wood.

***Polyporus brumalis* (Pers.) Fr.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2 km NW (WIF-46d); VIII-X; deciduous forest, pine forest; wood, *Betula* sp. branch. **Kaczawskie Foothills:** Belczyna 1 km SE (WIF-122f), Belczyna 1.5 km SE (WIF-124b), Bystrzyca 1.2 km SE (WIF-126n), near Wąwóz Lipy Res.; X-II; beech forest, riparian forest, set-aside; wood, *Betula* sp. branch, *Fagus sylvatica* branch.

***Polyporus ciliatus* Fr.**

Izerskie Foothills: Radomice (limestone quarry); V; overgrown quarry; *Betula* sp. log. **Kaczawskie Foothills:** Belczyna 0.9 km E (near WIF-122s), Bystrzyca 2.4 km NW (WIF-21m), Górczyca 1 km E (WIF-12Ao), Radomiłowice 1.1 km SW (WIF-11g), Ustronie 2.5 km W, Wleń 1.5 km N (WIF-31a), Wleń 1.9 km NE (WIF-28d); IV-XI; oak-hornbeam forest, birch forest, deciduous forest, mixed forest, leeve of a pond, spruce forest, wood, deciduous wood, *Tilia* sp. branch, ?*Quercus* sp. branch.

***Polyporus squamosus* (Huds.) Fr.**

Kaczawskie Mts.: Płoszczyńska 0.5 km W (PIF-253m); V; deciduous forest; *Fraxinus* sp. trunk. **Kaczawskie Foothills:** Bystrzyca 2.2 km N (WIF-13d); IX; deciduous forest; stump.

***Polyporus umbellatus* (Pers.) Fr.**

Kaczawskie Foothills: Bystrzyca 2.2 km N (WIF-15b), Wleń 1.4 km N (WIF-31c); VII-VIII; mixed forest; *Quercus* sp. roots, roots.

***Polyporus varius* (Pers.) Fr.**

Izerskie Foothills: Wleń 1.5 km SW (ŁuF – Gniazdo Mt), Wleń 2.1 km SW (ŁuF – Gniazdo Mt); V, VIII; mixed forest; wood. **Kaczawskie Foothills:** Radomiłowice 0.7 km SW (near WIF-11a); VII; deciduous forest; wood.

***Porodaedalea pini* (Brot.) Murrill (= *Phellinus pini* (Brot.) A. Ames)**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt); V; pine forest; *Pinus sylvestris* trunk. **Rudawy Janowickie Mts.:** Wielka Kopa Mt (Kolorowe Lakes); VIII; coniferous forest; *Pinus sylvestris* trunk.

***Porphyrellus porphyrosporus* (Fr.) E.-J. Gilbert**

Kaczawskie Mts.: Bystrzyca 1.5 km S (WIF-132); X; spruce forest; soil.

***Postia caesia* (Schr.) P. Karst. (= *Oligoporus caesius* (Schr.) Gilbertson & Ryvarden)**

Kaczawskie Foothills: Belczyna 0.7 km S (WIF-123g); X; spruce forest; *Picea abies* stump. **Izerskie Foothills:** Wleń 1.5 km SW (ŁuF – Gniazdo Mt); VIII; spruce forest; *Picea abies* log.

***Postia fragilis* (Fr.) Jülich (= *Oligoporus fragilis* (Fr.) Gilbertson & Ryvarden)**

Kaczawskie Foothills: Wleń 1.7 km NE (WIF-27g, 27h); IX; coniferous forest, pine forest; *Picea abies* stump, coniferous wood.

***Postia guttulata* (Peck) Jülich (= *Oligoporus guttulatus* (Peck) Gilbertson & Ryvarden)**

Kaczawskie Mts.: Bystrzyca 1.4 km SE (WIF-130g); VIII; mixed forest; *Picea abies* stump.

***Postia ptychogaster* (F. Ludw.) Vesterh. (= *Oligoporus ptychogaster* (Ludwig) R. & O. Falck)**

Kaczawskie Mts.: Bystrzyca 1.5 km S (WIF-133d); VIII; spruce forest; coniferous wood. **Izerskie Foothills:** Pilchowice (MaF-174a – near a Pilchowickie Lake); X; mixed sapling stand; stump. **Kaczawskie Foothills:** Bystrzyca 2.1 km N (WIF-14h), Bystrzyca 2.6 km NW (WIF-21d), Wleń 1.9 km NE (WIF-28d); IX-X; mixed forest, spruce forest; *Picea abies* branch, stump and log, log.

***Postia subcaesia* (A. David) Jülich**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); X; oak-hornbeam forest; *Alnus incana* log. **Kaczawskie Foothills:** Wleń 1.5 km N (WIF-31i); X; mixed forest; *Padus avium* log.

***Psathyrella candolleana* (Fr.) Maire**

Kaczawskie Mts.: Bystrzyca (WIF-134), Bystrzyca 1.2 km SE (WIF-131d); VII; mixed sapling stand, near a stream; litter. **Izerskie Foothills:** Wleń 1.4 km NW (WIF-46a), Wleń 1.5 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.7 km NW (WIF-46i), Wleń 1.8 km NW (WIF-45b); VII-IX; deciduous forest, mixed forest, logging road; wood, litter. **Kaczawskie Foothills:** Bełczyna 0.9 km E (near WIF-122s), Bystrzyca 2.2 km N (WIF-15a); VIII; mixed forest, ash tree sapling stand; wood, litter.

***Psathyrella cernua* (Vahl) G. Hirsch**

Kaczawskie Foothills: Bystrzyca 2 km S (WIF-126w); X; beech forest; *Fagus sylvatica* trunk.

***Psathyrella corrugis* (Pers.) Konrad & Maubl. (= *P. gracilis* (Fr.) Quél.)**

Kaczawskie Mts.: Bystrzyca 1.3 km SE (WIF-130c); IX; young deciduous forest; litter. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF-64d – Gniazdo Mt), Wleń 2 km SW (ŁuF – Gniazdo Mt); IX-X; beech forest, deciduous forest; litter. **Kaczawskie Foothills:** Bystrzyca 1 km SE (near WIF-126p), Bystrzyca 2.2 km N (WIF-14i), Bystrzyca 2.5 km NE (WIF-15h), Wleń 1.5 km NW (near WIF-31a), Wleń 2.6 km N (WIF-25a), between Górczyca and Przeździec (near WIF-24); IX-XII; deciduous forest, field, deciduous forest margin (nitrophilous vegetation), log landing; litter, soil.

***Psathyrella fagetophila* Örstadius & Enderle (= *P. murcida* (Fr.) Kits van Wav.)**

Kaczawskie Foothills: Bystrzyca 2.5 km NW (WIF-15k); X; deciduous sapling stand; litter.

***Psathyrella fatua* (Fr.) Konrad & Maubl.**

Izerskie Foothills: Wleń 1.6 km NW (WIF-46l); IV; deciduous forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.5 km NW (WIF-26); IV; birch forest; litter.

***Psathyrella impexa* (Romagn.) Bon**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-59a – Gniazdo Mt); X; riparian forest; litter. **Kaczawskie Foothills:** Wleń 2.5 km NE (WIF-26i); IX; mixed forest; litter.

***Psathyrella longicauda* P. Karst.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-132c); X; mixed forest; litter. **Kaczawskie Foothills:** Bystrzyca 2.5 km NE (WIF-15h); IX; deciduous forest; litter.

***Psathyrella lutensis* (Romagn.) Bon**

Kaczawskie Mts.: Buki Sudeckie Res.; X; *Carici remotae-Fraxinetum*; soil.

***Psathyrella microrrhiza* (Lasch) Konrad & Maubl.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); XI; riparian forest; litter. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 2.1 km SW (ŁuF – Gniazdo Mt); X-XI; riparian forest, spruce forest; litter.

***Psathyrella obtusata* (Pers.) A.H. Sm.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2.8 km N (WIF-44j); VI-XI; deciduous forest, riparian forest; wood, litter. **Kaczawskie Foothills:** Myślibórz (near Wąwóz Myśliborski Res.); XI; riparian forest; litter.

***Psathyrella olympiana* A.H. Sm.**

Kaczawskie Mts.: Bystrzyca 1.1 km SE (WIF-131g); IX; log landing; litter.

***Psathyrella orbicularis* (Romagn.) Kits van Wav.**

Kaczawskie Foothills: Bystrzyca 2.2 km N (WIF-14i), Myślibórz (near Wąwóz Myśliborski Res.); X-XI; riparian forest, log landing; litter.

***Psathyrella orbitarum* (Romagn.) M.M. Moser**

Kaczawskie Mts.: Buki Sudeckie Res.; X; *Carici remotae-Fraxinetum*; soil.

***Psathyrella panaeoloides* (Maire) Arnolds**

Kaczawskie Mts.: Płoszczyńska 0.5 km W (PiF-250n); V; meadow (log landing); litter. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF-58d); X; riparian forest; litter. **Kaczawskie Foothills:** Przeździec 1 km SW; X; roadside; litter.

***Psathyrella pennata* (Fr.) A. Pearson & Dennis**

Kaczawskie Foothills: Radomilowice 1.1 km SW (WIF-11h); VIII; spruce forest (old campfire); charcoal.

***Psathyrella pertinax* (Fr.) Örstadius**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a); IX; riparian forest; wood.

***Psathyrella piluliformis* (Bull.) P.D. Orton**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Wleń 2.8 km E (WIF-134z); IX-XI; oak-hornbeam forest, mixed forest; wood, stump. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); IX; riparian forest; stump. **Kaczawskie Foothills:** Bystrzyca 1.3 km SE (WIF-126k), Bystrzyca 2.2 km N (WIF-15c), Radomiłowice 1 km SW (near WIF-11b), Wleń 1.4 km NW (WIF-31b); IX-X; oak-hornbeam forest, deciduous forest, mixed forest; wood, stump.

***Psathyrella potteri* A.H. Sm.**

Kaczawskie Foothills: Bystrzyca 1.4 km N (WIF-17h); IX; oak-hornbeam forest; litter.

***Psathyrella prona* (Fr.) Gillet**

Kaczawskie Foothills: Bystrzyca 2.5 km NW (WIF-22p); X; forest margin; soil.

***Psathyrella pseudocorrugis* (Romagn.) Bon ss. Kits van Wav., Arnolds & Enderle**

Kaczawskie Foothills: Sikchów 2 km S (valley of the Gajka river); IV; riparian forest; litter.

***Psathyrella pseudogracilis* (Romagn.) M.M. Moser**

Kaczawskie Foothills: Wleń 1.9 km NE (WIF-28d); VIII; mixed forest; soil.

***Psathyrella spadiceogrisea* (Schaeff.) Maire**

Izerskie Foothills: Radomice (limestone quarry), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); V-VIII; riparian forest, overgrown quarry; soil. **Kaczawskie Foothills:** Belczyna 1.8 km SE (WIF-124a), Górczyca 1 km E (WIF-12Ao); IV-VI; birch forest, oak-hornbeam forest margin; litter, soil.

***Psathyrella supernula* (Britzelm) Örstadius & Enderle**

Kaczawskie Foothills: Wleń 2.6 km N (WIF-25a); X; log landing; litter.

***Psathyrella tenuicula* (P. Karst.) Örstadius & Huhtinen**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; *Sus scrofa* dung. **Izerskie Foothills:** Wleń 1.8 km NW (WIF-46i); VIII; spruce forest, *Sus scrofa* dung.

***Pseudoclitocybe cyathiformis* (Bull.) Singer**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1.1 km S (WIF-134a); X; oak-hornbeam forest, riparian forest; litter, soil. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); XI; deciduous forest; soil. **Kaczawskie Foothills:** Belczyna 1 km E (WIF-122b), Belczyna 1.3 km E (WIF-122d), Bystrzyca 2.5 km NW (WIF-22p), Przeździec 1.2 km E (near WIF-12); X; oak-hornbeam forest, deciduous forest, deciduous tree clump; litter, soil.

***Pseudoclitocybe expallens* (Pers.) M.M. Moser (= *P. obbata* (Fr.) Singer)**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry); X; grassland in quarry excavation; soil. **Izerskie Foothills:** Radomice (limestone quarry); XII; grassland in quarry excavation; soil. **Kaczawskie Foothills:** Przeździec 1.3 km S (quarry), Sobota 1.5 km S, Sobota 1.7 km SW (WIF-24Ab); X-XI; pasture, deciduous forest, melaphyre quarry; soil.

***Pseudocraterellus sinuosus* (Fr.) D.A. Reid (= *P. undulatus* (Pers.) Rauschert)**

Kaczawskie Mts.: Rząśnik 1.4 km SW (WIF-290k); IX; oak-hornbeam forest; soil. **Izerskie Foothills:** Wleń 1.3 km SW (ŁuF – Gniazdo Mt); VIII; mixed forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.8 km NW (WIF-21c), Wleń 2.6 km N (WIF-25a); VII-IX; oak-hornbeam forest, deciduous forest; soil.

***Pseudohydnum gelatinosum* (Scop.) P. Karst.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a), Tarczyn 0.9 km NE (WIF-134c); X-XI; riparian forest, spruce forest; *Pinus sylvestris* wood, *Picea abies* stump. **Izerskie Foothills:** Wleń 1.4 km NW (WIF-46g); VIII; mixed forest; coniferous stump. **Kaczawskie Foothills:** Bystrzyca 1.9 km N (WIF-14j), Bystrzyca 2.1 km N (WIF-14h); X; mixed forest, spruce forest; *Picea abies* wood.

***Psilocybe semilanceata* (Fr.) P. Kumm.**

Izera Mts.: Świeradów-Zdrój (ŚF-371a); X; grassy roadside; soil. **Karkonosze Mts.:** Szklarska Poręba (SzPF-347c); IX; roadside; soil. **Kaczawskie Foothills:** Radomiłowice 0.8 km W (near WIF-12Ar); IX; wet meadow; soil.

***Psilocybe serbica* M.M. Moser & E. Horak var. *arcana* (Borovicka & Hlaváček) Borovicka, Oborník & Noordel.**

Kaczawskie Mts.: near Bystrzyca (near WIF-134dx); riparian forest; litter. **Izerskie Foothills:** Pilchowice 1.1 km S (MaF – Dziki Valley); XI; logging road near a stream; litter (*Picea abies* branch).

***Psilocybe serbica* M.M. Moser & E. Horak var. *bohémica* (Šebek) Borovička, Oborník & Noordel.**

Lubawska Gate: Lubawka; XI; deciduous forest, litter.

***Pterula subulata* Fr. (= *P. multifida* (Chev.) Fr.)**

Kaczawskie Mts.: Bystrzyca 1 km SE (WIF-132a), Góra Miłek Res.; IX-X; spruce forest (old campfire), *Cephalanthero-Fagetum*; litter, soil.

***Pycnoporus cinnabarinus* (Jacq.) P. Karst.**

Kaczawskie Mts.: Bystrzyca 1.4 km SE (WIF-130f), Tarczyn 0.8 km N (near WIF-134/136), Tarczyn 1.2 km NE (WIF-133f), Tarczyn 1.8 km E (near WIF-133f), Wleń 1.3 km SE (WIF-137c); I-VIII; mixed forest, spruce forest, tree clump; ?*Quercus* sp. wood, *Betula* sp. branch, *Padus avium* branch, *Populus tremula* branch.

Izerskie Foothills: Wleń 1.4 km NW (WIF-46g), Wleń 2 km NW (WIF-46d); VIII; mixed forest, pine forest; deciduous wood. **Kaczawskie Foothills:** Bystrzyca 1.9 km N (WIF-13h), Bystrzyca 2.5 km NW (WIF-15k), Bystrzyca, Wleń 1.4 km N (WIF-33c), Wleń 1.5 km N (WIF-31a), Wleń 1.5 km NW (at the Bóbr river near WIF-31a), Wleń 1.5 km NE (near WIF-33), Wleń 1.6 km NW (WIF-28f); II-X; deciduous forest, mixed forest, deciduous sapling stand, roadside, spruce forest, riverside tree clump, roadside tree clump; *Padus avium* wood, deciduous wood, *Padus avium* branch, ? *Populus tremula* branch, *Betula* branch and trunk.

***Radulomyces confluens* (Fr.) M.P. Christ.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF-64a – Gniazdo Mt); X; riparian forest; ?*Quercus* sp. branch.

Ramaria abietina* (Pers.) Quél. var. *abietina

Kaczawskie Foothills: Belczyna 0.7 km S (WIF-123g), Bystrzyca 2.1 km N (WIF-14h); IX-X; mixed forest, young spruce forest (old campfire), spruce forest; charcoal, soil. **Sudetes Foreland:** Szklary 0.5 km S; IX; mixed forest; soil.

***Ramaria apiculata* (Fr.) Donk**

Izerskie Foothills: Wleń 1.8 km SW (ŁuF – Gniazdo Mt); IX; mixed forest; litter. **Rudawy Janowickie Mts.:** Łomnica; IX; mixed forest; soil.

***Ramaria eumorpha* (P. Karst.) Corner**

Kaczawskie Foothills: Bystrzyca 2.5 km NW (WIF-21l), Radomiłowice 0.7 km SW (WIF-11f), Wleń 1.6 km NE (WIF-27j); VIII-IX; spruce forest margin, mixed forest, spruce forest; litter, soil.

***Ramaria decurrens* (Pers.) R.H. Petersen**

Kaczawskie Mts.: Góra Milek Res.; IX; *Cephalanthero-Fagetum*; litter, soil.

Ramaria fennica* (P. Karst.) Ricken var. *fennica

Kaczawskie Mts.: Góra Milek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Ramaria flaccida* (Fr.) Bourdot**

Kaczawskie Foothills: Belczyna 1.8 km SE (WIF-124d), Radomiłowice 1 km SE (WIF-11f), Wleń 3 km NE (WIF-22r); VII-IX; deciduous forest, spruce forest; litter, soil.

***Ramaria flavescens* Schaeff. ex R.H. Petersen**

Kaczawskie Foothills: Wąwóz Lipy Res.; XI; oak forest; soil.

***Ramaria formosa* (Pers.) Quél.**

Kaczawskie Mts.: Bystrzyca 0.8 km SE (WIF-127a); VIII; oak-hornbeam forest; soil.

***Ramaria lutea* Schild**

Kaczawskie Mts.: Góra Miłek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Ramaria mairei* Donk**

Kaczawskie Mts.: Góra Miłek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Ramaria ochracea* (Bres.) Corner**

Kaczawskie Mts.: Rząśnik 2.7 km W (WIF-128b); VIII; mixed forest; deciduous wood.

***Ramaria stricta* (Pers.) Quél. var. *concolor* Corner**

Kaczawskie Mts.: Rząśnik 1.5 km W (WIF-290m); oak forest; wood.

Ramaria stricta* (Pers.) Quél. var. *stricta

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1.5 km SE (WIF-130c), Rząśnik 1.5 km W (WIF-290m), Góra Miłek Res.; VIII-XI; *Cephalanthero-Fagetum*, oak-hornbeam forest, mixed forest; wood, ?soil.

Karkonosze Mts.: Szklarska Poręba (SzPF-347b); IX; mixed forest; wood. **Izerskie Foothills:** Maciejowiec (MaF – Dziki Valley); X; mixed forest; wood. **Kaczawskie Foothills:** Bystrzyca 1 km E (WIF-126s), Radomiłowice 1.5 km NW (WIF-15c), Sobota 1.7 km SW (WIF-24Ab), Wleń 1.5 km NW (near WIF-31a); IX-XI; deciduous forest, forest margin; wood, ?soil.

***Ramariopsis citrina* Schild**

Kaczawskie Foothills: Bystrzyca 2.2 km N (WIF-14b); IX; mixed forest; soil.

***Ramariopsis crocea* (Pers.) Corner**

Kaczawskie Foothills: Bystrzyca 1.4 km NW (WIF-19g), Wleń 1.6 km NE (WIF-27d); IX-X; deciduous forest, beech forest, clearing; soil.

***Ramariopsis kunzei* (Fr.) Corner**

Kaczawskie Mts.: Buki Sudeckie Res., Góra Miłek Res., Miłek Mt; IX-X; *Cephalanthero-Fagetum*, *Dentario enneaphylli-Fagetum*, meadow; soil. **Izerskie Foothills:** Maciejowiec; X; lawn; soil.

***Ramariopsis pulchella* (Boud.) Corner**

Kaczawskie Mts.: Góra Miłek Res.; IX; *Cephalanthero-Fagetum*; soil.

***Ramariopsis subtilis* (Pers.) Corner s.l. (= *Clavulinopsis subtilis* (Pers.) Corner)**

Kaczawskie Mts.: Nowe Rochowice, Buki Sudeckie Res., Góra Miłek Res.; VIII-IX; *Cephalanthero-Fagetum*, *Dentario enneaphylli-Fagetum*; soil.

***Ramariopsis tenuiramosa* Corner**

Kaczawskie Mts.: Buki Sudeckie Res.; IX; *Dentario enneaphylli-Fagetum*; soil.

***Resupinatus applicatus* (Batsch) Gray**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII-XI; deciduous forest, mixed forest; wood. **Kaczawskie Foothills:** Bełczyna 0.7 km S (WIF-123b), Bystrzyca 1.9 km N (WIF-18d); VII-I; oak-hornbeam forest, deciduous forest; wood.

***Rhizopogon roseolus* (Corda) Th.M. Fr.**

Kaczawskie Mts.: Miłek Mt; VIII; overgrown limestone quarry; soil.

***Rhodocollybia butyracea* (Bull.) Lennox f. *asema* (Fr.) Antonín, Halling & Noordel.**

Kaczawskie Foothills: Bystrzyca 0.7 km E (WIF-20); XI; spruce forest; soil.

Rhodocollybia butyracea* (Bull.) Lennox f. *butyracea

Kaczawskie Mts.: Wleń 2.8 km E (WIF-134w); XI; mixed forest; litter

***Rhodocybe caelata* (Fr.) Maire**

Izerskie Foothills: Maciejowiec (MaF – Dziki Valley); X; mixed forest; soil.

***Rickenella fibula* (Bull.) Raithelh.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); VI; mixed forest; soil, mosses. **Kaczawskie Foothills:** Bełczyna 0.8 km N (WIF-122m); IX; deciduous sapling stand; soil.

***Rickenella swartzii* (Fr.) Kuyper (= *R. setipes* (Fr.) Raith.)**

Izerskie Foothills: Łupki (near tourist hostel); X; lawn (near apple-tree, spruce and larch); soil. **Kaczawskie Foothills:** Bystrzyca 1.2 km SE (WIF-126s), Wleń 2.6 km N (WIF-25a); VII-IX; deciduous forest, riparian forest; soil.

***Rigidoporus sanguinolentus* (Alb. & Schwein.) Donk (= *Physisporinus sanguinolentus* (Alb. & Schwein.) Pilát)**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF-64k – Gniazdo Mt); IX; deciduous forest; *Picea abies* branch.

***Rimbachia neckerae* (Fr.) Redhead**

Kaczawskie Foothills: Wleń (butchery site); IX; mosses.

***Ripartites tricholoma* (Alb. & Schwein.) P. Karst. s.l.**

Kaczawskie Mts.: Rzaśnik 2.3 km W (WIF-128c), Tarczyn 0.6 km E (near WIF-134), Wleń 1.4 km NE (WIF-136b); X-XI; deciduous forest, mixed forest, tree clump; soil. **Kaczawskie Foothills:** Bełczyna 0.6 km S (WIF-123g), Bełczyna 1.8 km SE (WIF-124a), Bystrzyca 1.5 km E (WIF-126d), Bystrzyca 1.9 km N (WIF-18d), Górczyca 1.1 km W (WIF-24d), Sobota 1.6 km SW (WIF-24Aa), Wleń 1.9 km N (WIF-30d); IX-XI; coniferous forest, deciduous forest, mixed forest, pine forest, alder forest, spruce forest; soil.

***Roridomyces roridus* (Scop.) Rexer (= *Mycena rorida* (Fr.) Quél.)**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); IX; oak-hornbeam forest; wood. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IX; mixed forest; litter. **Kaczawskie Foothills:** Bystrzyca 2.1 km N (WIF-14h); IX; mixed forest; *Picea abies* cone.

***Rugosomyces carneus* (Bull.) Bon (= *Calocybe carnea* (Bull.) Donk)**

Kaczawskie Mts.: Tarczyn 0.6 km E; X; slate excavation; soil. **Izerskie Foothills:** Wleń (Południowe Es.); VII-IX; lawn; soil. **Kaczawskie Foothills:** Bystrzyca 1.5 km NE (near WIF-131); X; meadow; soil.

***Russula grisescens* (Bon & Gaugué) Marti**

Izerskie Foothills: Zalesie 0.8 km N (LuF-200h); VIII; mire (mixed sapling stand); soil.

***Russula lepida* Fr. (= *R. rosea* Pers.)**

Kaczawskie Foothills: Przeździec 0.8 km SW (WIF-221); IX; mixed forest; soil.

***Russula mustelina* Fr.**

Kaczawskie Foothills: Wleń 1.7 km NE (WIF-27h); IX; spruce forest; soil.

***Russula nauseosa* (Pers.) Fr.**

Kaczawskie Foothills: Wleń 2.3 km NE (WIF-25c); IX; spruce forest; soil.

***Russula nigricans* (Bull.) Fr.**

Kaczawskie Foothills: Wleń 2.6 km N (WIF-25a); IX; deciduous forest; soil.

***Russula ochroleuca* Pers.**

Izerskie Foothills: Wleń 1.5 km SW (ŁuF – Gniazdo Mt); VIII; mixed forest (under birch); soil.

***Russula pseudoraoultii* Ayel & Bidaud**

Izerskie Foothills: Zalesie 0.8 km N (LuF-200h); VIII; mire (mixed sapling stand); soil.

***Russula rhodopoda* Zvára**

Izerskie Foothills: Zalesie 0.8 km N (LuF-200h); IX; mire (mixed sapling stand); soil.

***Russula sardonia* Fr.**

Kaczawskie Foothills: Wleń 1.7 km NE (WIF-27g); X; pine sapling stand; soil.

***Russula undulata* Velen.**

Izerskie Foothills: Zalesie 0.8 km N (LuF-200h); VIII; mire (mixed sapling stand); soil.

***Russula violeipes* Quél.**

Kaczawskie Foothills: Radomiłowice 1 km SW (WIF-11b); IX; mixed forest; soil.

***Russula virescens* (Schaeff.) Fr.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; soil.

***Sarcodon imbricatus* (L.) P. Karst.**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF-59c – Gniazdo Mt); VIII-X; wet spruce forest; soil.

***Sarcomyxa serotina* (Pers.) P. Karst. (= *Panellus serotinus* (Schrad.) Kühner)**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1.3 km SE (WIF-130c), Bystrzyca 1.6 km SW (WIF-134gx), Tarczyn 1.2 km NE (WIF-134n); XI-I; deciduous forest, oak-hornbeam forest, riparian forest; wood, *Alnus* sp. trunk. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.4 km NW (WIF-47k); XI; deciduous forest, riparian forest; ?*Salix* sp. log, wood. **Kaczawskie Foothills:** Bełczyna 1.1 km SE (WIF-123m), Radomiłowice 0.7 km SW (WIF-12a), Sobota 1.7 km SW (WIF-24Ab), near Wąwóz Myśluborski Res. towards Myślubórz; X-XI; deciduous forest, mixed forest, riparian forest; wood.

***Schizophyllum amplum* (Lév.) Nakasone (= *Auriculariopsis ampla* (Lév.) Maire)**

Kaczawskie Foothills: Bystrzyca 1.2 km SE (WIF-126n), Bystrzyca 2.4 km NW (WIF-21o, 22), Przeździec 1.2 km E, Radomiłowice 1.4 km W (WIF-12Ak), Sobota 1.5 km S; VIII-I; mixed forest, riparian forest, overgrown pasture; *Populus tremula* wood. **Sudetes Foreland:** Szklary 0.5 km SW; I-III; overgrown spoil tip, overgrown excavation; deciduous wood, *Populus tremula* wood.

***Schizophyllum commune* Fr.**

Kaczawskie Foothills: Górczyca 1.5 km W (WIF-24l), Wleń 2.6 km N (WIF-25a); III, VII; willow clump at the margin of oak-hornbeam forest, deciduous forest; *Salix* sp. wood, *Tilia* sp. branch.

***Schizopora flavipora* (Berk. & M.A. Curtis ex Cooke) Ryvarden (= *Hyphodontia flavipora* (Cooke) Sheng H. Wu)**

Kaczawskie Foothills: Przeździec 1.7 km E (near WIF-12), Wleń 1.1 km N (WIF-31k); I, X; deciduous forest, tree clump near a stream; wood, *Alnus* sp. branch.

***Schizopora paradoxa* (Schrad.) Donk (= *Hyphodontia paradoxa* (Schrad.) E. Langer & Vesterholt s.l.)**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); II; blackthorn shrubs; blackthorn branch.

Kaczawskie Foothills: Bełczyna 1.3 km E (WIF-122d), Bystrzyca 1.9 km N (WIF-18d), Bystrzyca 2.1 km NW (WIF-26c), Bystrzyca 2.7 km NW (near WIF-26f); X-I; riparian forest, oak-hornbeam forest, deciduous forest; *Quercus* sp. branch, *Corylus avellana* branch.

***Schizopora radula* (Pers.) Hallenb. (= *Hyphodontia radula* (Pers.) E. Langer & Vesterholt)**

Kaczawskie Foothills: Bełczyna 0.8 km E (near WIF-122); XI; riparian forest; ?*Fraxinus* sp. log.

***Scleroderma areolatum* Ehrenb.**

Izerskie Foothills: Wleń 1.2 km NW (WIF-46g); VIII; mixed forest; soil. **Kaczawskie Foothills:** Bełczyna 1.8 km (WIF-126b); X; mixed wood margin; soil.

***Scleroderma bovista* Fr.**

Kaczawskie Foothills: Wleń 1.9 km NE (WIF-28d); VIII; mixed forest; soil.

***Scleroderma cepa* Pers.**

Izerskie Foothills: Wleń 1.6 km NW (WIF-47f); IX; deciduous forest; soil.

***Scleroderma citrinum* Pers.**

Kaczawskie Foothills: Wleń 2.6 km N (WIF-25a); VII; deciduous forest; soil.

***Scleroderma verrucosum* (Bull.) Pers.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); X; oak-hornbeam forest; soil.

***Sebacina incrustans* (Pers.) Tul. & C. Tul.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; oak-hornbeam forest; mosses, litter. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII-XI; oak-hornbeam forest, deciduous forest; mosses, litter.

***Serpula himantoides* (Fr.) P. Karst.**

Kaczawskie Mts.: Bystrzyca 1.5 km SE (WIF-130d), Bystrzyca 1.7 km SE (WIF-130j), Tarczyn 1.2 km NE (WIF-133f); III-XII; mixed forest, spruce forest; deciduous wood, *Picea abies* log. **Izerskie Foothills:** Wleń 2 km SW (ŁuF – Gniazdo Mt, Białe Skały); X; pine forest; *Picea abies* log. **Kaczawskie Foothills:** Bystrzyca 2.1 km N (WIF-14h), Bystrzyca 2.2 km N (WIF-14d), Wleń 1.3 km N (WIF-31h); X-XI; mixed forest, spruce forest; ?deciduous wood, *Picea abies* log, singed ?*Picea abies* wood.

***Simocybe centunculus* (Fr.) Singer var. *centunculus* (= *Ramicola centunculus* (Fr.) Watling)**

Kaczawskie Foothills: Wleń 1.5 km N (WIF-31h); VIII; deciduous forest; wood.

***Simocybe sumptuosa* (P.D. Orton) Singer (= *Ramicola sumptuosa* (P.D. Orton) Watling)**

Kaczawskie Foothills: Bystrzyca 2.8 km NW (WIF-21c); VII; deciduous forest; wood.

***Sistotrema confluens* Pers.**

Sudetes Foreland: Szklary 0.5 km W; X; overgrown spoil tip; ?soil.

***Skeletocutis amorpha* (Fr.) Kotl. & Pouzar**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); IV; mixed forest; *Picea abies* stump. **Kaczawskie**

Foothills: Wleń 1.4 km N (WIF-31b); IV; mixed forest; *Pinus sylvestris* trunk.

***Skeletocutis carneogrisea* A. David**

Kaczawskie Mts.: Bystrzyca 1.6 km SE (WIF-131i); XII; mixed forest; *Picea abies* log. **Izerskie Foothills:**

Wleń 1.6 km SW (ŁuF – Gniazdo Mt); II; mixed forest; *Picea abies* log.

***Skeletocutis nivea* (Jungh.) Jean Keller**

Kaczawskie Foothills: Bystrzyca 2.5 km NE (WIF-15h), Górczyca 1.1 km W (WIF-24g), near Wąwóz Lipy Res.; VII-XI; riparian forest, deciduous forest, mixed forest; *Fraxinus* sp. log, *Sorbus aucuparia* branch. **Sudetes**

Foreland: Szklary 0.5 km SW; IX; overgrown outcrop; wood.

***Skeletocutis odora* (Peck ex Sacc.) Ginns**

Kaczawskie Mts.: Czernica 1.2 km S (PiF-241); IV; oak-hornbeam forest; *Larix* sp. stump.

***Sparassis crispa* (Wulfen) Fr.**

Izerskie Foothills: Wleń 1.3 km SW (ŁuF – Gniazdo Mt), Wleń 1.9 km SW (ŁuF – Gniazdo Mt), Wleń 2 km NW (WIF-46c); VIII-IX; mixed forest, pine forest; *Pinus sylvestris* roots, stump and trunk base. **Kaczawskie Foothills:** Bystrzyca 2.1 km N (WIF-14h), Bystrzyca 2.5 km NW (WIF-26f), Przeździec 0.8 km SW (WIF-22l), Wleń 0.9 km NW (WIF-32b), Wleń 1.4 km N (WIF-31c), Wleń 1.7 km NE (WIF-27h); VIII-X; oak forest, coniferous forest, mixed forest, pine forest; *Pinus sylvestris* roots.

***Sphaerobolus stellatus* Tode**

Kaczawskie Foothills: Wleń 2.6 km N (WIF-25a); VI; deciduous forest; deciduous stump.

***Steccherinum bourdotii* Saliba & A. David (= *Irpex bourdotii* (Saliba & A. David) Kotiranta & Saarenoksa)**

Kaczawskie Mts.: Wleń 1.4 km SE (WIF-137h); I; margin of pine sapling stand; *Corylus avellana* branch.

Kaczawskie Foothills: Bystrzyca 2.2 km N (WIF-13d), Radomiłowice 0.7 km W (near WIF-12Af), Radomiłowice 1.1 km SW (WIF-11g), Rząśnik 1.3 km NW (WIF-125g), Wleń 2 km N (WIF-29h), Wleń 2.7 km NE (WIF-26d); XI-III; deciduous forest, mixed forest, spruce forest; branch, *Quercus* sp. branch, *Alnus incana* branch, *Padus avium* branch.

***Steccherinum fimbriatum* (Pers.) J. Erikss. (= *Irpex fimbriatus* (Pers.) Kotiranta & Saarenoksa)**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); XI; deciduous forest; wood. **Kaczawskie Foothills:** Bełczyna 0.8 km E (near WIF-122), Bystrzyca 1.9 km N (WIF-18d), Bystrzyca 2 km NE (WIF-15s), Radomiłowice 0.8 km W (WIF-12Ad); XI-I; riparian forest, oak-hornbeam forest, mixed forest; deciduous log, *Alnus incana* branch, *Fraxinus* sp. branch, *Tilia* sp. branch.

***Steccherinum ochraceum* (Pers. ex J.F. Gmel.) Gray (= *Irpex ochraceus* (Pers.) Kotiranta & Saarenoksa)**

Kaczawskie Mts.: Rząśnik 1.3 km W (WIF-290a), Tarczyn 0.7 km N (WIF-134gx); I, IX; mixed forest, oak forest; deciduous wood, *Quercus* sp. branch. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); II; deciduous forest, riparian forest; *Quercus* sp. branch, *Corylus avellana* branch. **Kaczawskie Foothills:** Bystrzyca 2.1 km NW (near WIF-26c), Wleń 1.5 km NW (near WIF-31a); XII-I; forest margin, mixed forest; *Quercus* sp. branch.

***Stereum gausapatum* (Fr.) Fr.**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); VIII; mixed forest; *Quercus* sp. stump.

***Stereum hirsutum* (Willd.) Pers.**

Kaczawskie Foothills: Bełczyna 1.5 km SE (near WIF-124b); X; set-aside (meadow near a forest); deciduous stump (?birch, ?alder).

***Stereum ochraceoflavum* (Schwein.) Sacc.**

Kaczawskie Foothills: Radomiłowice 1.1 km SW (WIF-11g); III; mixed forest; *Quercus* sp. branch.

***Stereum rugosum* Pers.**

Kaczawskie Foothills: Bystrzyca 2.1 km NW (near WIF-26f), Bystrzyca 2.8 km NW (WIF-22o); I, X; deciduous forest; *Alnus* sp. log, stump.

***Stereum sanguinolentum* (Alb. & Schwein.) Fr.**

Kaczawskie Foothills: Bełczyna 1.8 km SE (WIF-124d); II; mixed forest; *Pinus sylvestris* branch.

***Stereum subtomentosum* Pouzar**

Izerskie Foothills: Wleń 1.4 km NW (WIF-47k), Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII-IX; deciduous forest; ?*Quercus* sp. log, *Alnus incana* branch. **Kaczawskie Foothills:** Bełczyna 0.9 km E (WIF-122t), Bystrzyca 1.2 km SE (WIF-126n), Przeździec 1.4 km E (WIF-12f), near Nad Groblą Res.; I, VII-XI; deciduous forest, riparian forest; *Alnus incana* branch, *Alnus* sp. log.

***Strobilomyces strobilaceus* (Scop.) Berk.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Wleń 1.1 km E; VIII-IX; oak-hornbeam forest, deciduous forest; soil. **Kaczawskie Foothills:** Wleń 1.4 km N (WIF-31c, 31f), Wleń 1.5 km N (WIF-30i), Wleń 1.6 km NE (WIF-27k), Wleń 2.1 km NW (WIF-30b), Wleń 2.7 km N (WIF-25b); VIII-IX; oak forest, deciduous forest, mixed forest, spruce forest; soil.

***Strobilurus esculentus* (Wulfen) Singer**

Kaczawskie Mts.: Wleń 2.3 km NE (WIF-136b); IX; mixed forest; *Picea abies* cone. **Izerskie Foothills:** Wleń 1.3 km NW (WIF-47o); XI; mixed forest; ?*Picea abies* cone. **Kaczawskie Foothills:** Bystrzyca 2.2 km N (WIF-13d), Rząśnik 1.3 km NW (WIF-125g); X-XI; birch forest, mixed forest; cone.

***Strobilurus stephanocystis* (Hora) Singer**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt); V; mixed forest; *Pinus sylvestris* cone.

***Strobilurus tenacellus* (Pers.) Singer**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.4 km NW (WIF-46g); VI-VII; spruce-pine forest, spruce forest; *Pinus sylvestris* cone. **Kaczawskie Foothills:** Bystrzyca 2.6 km NW (WIF-21d), Leszczyna 1 km N (near Ciche Szczęście mine), Wleń 1.9 km NE (WIF-28d); IV-VI; oak-hornbeam forest, mixed forest, spruce forest; *Pinus sylvestris* cone, *Picea abies* cone.

***Stropharia aeruginosa* (Curtis) Quél. (= *Psilocybe aeruginosa* (M.A. Curtis) Noordel.**

Kaczawskie Mts.: Rząśnik 1.5 km W (WIF-290m), Wleń 1.5 km E (WIF-137h); X; mixed forest, pine sapling stand; litter, soil. **Kaczawskie Foothills:** Bystrzyca 1.8 km W (WIF-126m), Bystrzyca 1.9 km N (WIF-13h), Wleń 1.7 km NE (WIF-27g, 28k); X-XI; mixed forest, pine sapling stand, spruce forest; litter, soil.

***Stropharia cyanea* (Bull.) Tuom. (= *Psilocybe caerulea* (Kreisel) Noordel.**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-131b), Tarczyn 1.1 km E (WIF-133f), Miłek Mt; X-XI; meadow, riparian forest, spruce forest; litter, soil. **Kaczawskie Foothills:** Wleń 2.8 km N (WIF-25b); XI; deciduous forest; soil.

***Stropharia coronilla* (Bull.) Quél. (= *Psilocybe coronilla* (Bull.) Noordel.)**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); IX; deciduous forest (roadside); soil.

***Stropharia semiglobata* (Batsch) Quél. (= *Psilocybe semiglobata* (Batsch) Noordel.)**

Karkonosze Mts.: Karkonosze NP (Szrenica Mt); IX; mountain pine shrubs; soil. **Kaczawskie Mts.:** Bystrzyca 1 km S (WIF-133a); IX; riparian forest; ?soil.

***Stropharia rugosoannulata* Murrill f. *rugosoannulata* (= *Psilocybe rugosoannulata* (Farlow ex Murrill) Noordel.)**

Kaczawskie Foothills: Bełczyna 0.8 km SW (near WIF-11), Bystrzyca 2.4 km NW (WIF-22o); VIII-X; field, mixed forest (log landing); soil.

***Suillus bovinus* (L.) Roussel**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); X; mixed forest; soil. **Izerskie Foothills:** Zalesie 0.6 km N (LuF-199g), Zalesie 0.8 km N (LuF-199g); IX-X; mixed forest near mire, pine forest; soil.

***Suillus cavipes* (Opat.) A.H. Sm. & Thiers f. *aureus* (Rolland) Singer**

Karkonosze Mts.: Szklarska Poręba (Złoty Potok Valley); XI; spruce forest (under larch); soil. **Izerskie Foothills:** Wleń 2 km NW (WIF-46d), Zalesie 0.8 km N (LuF); IX-X; roadside (under larch and pine), mire margin (pine sapling stand), pine forest; soil. **Kaczawskie Foothills:** Bystrzyca 2.1 km N (WIF-14i); mixed forest; soil.

***Suillus cavipes* (Opat.) A.H. Sm. & Thiers f. *cavipes* (= *Boletinus cavipes* (Klotzsch ex Fr.) Kalchbr.)**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry); IX; overgrown limestone quarry; soil. **Izerskie Foothills:** Wleń 1,5-1.6 km SW (LuF – Gniazdo Mt); VII-IX; spruce forest; soil. **Kaczawskie Foothills:** Przeździec 0.8 km SW (WIF-22l); IX; mixed forest; soil.

***Suillus collinitus* (Fr.) Kuntze**

Izerskie Foothills: Radomice (limestone quarry); IX-X; overgrown quarry; soil.

***Suillus flavidus* (Fr.) J. Presl**

Izerskie Foothills: Zalesie 0.8 km N; X; mire; soil.

***Suillus luteus* (L.) Roussel**

Karkonosze Mts.: Karkonosze NP (Równia pod Śnieżką – 0.45 km W from “Dom Śląski” tourist hostel); IX; mountain pine shrubs; soil.

***Suillus tridentinus* (Bres.) Singer**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry); VIII-X; overgrown quarry (under larch); soil.

***Suillus variegatus* (Sw.) Kuntze**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); VIII; mixed forest; soil. **Izerskie Foothills:** Zalesie 0.8 km N (LuF-200h); VIII-X; dry mire margin, mixed sapling stand; soil.

***Suillus viscidus* (L.) Roussel (= *S. aeruginascens* (Opat.) Snell.)**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry), Nowe Rochowice (limestone quarry), Góra Miłek (limestone quarry); VIII-X; overgrown quarry; soil. **Karkonosze Mts.:** Szklarska Poręba (Złoty Potok Valley); XI; spruce forest (under larch); soil. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF-59c – Gniazdo Mt); VIII, X; wet spruce forest (under larch); soil.

***Tapinella panuoides* (Fr.) E.-J. Gilbert f. *panuoides* (= *Paxillus panuoides* (Fr.) Fr.)**

Kaczawskie Mts.: Bystrzyca 1.6 km SE (WIF-131i), Wleń 1.3 km NE (WIF-136b); VIII-IX; mixed forest; ?*Quercus* sp. wood, ?*Picea abies* stump. **Izerskie Foothills:** Wleń 1.5 km SW (ŁuF – Gniazdo Mt); VIII-IX; spruce forest; *Picea abies* stump. **Kaczawskie Foothills:** Wleń 1.7 km NE (WIF-27h); IX; spruce-pine forest; *Picea abies* stump.

***Thelephora anthocephala* (Bull.) Fr.**

Kaczawskie Mts.: Tarczyn 0.9 km NE (WIF-134c); VIII; spruce forest; soil.

***Thelephora caryophyllea* (Schaeff.) Pers.**

Sudetes Foreland: Szklary 0.5 km SW; IX-X; overgrown excavation; soil.

***Thelephora palmata* (Scop.) Fr.**

Kaczawskie Mts.: Rząśnik 1.4 km SW (WIF-290k); IX; deciduous forest; soil. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); IX; oak-hornbeam forest; soil.

***Thelephora penicillata* (Pers.) Fr.**

Kaczawskie Mts.: Rząśnik 1.4 km SW (WIF-290k); IX; deciduous forest; litter. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; riparian forest; soil.

***Thelephora terrestris* Ehrh.**

Izera Mts.: Sępik Mt (ca. 900 m a.s.l.); X; spruce forest; *Picea abies* stump. **Kaczawskie Mts.:** Rząśnik 3 km SW (near WIF-135); X; slate excavation; soil, litter, wood. **Kaczawskie Foothills:** Wleń 1.4 km NW (WIF-31b); mixed forest; soil, litter, wood.

***Trametes gibbosa* (Pers.) Fr.**

Izerskie Foothills: Wleń 2 km SW (ŁuF – Gniazdo Mt); VII; spruce forest; *Fagus sylvatica* stump. **Kaczawskie Foothills:** Bystrzyca 2.4 km NW (WIF-211); III; deciduous forest; *Alnus incana* log.

***Trametes hirsuta* (Wulfen) Pilát**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); VII; deciduous forest; deciduous log. **Kaczawskie Foothills:** Przeździec 1.7 km E (near WIF-12); X; tree clump near a stream; wood.

***Trametes ochracea* (Pers.) Gilb. & Ryvarden**

Kaczawskie Foothills: Bystrzyca 1.2 km SE (WIF-126n), Bystrzyca 1.5 km N (WIF-18g), Wleń 2 km N (WIF-29d); I, X; mixed forest, coniferous forest, riparian forest; log, *Acer pseudoplatanus* log, *Quercus* sp. branch.

***Trametes pubescens* (Schumach.) Pilát**

Kaczawskie Mts.: Wleń 1.1 km E (near WIF-137i); I; deciduous forest; *Populus tremula* log. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF-64a – Gniazdo Mt); XI; deciduous forest; ?*Alnus incana* log. **Kaczawskie Foothills:** Bystrzyca 1.4 km N (WIF-16k), Bystrzyca 1.6 km N (WIF-16h), Bystrzyca 2.1 km NW (near WIF-26f), Bystrzyca 2.4 km NW (near WIF-22/26), Bystrzyca 2.5 km NE (WIF-15h), Sobota 1.7 km SW (WIF-24Ab); X-III; oak-hornbeam forest, deciduous forest, mixed forest; *Fagus sylvatica* branch, *Populus tremula* branch, ?*Alnus* sp. log, *Padus avium* log, deciduous log, wood.

***Trametes suaveolens* (L.) Fr.**

Sudetes Foreland: Szklary 0.3 km S; III; tree clump; *Salix* sp. stump.

***Trametes versicolor* (L.) Lloyd**

Kaczawskie Foothills: Wleń 2.6 km N (WIF-25a); XI; oak-hornbeam forest; *Corylus avellana* branch.

***Trechispora mollusca* (Pers.) Liberta**

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); XI; riparian forest; *Acer pseudoplatanus* wood.

***Tremella globospora* D.A. Reid**

Izerskie Foothills: Wleń 1.3 km SW (ŁuF – Gniazdo Mt), Wleń 1.4 km SW (ŁuF – Gniazdo Mt); II-V; oak-hornbeam forest, deciduous forest; branch (*Pyrenomyces stromata*). **Kaczawskie Foothills:** Przeździec 1.7 km E (WIF-12k); I; deciduous forest; *Quercus* sp. branch (*Pyrenomyces stromata*).

Tremella mesenterica* (Schaeff.) Retz. f. *mesenterica

Izerskie Foothills: Wleń 1.4 km SW (ŁuF – Gniazdo Mt); VI; deciduous forest; *Fraxinus* sp. wood.

***Trichaptum hollii* (J.C. Schmidt) Kreisel (= *T. fuscoviolaceum* (Ehrenb.) Ryvarden)**

Kaczawskie Mts.: Bystrzyca 1.4 km SE (WIF-130f); II; spruce forest; *Picea abies* bark.

***Tricholoma argyraceum* (Bull.) Gill.**

Izerskie Foothills: Radomice (limestone quarry); IX; overgrown quarry; soil.

***Tricholoma boudieri* Barla**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); X; oak forest; soil.

***Tricholoma cingulatum* (Almfelt) Jacobasch**

Kaczawskie Mts.: Mysłów (Mysłów-Sobocin quarry); XII; overgrown limestone quarry; soil. **Izerskie Foothills:** Radomice (limestone quarry); XII; overgrown quarry (under aspen); soil.

***Tricholoma columbetta* (Fr.) P. Kumm.**

Kaczawskie Foothills: Wąwóz Lipy Res.; XI; oak forest; soil.

***Tricholoma frondosae* Kalamees & Shchukin**

Kaczawskie Mts.: Tarczyn 0.6 km E (Tarczynka Mt); XI; overgrown hill; soil. **Izerskie Foothills:** Radomice (limestone quarry); X; overgrown quarry (under aspen); soil. **Kaczawskie Foothills:** Bełczyna 0.6 km S (near WIF-123h), Bełczyna 0.7 km S (WIF-123g), Bełczyna 0.9 km E (WIF-122a), Sobota 1.5 km S; X-XI; young spruce forest (under aspen), coniferous forest (under aspen), overgrown pasture (under aspen); soil. **Sudetes Foreland:** Szklary 0.6 km S; IX; overgrown meadow; soil.

***Tricholoma imbricatum* (Fr.) P. Kumm.**

Kaczawskie Foothills: Bełczyna 1.2 km E (WIF-122c), Wleń 1.7 km NE (WIF-27g); X; young spruce forest, young pine forest; soil.

***Tricholoma orirubens* Quél.**

Kaczawskie Mts.: Grudno (Bukowa Mt); IX; *Cephalanthero-Fagetum*; soil.

***Tricholoma pardinum* (Pers.) Quél.**

Kaczawskie Mts.: Wleń 1.1 km E (WIF-136g); X; spruce forest; soil.

***Tricholoma portentosum* (Fr.) Quél.**

Kaczawskie Mts.: Tarczyn 1.2 km NE (WIF-133f); X; mixed forest; soil.

***Tricholoma psammopus* (Kalchbr.) Quél.**

Izera Mts.: Świeradów-Zdrój (ŚFD-238k); X; mixed forest, under larch; soil. **Kaczawskie Foothills:** Wleń 1.9 km NE (WIF-28d); IX; mixed forest; soil.

Tricholoma saponaceum* (Fr.) P. Kumm. var. *saponaceum

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a), Bystrzyca 1 km S (WIF-132c); VIII, X; oak forest, spruce forest; soil.

***Tricholoma sculpturatum* (Fr.) Quél.**

Kaczawskie Mts.: Tarczyn 0.6 km E (Tarczynka Mt); XI; overgrown hill; soil. **Kaczawskie Foothills:** Bystrzyca 1.8 km E (WIF-126m); pine sapling stand; soil.

***Tricholoma sciodes* (Pers.) C. Martín**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); X; mixed forest; soil.

***Tricholoma stiparophyllum* (N. Lund) P. Karst.**

Izerskie Foothills: Wleń 1.5 km SW (ŁuF – Gniazdo Mt); X; mixed forest; soil.

***Tricholoma vaccinum* (Schaeff.) P. Kumm.**

Izera Mts.: Świeradów-Zdrój (ŚFD-238f), Kobyła Mt (N slope); X; coniferous forest, spruce forest; soil.

***Tricholomopsis decora* (Fr.) Singer**

Karkonosze Mts.: Karkonosze NP (Łabski Kocioł); IX; spruce forest; *Picea abies* wood. **Izerskie Foothills:** Zalesie 0.6 km N (LuF); IX; pine forest; *Pinus sylvestris* wood.

***Tricholomopsis rutilans* (Schaeff.) Singer**

Kaczawskie Foothills: Bystrzyca 0.7 km E (WIF-20ax), Wleń 1.4 km N; X; mixed forest; stump.

***Tubaria conspersa* (Pers.) Fayod**

Kaczawskie Mts.: Bystrzyca 1 km S (WIF-133a), Bystrzyca 1.1 km S (WIF-133c); IX-XI; beech forest, riparian forest; litter. **Izerskie Foothills:** Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 2 km SW (ŁuF – Gniazdo Mt); VIII; mixed forest, spruce-pine forest; deciduous twigs. **Kaczawskie Foothills:** Przeździec 0.8 km SW (WIF-22l), Sobota 1.7 km SW (WIF-24Ab), Wleń 2.6 km N (WIF-25a); VII-X; deciduous forest; litter, *Populus tremula* wood.

***Tubaria furfuracea* (Pers.) Gillet s.str.**

Kaczawskie Mts.: Bystrzyca 1.1 km SE (WIF-131f), Bystrzyca 1.1 km S (WIF-133c); XI-XII; beech forest, log landing; woodchips, litter. **Kaczawskie Foothills:** Belczyna 1.8 km SE (WIF-124a), Bystrzyca 1.4 km E (WIF-126k), Radomiłowice 0.7 km SW (WIF-11f), Wleń 2.7 km NE (near WIF-25b); IX-XI; meadow near a forest, between oak forest and spruce forest, riparian forest, log landing; litter.

***Tubaria hiemalis* Romagn. ex Bon**

Izerskie Foothills: Radomice (limestone quarry); XII; overgrown quarry; litter. **Kaczawskie Foothills:** Belczyna 1 km SE (WIF-122f), Bystrzyca 2 km NE (WIF-15s), Bystrzyca 2.1 km NW (WIF-26c), Bystrzyca 2.5 km NW (WIF-15k), Radomiłowice 0.8 km W (WIF-12Ar); XII-II; deciduous forest, mixed forest, between spruce forest and deciduous tree clump; litter.

***Tubaria minutalis* Romagn.**

Kaczawskie Foothills: Bystrzyca 0.5 km N (WIF-20l); X; basalt quarry; litter.

***Tubaria romagnesiana* Arnolds**

Kaczawskie Foothills: Bystrzyca 2.2 km N (WIF-15c); IX; oak-hornbeam forest; litter.

***Tulasnella violea* (Quél.) Bourdot & Galzin**

Kaczawskie Foothills: Bystrzyca 1.4 km N (WIF-19h); I; mixed sapling stand; *Populus* sp. log.

***Tyromyces kmetii* (Bres.) Bondartsev & Singer**

Izerskie Foothills: Wleń 1.6 km SW (ŁuF – Gniazdo Mt); VIII; deciduous forest; *Alnus incana* branch.

***Volvariella bombycina* (Schaeff.) Singer**

Kaczawskie Foothills: between Belczyna and Radomiłowice; VIII-IX; roadside; *Populus* sp. trunk.

***Volvariella gloiocephala* (DC.) Boekhout & Enderle**

Kaczawskie Mts.: Bystrzyca 1 km SE (WIF-131a); IX; log landing; sawdust. **Kaczawskie Foothills:** Bystrzyca 0.5 km N (WIF-20l), Bystrzyca 1 km SE (near WIF-126p), Bystrzyca 2.2 km N (WIF-15a/14i), Górczyca 0.7 km W, Rząśnik 1.2 km NW (near WIF-125), Wleń (near sewage treatment plant); IX-XI; farmland, log landing, stubble field, overgrown basalt excavation; woodchips, soil.

***Volvariella hypopithys* (Fr.) M.M. Moser**

Izerskie Foothills: Radomice (limestone quarry); VIII; overgrown quarry; soil.

***Xerocomus badius* (Fr.) E.-J. Gilbert**

Karkonosze Mts.: Szklarska Poręba (Złoty Potok Valley); XI; spruce forest; soil.

***Xerocomus pelletieri* (Lév.) Bresinsky & Manfr. Binder (= *Phylloporus rhodoxanthus* (Schwein.) Bres.)**

Kaczawskie Foothills: Bystrzyca 1.2 N (WIF-17l), Górczyca 1.4 km W (WIF-24f), Radomiłowice 1.4 km W (WIF-12Ai); VIII-IX; beech forest, deciduous forest, lime forest; soil.

***Xerocomus pruinatus* (Fr. & Hök) Quél.**

Karkonosze Mts.: Karkonosze NP (SzPF-186a); IX; spruce forest; *Picea abies* wood. **Izerskie Foothills:** Wleń 1.4 km SW (ŁuF – Gniazdo Mt); X; deciduous forest; soil.

***Xerocomus rubellus* (Krombh.) Quél.**

Kaczawskie Foothills: Bystrzyca 1.5 km N (WIF-18g), Bystrzyca 2.4 km NW (WIF-21l); VIII-IX; deciduous forest, mixed forest; soil. **Izerskie Foothills:** Wleń 1.5 km SW (ŁuF – Gniazdo Mt), Wleń (Południowe Es.); VII-VIII; garden (lawn), spruce forest; soil.

***Xerocomus subtomentosus* (L.) Quél.**

Kaczawskie Mts.: Bystrzyca 0.8 km S (WIF-127a); IX; mixed forest; soil.

***Xeromphalina brunneola* O.K. Mill.**

Izera Mts.: Sępik Mt (N slope); IX; spruce forest; wood.

***Xeromphalina campanella* (Batsch) Kühner & Maire**

Izerskie Foothills: Wleń 1.3 km SW (ŁuF – Gniazdo Mt), Wleń 1.6 km SW (ŁuF – Gniazdo Mt), Wleń 1.8 km SW (ŁuF – Gniazdo Mt); IV-X; mixed forest, spruce forest; wood. **Kaczawskie Foothills:** Wleń 2.4 km N (WIF-25c); VII; mixed forest; wood.