Cytogenetic Investigations on *Anemone* L. I. *Anemone Janczewskii* a New Amphidiploid Species of Hybrid Origin

by

W. Gajewski

(Plates III—IV)

I. Introduction

The chief suggestion to the present paper were the highly interesting conclusions of Janczewski's researches on the interspecific hybrids in the genus *Anemone*. This author has published in the years 1889—1892 the results of his vast investigations on *Anemone* hybrids. Following points drew especially my attention:

1. The hybrids among *Anemone silvestris*, *A. multifida* and *A. virginiana* present intermediate characters between the parental species and are completely or almost completely sterile. Only the hybrids between *Anemone silvestris* and *A. multifida* v. *magellanica* have produced twice in two years a quite fertile shoot from a sterile plant. On the other shoots of this hybrid, which were sterile, some good achenes were formed every year. The plants raised from the good achenes, both from fertile and sterile shoots, have formed an uniform second hybrid generation.

2. The plants of the second generation were fertile and breed true in following generations. Basing on these observations, Janczewski has stated that this fertile and constant hybrid merits a rank of a new species. He wrote: „une nouvelle espèce végétale (dans l'acception rigoureuse de ce mot) peut être créée par la voie de „hybridité” (Janczewski, 1892).

These highly interesting statements were recognised by geneticists only in the last years. Renner (1929) and Skalińska (1935) have suggested that the results of Janczewski's experiments present the first carefully described case of the apparition of a constant

and fertile species hybrid, probably on the way of the doubling of the chromosome number.

In Janczewski's experiments the cytological side is entirely missing. Nothing certain can be stated concerning the mode of apparition of fertile hybrids without cytological investigations.

I have attempted to repeat Janczewski's crosses, and in the case of obtaining fertile hybrids, to test their chromosome numbers and meiotic divisions. Unfortunately I have not obtained the production of fertile shoots, but the production of unreduced gametes as the cause of the apparition of good achenes on the sterile shoots has been stated beyond any doubt. The results of the present experiments have demonstrated that the doubling of the chromosome number is the cause of fertility and constancy of new hybrid form. This new species of hybrid origin is named *Anemone Janczewskii*.

During my researches it has been obtained from the Botanical Gardens of Edinburgh, Freiburg, Stäfa and Dublin the seeds of *Anemone*, named *A. magellanica* and *A. decapetala*, from which I have raised the plants quite identical with the experimentally produced *A. Janczewskii*. It seems then, that *Anemone Janczewskii* has probably arisen in the Botanical Gardens, where the parental species are grown together and can intercross easily.

This work was carried in the Botanical Garden in Warsaw. I am much indebted to prof. B. Hryniewiecki under whose guidance this work was carried, to prof. R. Kobendza for helpful advice and for preparation of all photographs included in this paper. I wish also to express my thanks to prof. K. Piech for the permission to use his library and to dr. S. Krupko and H. Teleżyński for many valuable advices in cytological technique.

II. Material and methods

Plant material was partly obtained from the Botanical Garden in Warsaw and partly from seeds obtained from foreign Botanical Gardens. *Anemone silvestris* was collected in nature in Podolia and cultivated in Botanical Garden in Warsaw. *A. multifida* was grown from seeds. *A. Janczewskii*, described in this paper, was cultivated from seeds obtained from Botanical Gardens in Edinburgh, Freiburg and Dublin under the name *A. magellanica* and from the Botanical Garden Stäfa under the name *A. decapetala*. All these species belong, according to Ulbrich (1916), to the section *Eriocephalus* Hook. et Thoms. This section contains about 20 species with a wide

distribution over Europe, Asia and America, the species are characterised by elongated receptaculum and densely, woolly achenes.

The cytological studies were made chiefly from permanent preparations. The root tips were fixed in Navaschin fixative modified by Langlet (1927). The floral buds before fixation in Langlet were forfixed for 1—2 minutes in Carnoy fixative. The somatic chromosomes were studied only from root tips and meiotic chromosomes from P. M. C. The root tips preparations were cut 20μ thick and the floral buds — 25μ thick. All material was stained by Newton's gentian-violet technique. Aceto-carminine smears were also sometimes used predominantly from anthers and rarely also from root-tips according to the „Kochmethode" of Heitz.

The pollen grains and the stomata were studied in water preparations. The pollen grains were stained in aceto-carminine.

Drawings were made with a camera lucida at magnification of 3000 or 1700, and were reduced to one-half this size in reproduction.

III. The description of *Anemone silvestris*, *A. multifida* and their hybrids

1. *Anemone silvestris* L. $2n = 16$. Perennial, the roots very long and creeping, produce abundantly adventitious shoots. The stem very short, branched and ramping on the earth. The basal leaves longpetioled, 3-parted, the divisions broadly cuneate-oblong, variously cleft and divided into serrate lobes on both sides sparsingly and shortly haired. Those of involucre very similar. The floral shoot usually with involucre of three leaves and terminated with one flower. The floral buds conic, on hanging peduncles which straight again only before anthesis. Sepals usually 5, ovate, obtuse, ca. 28 mm. long and 19 mm. broad, white, often with traces of violet tint on the outer side and silky hairy. The stamina very numerous with gold-yellow anthers. Pollen grains with three furrows when expanded circular, ca. 20μ in diameter. Achenes on short pedicels, cylindrical, densely woolly and tipped with short subulate styles. The more precise biometric data bearing on this species are given together with the description of his hybrid with *A. multifida*.

2. *Anemone multifida* Poir. $2n = 32$. Perennial, the root system with a long principal root and a mass of lateral roots growing oblique down into the earth. Adventitious shoots from roots very rare and only from the base of the principal root. The stem short, deep branched and ramping on the earth. Basal leaves longpetioled, 3-parted, the

divisions cleft into linear acute lobes, 2—4 mm. broad with entire margins. Those of involucre shortpetioled and more or less cuneate, otherwise similar. The leaves are pale-green and densely hairy. Floral shoots with 3 leaves in involucre, branched usually with 2—5 flowers, the lateral peduncles involucre with 2 leaves. Floral buds globoid on erect peduncles. Sepals 5—10 oblong, white, greenish, rose or purple, 1,5—1,8 cm. long and 0,7—0,9 cm. broad, from outer side silky sometimes with a claw. The stamina less numerous than in *A. silvestris* with greenish-yellow anthers. Pollen grains with twelve furrows, when expanded circular ca 30 μ in diameter. Achenes sessile, compressed, densely woolly and tipped with the short erect style.

In connection with a very large distribution of *A. multifida*, ranging from north Canada to the Maghellan Strait through the whole American Continent, this species form numerous forms, some of which can be regarded as geographical races. According to Janeczewski (1899) this species possesses three chief varieties: 1) *var magellanica* — 25 cm. high, leaves small narrowly incised, sepals pale yellow (Sulphur-yellow nr 1/3 according to Horticultural Colour Cart), 18 mm long and 9 mm. broad. Less deeply hairy than others varieties. 2) *var alba* — to 40 cm. high, sepals white often with greenish, yellow or rose tint, 15 mm. long and 7 mm. broad. 3) *var hudsoniana* — 20—30 cm. high, sepals purple (Solferino-purple nr 16 according to Horticultural Colour Cart) often with a small claw on the base, 13—14 mm. long and 5—8 mm broad. More precise biometric data bearing on this species are given together with the description of his hybrid with *Anemone silvestris*.

3. *F*₁ *Anemone silvestris* x *A. multifida* var. *magellanica*

This species hybrid is strictly perennial like *A. silvestris*. It was grown already 5 years without any traces of degeneration. From year to year it is more robust and in the year 1940 it has produced 69 floral shoots on one exemplar, what I never found in the parental species. In the vegetative growth it shows very distinct heterosis and in height greatly exceeds both parental species (Fig. nr 7). It produces new leaves until the first frosts, like *A. silvestris*. It flowers in May together with *A. multifida* and a little later than *A. silvestris*, in autumn there appear new floral shoots every year. Besides a compact group of roots growing oblique down into the earth there exist also horizontal growing roots with adventives buds. This is a mixture of both parental characters. The basal leaves are intermediate in their

shape (Fig. nr 1), between the two parental species. They are greater however in all parts in comparison with both parental species. The mean values of the petiole length and the length of the central blade segment are as follows:

<i>A. silvestris</i>	length of the petiole	13,5cm	length of the blade	6cm
$F_1 A. silv. \times A. mult.$	„ „ „ „	19,0cm	„ „ „ „	6,5cm
<i>A. multifida</i>	„ „ „ „	6,0cm	„ „ „ „	4,0cm

The size of leaves and also of shoots is very variable in different years and the heterosis is not always distinct. The leaves of

Fig. 1. Basal leaves of *A. silvestris* (left), $F_1 A. silvestris \times A. multifida$ v. *magellanica* (middle) and *A. multifida* v. *magellanica* (right).

involucres are also intermediate in shape like the basal-ones (Fig. nr 2). The hairs on whole the plant are intermediente in length and density between the parental species.

Very interesting is the distribution of the stomata by parental species and their hybrid. Miczyński (1892), a coworker of Janczewski, has described on Janczewski's material the following relations in stomatal distribution: by *A. silvestris* stomata are only on the lower side of the leaves, 60 on one sq. mm of the surface, by *A. multifida* they are on both sides of the leaves, 50 on one sq. mm of the lower side surface and 28 — on the upper side. By the hybrid they are 13 stomata on one sq. mm of the upper side and 55 on the lower side, which is quite exactly an arithmetic midway between the

two parental species. My own observations have confirmed Miczyński's results but the values, that I have obtained, are quite different. Firstly, it has been stated, that the parental species differ not only

Fig. 2. Involucral leaves of *A. silvestris* (left), F_1 *A. silvestris* $\times$ *A. multifida* v. *magellanica* (middle) and *A. multifida* v. *magellanica* (right).

in the number of stomata on 1 sq. mm but also in the shape of their stomata (the length/breadth index), in the size of stomata and in the stomatal density (the relation between the number of stomata and the number of epidermal cells on surface unity). The stomatal density were calculated as the percent of stomata to the general sum

of epidermal cells on surface unity, counting each stomata as one cell. This is the „stomatal index” of Salisbury. According to Salisbury stomatal index is a constant value for every plant species. The results of my measurments are given in the tables nr 1 and 2.

TABLE 1. Size of the stomata in μ

Name of the plant	μ 27 30 33 36 39 42 45 48 51 54 57 60 63															n	M	m
<i>A. silv.</i> lgth. bread.			1	19	45	26	8	1							100	38,2	0,25	
		1	7	38	41	12	1								100	36,2	0,25	
F_1 lgth. bread.			1	2	7	18	44	21	7						100	43,2	0,30	
		17	53	23	5	2									100	31,4	0,25	
<i>A mult.</i> lgth. bread.								8	18	30	38	4	2		100	53,0	0,30	
			6	24	44	16	10								100	37,4	0,20	

TABLE 2. Number of stomata and epidermal cells on 1 sq. mm.

Name of plant	Side of leaf	Number of stomata	Number of epid. cells	Stomatal index
<i>Anemone silvestris</i>	upper	0	544,8	0%
	lower	168,7	537,6	23,8%
F_1 <i>silvestris</i> $\times$ <i>multifida</i>	upper	29,2	416,1	6,5%
	lower	91,5	387,0	19,1%
<i>Anemone multifida</i>	upper	42,9	378,9	10,1%
	lower	71,5	361,7	16,5%

Fig. 3. Epidermis from the lower side of the leaf of *A. silvestris* ($\times 520$).

From the tables 1 and 2 and from Figs. 3, 4 and 5 it is evident that the F_1 hybrid possesses in comparison with parental species an

Fig. 4. Epidermis from the lower side of the leaf of F_1 *A. silvestris* $\times$ *A. multifida* v. *magellanica* ($\times 520$).

Fig. 5. Epidermis from the lower side of the leaf of *A. multifida* v. *magellanica* ($\times 520$).

intermediate number, size and shape of the stomata. Also the value of his stomatal index is intermediate. The number of stomata and epidermal cells on 1 sq. mm of leaf surface are by the hybrid closer to *A. multifida* than to *A. silvestris*. The elongated shape of stomata

by *A. multifida* is dominant over the circular shape of stomata by *A. silvestris*. The length/breadth index is by *A. silvestris* 1, 11, by *A. multifida* — 1,41 and by F_1 — hybrid — 1,37. By the measure-

Fig. 6. Floral shoots of *A. silvestris* (left), F_1 *A. silvestris* $\times$ *A. multifida* v. *magellanica* (middle) and *A. multifida* v. *magellanica* (right).

ments of stomatal size and number great care must be taken to compare merely the leaves of exactly the same state of development.

By *A. silvestris* the floral shoot has one flower or rarely two, by *A. multifida* v. *magellanica* it has 2—3 or rarely 4—5 flowers and the hy-

brid between them produces on one shoot from 1 to 4 flowers (table nr 3).

TABLE 3. The number of flowers on one shoot.

Numb. of. flowers	1	2	3	4	5	n	M	m
<i>A. silvestris</i>	46	9				55	1,1	0,04
F ₁	2	29	16	1		48	2,3	0,08
<i>A. mult. v. magell.</i>		25	33	5	1	64	2,7	0,08

As we see the number of flowers is by the hybrid intermediate when compared to parental species. The height of floral shoots by the hybrid greatly exceeds the height of the floral shoots of his parents (Figs. 6 a. plate III). The measurments made in 1938 have given following results:

<i>A. silvestris</i>	length of the floral shoots (mean)							35 cm
F ₁	"	"	"	"	"	"	"	42 cm
<i>A. mult. v. magell.</i>	"	"	"	"	"	"	"	26 cm

The floral buds are on erect peduncles like by *A. multifida*, this character dominates completely over the dropping peduncles of *A. silvestris*. Also the colour of the sepals is identical with the colour of the paternal species (Sulphur-yellow 1/3 according to Horticultural Colour Cart). The number of sepals by the hybrid is quite exactly intermediate as the table 4 shows.

TABLE 4. The number of sepals.

Number of sepals	4	5	6	7	8	9	10	11	n	M	m
<i>A. silvestris</i>	4	179	20	6	1				210	5,1	0,03
F ₁		24	25	16	5				70	6,0	0,1
<i>A. mult. v. magell.</i>		1	7	21	40	7	1	1	78	7,7	0,08

Also the length and breadth of sepals by the hybrid is quite intermediate, in comparison with parental species as it is evident from the data of tables 5 and 6 and from the fig. 7.

TABLE 5. Length of the sepals in mm.

Name of plant	Length of sepals in mm.																																				M	m
	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	n								
<i>Anemone silvestris</i>														2	15	17	21	22	30	33	43	61	69	42	40	22	15	12	5	4	453	28,2	0,15					
F ₁ <i>A. silvestris</i> x <i>multifida</i>																															266	21,4	0,12					
<i>Anemone multifida</i>	1	1	9	15	25	27	31	46	47	52	31	19	17	6																	327	15,9	0,13					

TABLE 6. Breadth of the sepals in mm.

Name of plant	Breadth of sepals in mm.																											M	m	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27			n
<i>Anemone silvestris</i>													5	5	6	12	18	22	32	45	50	35	26	21	4	3	3	287	19,8	0,16
F ₁ A. <i>silvestris</i> x <i>multifida</i>					2	9	10	20	36	38	39	51	30	18	8	4	1											266	11,3	0,13
<i>Anemone multifida</i>	2	—	4	11	21	60	67	90	51	19	2																327	7,8	0,08	

TABLE 7. Pollen grains diameter in μ .

Name of plant	7,5 9 10,5 12 13,5 15 16,5 18 19,5 21 22,5 24 25,5 27 28,5 30 31,5 33 34,5 36 37,5																											M	m																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																	
						2	4	34	83	69	4																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																			

The length/breadth index of the sepals in *A. silvestris* is 1, 4, in *A. multifida* — 2,0 and in the hybrid 1,9 which is very near to paternal value. By *A. silvestris* it appears in the later phases of growth a great amount of anthocyanin both in vegetative parts and in sepals giving violet tint to them; this anthocyanin by *A. multifida* v. *magellanica* and by the hybrid is entirely lacking. In other parts of the flower the two parental species differ slightly. The number of stamina is greater by *A. multifida* than by *A. silvestris* and by the hybrid the number is similar to paternal species. The colour of anthers is gold-yellow by *A. silvestris* and greenish-yellow by *A. multifida*, by the hybrid the anthers are shrunken and nearly empty and are of grayish colour.

Fig. 7. Sepals of *A. silvestris* (top row), *A. multifida* v. *magellanica* (bottom row) and their F_1 hybrid (middle row).

The pollen grains of both parental species differ greatly in size and number of furrows. *A. silvestris* possesses smaller pollen grains with three furrows, *A. multifida* has greater pollen grains with twelve furrows. When a grain is quite dry it is of an elongate form, but when it is expanded the grain becomes ellipsoidal and nearly circular in outline. The pollen grains were measured when expanded and stained in aceto-carmin. In such preparations both nuclei are quite clearly visible of which one is more globular and the second discoid in outline and more deeply staining. By *A. multifida* the nuclei are greater together with greater dimensions of pollen grains in comparison with *A. silvestris*. The length of the longer axis of the discoid nucleus is $6,7\mu$ by *A. silvestris* and $9,9\mu$ by *A. multifida*. By the hybrid the pollen grains are quite completely degenerated without nuclei, with an unstained or very weak stained plasm,

sometimes entirely empty or with the chromatin mass agglomerated in an amorphous body or in numerous little droplets. The pollen grains are not or little expanded and of very different size and shape with a variable number of furrows. Very numerous are quite little

Fig. 8. Pollen grains from left to right of *A. silvestris*, *A. multifida*, F_1 hybrid and F_2 amphidiploid.

Fig. 9. One viable pollen grain and numerous degenerated from the F_1 hybrid *A. silvestris* $\times$ *A. multifida* v. *magellanica*.

grains with 1—2 furrows, together with the increasing dimensions of pollen grains increases also the number of furrows ranging from 1 to 12. The well stained pollen grains with a good appearance are very rare and form only 0,3% of the total pollen grains. They possess 12 furrows like *A. multifida* but are much greater (figs. 8 a. 9) In the table 7 are put together the results of the measurements of pollen grains diameters.

The diameter of good pollen grains of the hybrid is greater than the diameter of pollen grains of the tetraploid parent. This unexpected fact suggests the supposition that the good pollen of this hybrid possess unreduced number of chromosomes. As a matter of fact it was really demonstrated later.

The fertility of this hybrid is very low. *A. silvestris* and *A. multifida* are quite fully fertile and give only few bad achenes. Following self-pollination the hybrid sets no seeds at all and by open pollination only few seeds with good appearance were produced. The rarity of good seeds indicates that the functioning embryo-sacs are like pollen grains very rare. In the autumn of 1937 this hybrid has produced some flowers and has set 6 good achenes in absence of any other flowering *Anemone* species, what proofs that the good achenes are set as the result of selfing or parthenogenetic development. The selfpollination is possible as the anthers burst and the pollen are thrown, in spite of its quite bad quality. The fertility of the hybrid and of the parental species is represented in the table 8.

TABLE 8. The fertility of hybrid and parental species.

	Season	Nmb. of head	Nmb. of achen.	Nmb. of good achen	%
<i>A. silvestris</i>	sum. 36	8	1002	965	96,4
<i>A. mult. v. mag.</i>	sum. 36	10	1461	1362	93,3
F_1 hybrid	aut. 36	4	486	0	0,0
	sum. 37	44	5093	30	0,6
	aut. 37	7	810	6	0,7
	sum. 38	46	5115	5	0,1
	sum. 39	23	2270	5	0,2

Besides the quite good achenes full and hard, the hybrid produces also some achenes which are shrunken and soft and make impression that the fertilization has taken place but the embryonal development has been soon arrested. The achenes of *A. silvestris* and of *A. multifida* differ both in shape and size. The achenes of the hybrid are intermediate in shape although more similar to *A. multifida*. They are however greater in all dimensions than the achenes of both parental species, as it is presented in table 9.

The length of achenes given on table 9 is measured without the persistent styles on the top of achenes. The greater dimensions of the hybrid achenes, as compared with parental ones, are probably due to the fact that the embryo in all germinating seeds was found to be hexaploid, what has been proved by cytological analysis of

plants raised from these achenes. The ratio length/breadth of the achenes is 3,0 for *A. silvestris*, 1,7 for *A. multifida* v. *magellanica* and 1,9 for their hybrid. This value is intermediate between the parents but nearer to the paternal species. The germinating power of the seemingly good achenes of the hybrid is very low. From the

TABLE 9. Dimensions of achenes in mm.

<i>Anemone silvestris</i>	length	$M=2,32$	$m=0,02$	$n=115$
	bread.	$M=0,75$	$m=0,015$	$n=115$
	thick.	$M=0,26$	$m=0,007$	$n=115$
F ₁ <i>A. silvestris</i> x <i>A. multifida</i>	length	$M=3,24$	$m=0,04$	$n=50$
	bread.	$M=1,64$	$m=0,02$	$n=50$
	thick.	$M=0,5$	$m=0,01$	$n=50$
<i>Anemone multifida</i>	length	$M=2,7$	$m=0,018$	$n=115$
	bread.	$M=1,57$	$m=0,018$	$n=115$
	thick.	$M=0,49$	$m=0,011$	$n=115$

30 achenes harvested in summer of 1937 and sown immediately after ripening have germinated only 12 of which 2 have died in cotyledon stage and only 8 have been raised to maturity. As these plants were cultivated in rather unfavourable conditions, it is possible, that in optimal conditions the percentage of mature plants would be somewhat greater. Nevertheless the germination of these achenes is disturbed.

IV. The description of amphidiploids with $2n=48$

The 8 plants of the second generation of the cross *A. silvestris* x *A. multifida* v. *magellanica* have a double number of chromosomes in comparison to F₁ maternal plants. The F₁ hybrid has $2n=24$ and the F₂ plants have $2n=48$ or nearly so. Morphologically these plants were quite uniform and differed greatly from F₁ plants. The first impression was that they are greater and bigger in all parts.

A careful comparison of these plants with anemones received from Edinburgh and Freiburg under the name *Anemone magellanica* has demonstrated that they are in all details identical. They are identical not only in their morphological traits but also in their chromosome number, which is 48 in root tips. Only few plants have a somewhat lower somatic number of chromosomes ranging from 42 to 47. It is highly probable that the hexaploids obtained from the Botanical Gardens have originated in a similar way as our amphidiploids, i. e. by the doubling of the chromosome number on somatic or gametic way in the hybrids between *A. silvestris* and *A. multifida*.

For comparison I have tabulated below some biometric values from amphidiploids experimentally obtained together with hexaploids obtained from Edinburgh. They are very similar and sometimes quite identical. The description of amphidiploids is based exclusively on the 8 amphidiploids obtained from my cross but it concerns also the hexaploids obtained from the Botanical Gardens. Janczewski has cultivated his constant hybrid through 3 generations and I have cultivated the amphidiploids to the third generation, and like Janczewski, I never found any trace of segregation. This hybrid-form quite constant and considerably fertile merits

Fig. 10. Basal and involucral leaves and sepals of triploid F_1 hybrid (right) and his hexaploid descendants in F_2 (left).

a specific rank, as has already stated Janczewski. It differs greatly in nearly all characters and chromosome number from the parental species from which it is separated also by barriers of cross-sterility. Since Janczewski has not given any name to his new „good” species, I propose to name them *Anemone Janczewskii*. The description given below can be regarded as his diagnosis.

Anemone Janczewskii $2n=48$. Perennial, besides a big central root growing vertically it is a mass of lateral roots growing oblique down into the earth and some long horizontal growing roots with adventitious buds. The adventitious buds are also on the base of central and lateral roots. The stem short, ramping on earth, covered with destroyed petiole bases, branched. Basal leaves longpetioled,

3-parted, the divisions cleft into broad and irregular lobes 0,5—1,5 cm. broad. (figs. 10 a. plate IV).

The mean petiole length is 21,5 cm. and the mean length of central blade division — 8 cm. The leaves are thick, stiff, dark

Fig. 11. Epidermis from the lower side of the leaf of *A. Janczewskii* ($\times 520$).

green and sparsingly haired with long hairs. The stomata are on both sides of the leaves and are greater and in minor number on 1 sq. mm. of the leaf surface than by *A. silvestris* and *A. multifida* as is shown in tables 10 and 11 and on the fig. 11.

TABLE 10. Size of the stomata of *A. Janczewskii* in μ .

Name of plant	42	45	48	51	54	57	60	63	66	6+	72	75	78	81	84	87	n	M	m
<i>A. Jancz.</i> length								1	2	5	21	33	23	7	5	3	100	73,7	0,14
Warsaw bread.			5	10	42	33	5	3	1	1							100	55,9	0,11
<i>A. Jancz.</i> length									1	10	16	30	18	13	8	4	100	73,9	0,15
Edinb. bread.	2	6	10	15	17	34	9	3	3	1							100	57,6	0,17

TABLE 11. Number of stomata and epidermal cells on 1 sq mm.

Name of plant	Side of leaf	Number of stomata	Number of epid. cels	Stomatal index
<i>A. Janczewskii</i> Warsaw	upper	12,0	185,0	6,0%
	lower	28,6	171,5	14,4%
<i>A. Janczewskii</i> Edinburgh	upper	13,1	193,0	6,3%
	lower	30,0	177,3	14,8%

In the tables 10 and 11 and in the following pages I denote the amphidiploids obtained by me in F_2 of the cross *A. silvestris* x *A. multifida* v. *magellanica* as *Anemone Janczewski* Warsaw, other hexaploids are denoted also as *Anemone Janczewskii* but with the name of Botanical Garden from which they were obtained. We see from these tables that the size and number of stomata of *A. Janczewskii* Warsaw and Edinburgh are very similar. The floral shoots are 35 to 60 cm. high, usually branched with 2 to 4 flowers. The shoots are thick with long erect hairs. The involucral leaves are shortpetioled, otherwise similar to the basal-ones (fig. 10). The lateral peduncles are also involucre with two leaves. The number of flowers on the shoot is as follows:

The number of flowers	1	2	3	4	5	n	M	m
<i>A. Janczewskii</i> Warsaw	1	6	10	1	—	18	2,7	0,20
<i>A. Janczewskii</i> Edinburgh	7	32	38	8	3	88	2,7	0,10

The floral buds are conic and on erect peduncles. The flowers are large to 5 cm in diameter, largely opened when blooming. The sepals 5 to 9 oblong or ovoid. The numbers of sepals in flower are as follows:

The number of sepals	5	6	7	8	9	n	R	m
<i>A. Janczewskii</i> Warsaw	8	17	11	6	2	44	6,48	0,16
<i>A. Janczewskii</i> Edinburgh	35	79	62	18	4	198	6,38	0,06

The mean values are here quite identical. The length and breadth of the sepals are represented on table 12.

The sepal dimensions of *Anemone Janczewskii* of different sources are statistically identical. The colour of the sepals is sulphur-yellow 1/3 according to Horticultural Colour Cart that is one shade lighter in comparison with *A. multifida* v. *magellanica* and its hybrid with *A. silvestris*. The sepals are stiff and breaking by bending which never occurs by parental species. On the outer side the sepals are silky hairy. The stamina are very numerous with great gold-yellow anthers. The good pollen grains are much greater than by *A. multifida* and with twelve furrows. The amount of good pollen grains varies from 42,8% to 78,9% by various plants of *Anemone Janczewskii* Warsaw. By *Anemone Janczewskii* from Edinburgh and other Botanical Gardens similar conditions exist. The question of pollen and seed fertility of the amphidiploids is discussed in a more detailed way after the description of meiotic irregularities in amphidiploids and F_1 plants

The course of meiosis is in direct relationship with the partial fertility of amphidiploids and these both questions must be discussed together.

The diameters of pollen grains of the amphidiploids are put together in the table 13.

It may be seen from table 13 and fig. 8 that diameters of the good pollen grains of F_1 hybrid and of *A. Janczewskii* are of nearly the same range. These results can be interpreted in assuming that the good pollen grains of triploid F_1 hybrids possess unreduced chromosome number. The achenes of the amphidiploids are of the same shape as by F_1 hybrids but are still greater. The weight of the achenes of *Anemone Janczewskii* greatly exceeds the weight of the achenes of parental species. 100 achenes of *A. Janczewskii* Warsaw weigh $2,55 \pm 0,05$ mg., *A. Janczewskii* Edinburgh — $2,38 \pm 0,05$ mg., of *A. silvestris* — $0,70 \pm 0,05$ mg. and of *A. multifida* v. *magellanica* — $1,79 \pm 0,05$ mg. The dimensions of the achenes of *A. Janczewskii* Warsaw are given below:

length	of the achenes	M=3,64 mm	m=0,02 mm	n=115
breadth	„ „ „	M=1,92 mm	m=0,01 mm	n=115
thickness	„ „ „	M=0,51 mm	m=0,01 mm	n=115

The seed fertility like the pollen fertility is also only partial. The percentage of fertile seeds varies by different plants from 22,5 to 74,8 by *Anemone Janczewskii* Warsaw. The fertility of other amphidiploids is represented in part VI. The germinating power of good seeds reach to 85% and the seeds germinate quickly in at last four weeks. In spite of the only partial fertility of these plants they seem to possess a not minor capacity of reproduction when compared to parental species, because they produce great number of flowers on one plant and a greater number of ovulae in one flower. Now, it may also be remembered, that beside normal achenes with full developed embryos, achenes also occur with embryos the development of which was stopped (Fig. 12).

Reasuming our observations on hexaploid *Anemone Janczewskii* it must be stated that it differs distinctly from the parental species in a great number of morphological traits. It is constant as have shown my observations on the second generation. Only the seed and pollen fertility show a segregation in more and less fertile types in different generations. Beside a more sterile forms they appear also highly fertile ones. These latter-ones permit to hope that by proper selection it would be possible to obtain a fully fertile and constant forms like any other natural species. In nature *Anemone Janczewskii*

most probably don't exist, as the two parental species in any part of their distributional area grow together and then the crossing between them is impossible. *A. Janczewskii* present however an excellent example of the origin of new species in culture. This species has arisen most probably in some of Botanical Gardens, where both parental species are cultivated side by side, on the way of the doubling of the chromosome number in a sterile triploid species hybrid. Then it was propagated by seeds. In the same family of *Ranunculaceae* we have an other example of analogical process in the genus *Delphinium*. The very commonly cultivated for high ornamental values, *Delphinium belladonna*, is a hexaploid with $2n=48$. It is also, like

Fig. 12. Different kinds of achenes development in *A. Janczewskii*, Warsaw ($\times 5$).

Anemone Janczewskii, not known from natural conditions. Lawrence (1936) has found in the material of Ruys a triploid hybrid species with $2n=24$ very near morphologically to *D. Belladonna* but quite sterile. This hybrid has produced some seeds probably as a result of somatic doubling of the chromosome number in a part of inflorescence. From these seeds there were obtained some hexaploid plants with $2n=48$ quite indetical morphologically and in their fertility with *Delphinium belladonna*. Propach (1939) has suggested that the parental species of *Delphinium belladonna* are either of following diploid species with $2n=16$: *D. grandiflorum*, *D. tatsiense* or *D. cheilanthum* and the tetraploid species with $2n=32$ — *D. elatum*. Now, on the other hand, it is known from many examples that analogical speciesforming processes occur in natural conditions too.

V. The comparison of the morphological characters of amphidiploids with parental species and F_1 hybrids.

In the recent years there have appeared many papers describing the changes in morphological and physiological characters in plants due to euploid chromosome number increase. These changes were described by autopoliploids and also by amphidiploids. Recently

Kostoff (1938a) has devoted to this question a special study and on the base of numerous examples has formulated some general rules governing the changes in plants caused by chromosome number increase or reduction. He has stated that by increasing or reduction of polyploidy degree the breadth / length index and the thickness of the leaves, the size and the weight of the seeds, the size of nuclei and cells and the duration of vegetation period increase or decrease respectively. Other morphological and physiological characters as the height of plant, the size of flowers, the size of chloroplasts behave by chromosome number change in an indefinite way depending on the plant material used. It must be already stated, that Kostoff's rules are not for all polyploids absolutely true. Kostoff has formulated his rules chiefly on the base of his experiments in *Nicotiana*, *Petunia* and *Triticum*. The processes described by him are founded on the premises that by the change of chromosome number it occurs a proportionate change in nuclear and cell size. This is true for majority of plants but not for all. The plasmo-nuclear relation and nucleus bulk cell size relation are based on the fact that with the increasing volumen of nucleus the mass of protoplasm and the cell size increase proportionally. The volumen of nucleus depends in first line on the amount of chromatine substance. By a constant size of chromosomes by different polyploidy degree, the nucleus bulk is proportionate to the chromosome number. Only in this case Kostoff's rules hold. Fagerlind (1937) has described many autopolyploid forms in different species of *Galium* in which the chromosomes dimensions decrease with the increase of polyploidy degree. Fagerlind has demonstrated on the base of litterature that such a behaviour of chromosomes is not so rare in plants. By *Galium* only one species, namely *G. laevigatum*, has constant chromosome dimensions by different autopolyploid forms and only in this species the autopolyploids present very distinct „gigas” characters. This is an example that in the plasmo-nuclear relation the deciding factor is the nuclear bulk determined merely by the chromatine mass and not by the chromosome number only.

Recently Wettstein (1938) has described that by autopolyploidal form of *Bryum* the cell dimensions are immediately after the doubling of the chromosome number much greater when compared to the cell dimensions of the haplont. The cell dimensions however diminish from one generation to the next, so that after eleven generations they reach the initial dimensions that they had before the doubling of the chromosome number. After the doubling of the

chromosome number the polyploid form is quite sterile but after eleven generations the fertility is restored. Perhaps similar processes are more common in flowering plants and cause that the polyploids known from nature don't present „gigas” characters and are fertile in opposition to the most of experimental polyploids.

From these grounds it follows that the polyploids can be divided in two classes: these which immediately after the doubling of the chromosome number have the same chromosome dimensions and those by which the chromosomes immediately after doubling are smaller. The Kostoff's rules concern only the first class of polyploids.

The physiological properties seem to be in a less direct relation to the polyploidy degree. Only the rate of cell divisions and osmotic value of the cells diminish with the increasing polyploidy degree in a rather regular way in the great majority of studied cases (Becker 1932 Schlösser 1936).

Anemones belong to the first class of polyploids, as the chromosome dimensions are constant after chromosome doubling. The size of nuclei and cells increase with the increasing number of chromosomes but not in a strictly proportionate way. The behaviour of organ dimensions is more variable as they depend not only on the cell size but also on the cell number and genetic factors. At first it must be stated that the effect of the doubling of the chromosome number can be studied only by comparison of triploid F_1 hybrid with its hexaploid progeny. The diploid *A. silvestris* and tetraploid *A. multifida* are two different species which differ not only in their chromosome number but also in their genetical constitution. To demonstrate the relations between the chromosome numbers and the expressions of different plant characters I have tabulated together the mean values of cell and organs dimensions for all anemones which were used in my experiments.

A. The size of nuclei and chloroplasts

The nuclei were measured only in the pollen grains. The measurements were made in aceto-carmin preparations only from grains which were undamaged and well stained. The results are as follows (in table 14).

The volumen of nuclei was calculated using the formula $V=4/3 ab^2$. We see that nuclear volumen by diploid chromosome number is three times greater, and by triploid number — five times greater than the volumen of nuclei with haploid chromosome number.

From this table it is evident that the good pollen grains of triploid hybrid have unreduced chromosome number as the nuclei are of similar size as by hexaploids. The nuclei in P. M. C. and in meristematic tissues of root tips show similar relations but were not measured.

TABLE 14. The dimensions of generative nucleus from pollen grains.

Name of plant	n	Length of long. axis in μ a	Length of short axis in μ b	Volumen in μ^3	Linear relat. of long axis	Cubic relat.
<i>A. silvestris</i>	8	6,7 $\pm$ 0,1	3,5 $\pm$ 0,1	342	1	1
<i>F₁ silv. x mult.</i>	12	11,5 $\pm$ 0,2	6,2 $\pm$ 0,1	1802	1,7	5,2
<i>A. multifida</i>	16	9,9 $\pm$ 0,1	5,1 $\pm$ 0,1	1055	1,47	3,0
<i>A. Janczewskii</i>	24	11,3 $\pm$ 0,2	6,1 $\pm$ 0,1	1717	1,6	5,0

The size of chloroplasts in my *Anemone* material was the same in all plants, regardless the chromosome number, and varies from 3 to 4,5 μ . Their number per cell seems to increase with the increasing cell dimensions by polyploid forms.

From the literature data it seems that there is no rule as to the chloroplasts behaviour in polyploids. Sax (1937) has stated by *Tradescantia* and *Secale* the increase of chloroplast size by the doubling of the chromosome number, but Kostoff and Kendall (1937) by tomato have found their size to be constant and the first author (Kostoff 1938a) has even reported that by some plants the chloroplasts may diminish with the increasing polyploidy degree.

B. The cell size

The size of the cells increases together with the increasing chromosome number. This seems to be true for all kinds of cells but I have measured only P. M. C. and pollen grains. The increase of cell dimensions is very distinct by meristematic cells of root tips but unfortunately they were not measured. The P. M. C. were measured in the study of tetrads when their shape is the most regular and circular in outline. The results are shown in table 15.

TABLE 15. The dimensions of P. M. C.

Name of the plant	2n	Diameter in μ	Linear relation
<i>A. silvestris</i>	16	13,9 $\pm$ 0,2	1
<i>F₁ silv. x mult.</i>	24	18,3 $\pm$ 0,2	1,31
<i>A. multifida</i>	32	22,2 $\pm$ 0,2	1,59
<i>A. Janczewskii</i>	48	25,3 $\pm$ 0,2	1,81

The results of pollen grains measurements are to be shown in table 16.

TABLE 16. The pollen grains dimensions.

Name of the plant	2n	Diameter in μ	Volumen in μ^3	Linear relation	Cubic relation
<i>A. silvestris</i>	16	$19,0 \pm 0,04$	3951	1	1
<i>F₁ silv. x mult.</i>	24	$33,8 \pm 0,36$	20218	1,7	5,6
<i>A. multifida</i>	32	$28,8 \pm 0,12$	12507	1,5	3,4
<i>A. Janczewskii</i>	48	$32,3 \pm 0,27$	17644	1,7	4,9

The comparison of the tables 14 and 16 shows a close parallelity in the linear and cubic relations of nuclei and cell dimensions in pollen grains. Unfortunately they did not show the real dimensions of the triploid condition as here the pollen possess unreduced chromosome number. The coefficients of magnification in linear dimensions for nuclei, P. M. C. and pollen grains are for $4n/2n=1,5$.

C. The size of the organs

1. The stomata. The stomatal apparatus is composed of small and fixed cell number and shows a distinct direct relation between size and the chromosome number. The stomata of *A. silvestris* and *A. multifida* differ however not only in their size but also in their shape. The dimensions of stomata seem to be not only the function of cell size but also of genetic constitution of the plant. In fig. 13 we see the stomata of all investigated forms and in the table 17 are given the average data of their measurements.

TABLE 17. The dimensions of stomata.

Name of the plant	2n	Length in μ	Breadth in μ	Length: breadth ratio
<i>A. silvestris</i>	16	$38,2 \pm 0,25$	$36,2 \pm 0,25$	1,11
<i>F₁ silv. x mult.</i>	24	$43,2 \pm 0,30$	$31,4 \pm 0,25$	1,37
<i>A. multifida</i>	32	$53,2 \pm 0,30$	$37,4 \pm 0,20$	1,41
<i>A. Janczewskii</i> Warsaw	48	$73,7 \pm 0,14$	$55,9 \pm 0,11$	1,31

These data show that the length/breadth index for triploid is intermediate between the parents. The shape of stomata by hexaploid is very similar to shape of the stomata by triploid, which indicates that the magnification of stomatal size is due only to the greater size of the constituent cells. In this case we have only a pure numerical influence of chromosome number. The relations in the distribu-

tion of the stomata on the leaf surface are very interesting. Taken as example the lower side of the leaves. The stomatal index is for:

<i>A. silvestris</i> $2n=16$	23,8%
F_1 <i>silv. x mult.</i> $2n=24$	19,1%
<i>A. multifida</i> $2n=32$	16,6%
<i>A. Janczewskii</i> $2n=48$	19,0%

From these data we see that stomatal index differs distinctly in the two crossed species. By the hybrid it is intermediate. By the

Fig. 13. The comparison of stomatal dimensions of *A. silvestris* ($n=8$) (upper right), F_1 *A. silvestris* $\times$ *A. multifida* v. *magellanica* ($n=12$) (upper left), *A. multifida* v. *magellanica* ($n=16$) (bottom left) and *A. Janczewskii*, Warsaw ($n=24$) (bottom right).

amphidiploid, however, it is identical with the stomatal index of triploid hybrid in spite of the greater stomatal size and smaller number of stomata on surface unity. This indicates that by the doubling of the chromosome number the distribution of stomata rests without any change and the stomata are enlarged in the same degree as the epidermal cells. The stomatal distribution is a character genetically based.

2. The sepals. It is no correlation between the size of the sepals and the chromosome number. The tetraploid species have smaller sepals than the diploid. But the tetraploid has more sepals than

the diploid species. Also the shape of sepals is different by the two investigated species. The mean values of sepals dimensions and their length/breadth index are given below in table 18.

TABLE 18. Sepals dimensions.

Name of the plant	2n	Sepals length in mm	Sepals breadth in mm	Length breadth index
<i>A. silvestris</i>	16	28,2±0,15	19,8±0,16	1,42
F ₁ <i>silv.</i> x <i>mult</i>	24	21,4±0,12	11,3±0,13	1,89
<i>A. multifida</i>	32	15,9±0,13	7,8±0,08	2,03
<i>A. Janczewskii</i>	48	24,5±0,17	14,7±0,14	1,66

From this table it is to be seen that size and shape of the triploid hybrids sepals are intermediate between the parental species and nearer to the tetraploid parent. By the hexaploid the length/breadth index is near to the index of the triploid hybrid. The rate of sepals length enlargement in the case $6n/3n$ is 1,14 and for sepals breadth — 1,40. Both these values are smaller than the corresponding values of stomatal enlargement. On the visual impression the cells of sepals epidermis are just as enlarged in comparison to the triploid as the stomata and epidermal cells of the leaves. The minor rate of sepals enlargement seems to be caused by slower rate of cell divisions by the amphidiploid. Also the changes in the length/breadth index in leaves and sepals may be explained as due to changes in cell divisions rates. In the case of stomata, with small and constant cell number, this process is evidently without influence. But by the organs, composed of a great and undetermined cell number their size after chromosome number increase is chiefly influenced by two processes acting in an opposite way and namely increase of cell size and decrease of cell number. From the literature it is known that in several instances the plants after the doubling of the chromosome number are smaller in general size and in some organs, in spite of greater cell size as compared with the initial form. This is evidently due to the suppression of greater cell size effect by the effect of decreased cell number.

3. The leaves. The leaves of *A. silvestris* and *A. multifida* are so different that an exact biometric comparison is impossible. The leaves of hexaploids as compared with the leaves of triploids are similar but thicker and greater. They seem to be somewhat more enlarged in breadth than in length which would be in good conformity with Kostoff's general rule. As their shape is too irregular to be measured I have no quantitative data. The greater increase of the

breadth of the leaves in respect to their length by amphidiploid when compared to his triploid ancestor can be judged from the fig. 10.

4. The achenes. The achenes of *A. silvestris* differ greatly both in their size and shape from the achenes of *A. multifida*. The measurements of achenes dimensions and their weight are given in table 19.

TABLE 19. The dimensions and weight of the achenes.

Name of the plant	2n	Length in μ	Breadth in μ	Thickness in μ	Weight of 100 achenes in mg.
<i>A. silvestris</i>	16	$2,32 \pm 0,02$	$0,75 \pm 0,015$	$0,26 \pm 0,007$	$0,70 \pm 0,05$
F_1 <i>silv.</i> x <i>mult.</i>	24	$3,24 \pm 0,04$	$1,64 \pm 0,025$	$0,50 \pm 0,013$	
<i>A. multifida</i>	32	$2,70 \pm 0,02$	$1,57 \pm 0,018$	$0,49 \pm 0,013$	$1,79 \pm 0,05$
<i>A. Janczewskii</i>	48	$3,64 \pm 0,02$	$1,92 \pm 0,01$	$0,51 \pm 0,01$	$2,38 \pm 0,05$

The shape of the achenes of the triploid is intermediate between the parental species, but their dimensions are greater. The achenes of hexaploids are of similar shape to those of triploids but are still somewhat greater. The weight of the amphidiploid seeds exceeds greatly the weight of the seeds of both parental species according to the Kostoff's general rule.

VI. The cytology of the investigated plants

1. *Anemone silvestris* L.

A. The somatic complement. Complete somatic metaphasal plate from root tip is shown in fig. 14a. In each plate there are 16 chromosomes. The chromosomes can be divided, according to their point of spindle attachment, the length of both arms and the presence or absence of trabants in eight types. The different types of chromosomes are marked here with capital letters in alphabetical order, A-denoting the longest type and H-the shortest. Each type is represented twice. The type A has median constriction and the arms ca. $4,5\mu$ long, the type B has also median constriction and the arms ca. $3,5\mu$ long, type C has submedian constriction with the arms ca. $3,5\mu$ and 2μ long, types D and E have median constriction and the arms ca. 3μ long, types F and G have subterminal constriction and a total length of ca $4,5\mu$ and the type H is similar to the precedings but has a trabant on the shorter arm. A complete chromosome morphological analysis was possible in this species from several plates as the chromosomes

lay very beautifully scattered and the constrictions were sharply defined.

B. Meiosis. In preparations from one floral bud fixed in a suitable time it is possible to have in P. M. C. from different anthers all stages from prophase to tetrads and young pollen. The course of meiosis is here quite normal. The early prophase stages have large nuclei with a dense mass of chromatin threads and one or two nucleoli. In leptotene the chromosomes are visible as long strings

Fig. 14. *Anemone silvestris*: a-somatic metaphase, b-diplotene, c-late diakinesis, d, e, f-metaphase I, g-anaphase, h-telophase, i-anaphase II, j-tetrad (a-i $\times 3000$, j $\times 1700$).

of beads strung together. In pachytene the threads are evidently double but they are so crowded in the nucleus that an attempt to analyse the full complement proved a failure. Only in diplotene it was possible to differentiate all 8 bivalentst. The bivalents are much shortened and the chiasmata are visible and can be counted in some propitious P. M. C. (fig. 14b). I always found in this stage and in the latter-ones only 8 bivalents. The number of chiasmata per bivalent is in this stage relatively easy to establish, as between the points of chiasmata large loops are formed. The number of chiasmata per bivalent seems to be proportionate to the chromosome length.

The number of chiasmata per bivalent varies from 2 to 6, the mean value being 3 chiasmata per bivalent (see table 20).

TABLE 20. The number of chiasmata and terminalization coefficient in *A. silvestris*.

Stage	No. of bivalents	Total no. of XX	No. of term. XX	Total no. of XX per bivalent	No. of term. XX per bivalent
Diplotene	20	61	21	3,05	1,05
Diakinesis	27	64	16	2,37	0,58
Metaphase I	50	63	8	1,26	0,16

In diplotene the terminalization of chiasmata is great and almost every bivalent has one terminal chiasma. In late diakinesis (fig. 14c) the chromosomes are deeply condensed and are evenly disposed through the nucleus. From the table 20 it is to be seen that the number of chiasmata per bivalent has decreased and the terminal chiasmata are also less numerous. This stage is the best for studying the number of bivalents, when the fixation is good. In the 31 P. M. C. analysed carefully at diakinesis I always found only 8 bivalents.

At metaphase I the bivalents are in form of crosses with arms of different length or in form of rings with two loose ends (figs. 14d, e, f). The great majority of bivalents possesses one interstitial chiasma, two or three chiasmata are very rare (table 20). The process of the reduction of terminal chiasmata together with the general decrease of chiasmata number from prophase to metaphase I is a rather exceptional phenomenon by plants. In the majority of plants it was described by several authors an increase of terminalization coefficient by the decrease of total chiasmata number. This peculiar mode of chiasmata behaviour was firstly described by Moffet (1932) for several species of *Anemone* and for *Anemone silvestris* too. His table is quite identical with the table 20 of the present paper. Similar conditions in chiasmata behaviour seem to be rather common in the whole family of *Ranunculaceae* and were described for *Pulsatilla* by Rosenthal (1936) and for *Ranunculus* by Larter (1932). This peculiar mode of chiasmata evolution Moffet ascribed to the lack of terminal affinity.

The anaphase and telophase I have a quite normal course (figs. 14g, h). The bivalents separate normally and quite simultaneously and I have never found any chromosome or fragment lying on equator or lost in cytoplasm through failure to reach the poles. In the interkinesis the dyad nuclei have a quite normal appearance of resting nuclei with one nucleolus.

The second division, like the first, is quite normal. In anaphase II I have stated in 12 cases the presence of four chromosome groups each of 8 chromosomes (fig. 14i). The axis of two second divisions are usually parallel seldom perpendicular. The cytokinesis is performed according to so called „Furchungstyp”.

Fig. 15. *Anemone multifida* v. *magellanica*: a-somatic metaphase, b, c, d-metaphase I, e, f, g-anaphase I, h-telophase I, i-telophase II, j-tetrad. (a-g $\times 3000$, h-j $\times 1700$).

The tetrads are quite normal with four microspores, each with one nucleus of similar size (fig. 14j). The young pollen has a quite good appearance and in size is quite uniform.

2. *Anemone multifida* Poir.

A. The somatic complement. From this species I have studied chiefly its variety *magellanica*. The varieties *hudsoniana* and *alba* from different sources were also cytologically examined and I have never found any noticeable difference in somatic complement or meiotic course among them. Thus holds the following description for all varieties of *A. multifida* which don't differ in their cytology.

In the metaphasal plates from root-tips it is always 32 chromosomes (fig. 15a). A detailed morphological analysis is here difficult owing to the high number of chromosomes which are usually bent together. Nevertheless it seems that there exist here 4 groups of 8 chromosomes very similar or even identical in their external morphology with the 8 types of chromosomes of *A. silvestris*. The chromosomes of the type H with subterminal constriction and a trabant, which are easy to distinguish are quite the same as by *A. silvestris* and in several plates they are represented four times in other two or three times but never more than four times. The same holds also for other types of chromosomes but it was very difficult to establish all types of chromosomes in one plate.

B. Meiosis. Observations at meiosis showed no abnormal behaviour. The prophasal stages up to diplotene are too obscure to be analysed. In diakinesis in 25 analysed P. M. C. I have always stated the presence of 16 bivalents. Other configurations were never found. In metaphase I in 52 P. M. C. it was 16 bivalents (figs. 15b, d) and only in one cell I have found 15 bivalents and 2 univalents (Fig. 15e). The presence of these univalents was probably due to a premature separation of one of the bivalents. The bivalents are of the same form as by *A. silvestris*, the most common are the bivalents with one interstitial chiasma forming a cross with unequal arms. The mean value of chiasmata per bivalent is at metaphase 1,16 and the terminalization coefficient is 0,10 both these values are very near to the corresponding values for *A. silvestris*. The anaphasal separation of bivalents is normal. In 12 P. M. C. analysed in this stage I have counted 16 chromosomes in each of anaphasal chromosome group (fig. 15e, f, g). The heterotypic division does not present any abnormality. In 189 P. M. C. in the stage of telophase II there were always only 4 nuclei of equal size (fig. 15i). Also the tetrads and the young pollen are quite normal (fig. 15j).

3. *F₁* *Anemone silvestris* x *A. multifida* v. *magellanica*

A. The somatic complement. In general, except some minor details, the course of meiosis in hybrids between *A. silvestris* and different varieties of *A. multifida* is the same. Also their somatic complements have the same chromosome number. In numerous metaphasal plates from root-tips it was found 24 chromosomes (fig. 16a). This number was expected from parental chromosome numbers and was also found by Moffet (1932). The 8 types of chromosomes are here represented 3-times. On some preparations it is easy to distinguish

the three longest chromosomes of the type A and the three shortest chromosomes with the trabant of the type H.

B. Meiosis. Meiosis runs on the whole a quite abnormal course. The early prophase stages are similar as by parents and were not studied. In diplotene it is already visible that, beside double strands, single strands also occur (fig. 16b). In this stage I was never able to

Fig. 16. F_1 *A. silvestris* $\times$ *A. multifida* v. *magellanica*: a-somatic metaphase, b-diplotene, c-diakinesis with 4_{II} and 16_I , d-diakinesis with 7_{II} and 10_I , e-diakinesis with $5_{II}+14_I$, f-metaphase I with 3_{II} and 18_I , g-metaphase I with 7_{II} and 10_I , h-metaphase I with 4_{II} and 16_I , i-side view of metaphase I with 6_{II} and 12_I , j, k-me, taphase I with 5_{II} and 14_I , l-metaphase I with 7_{II} and 10_I . ($\times 3000$).

establish the precise number of double and single strands in the whole nucleus, the strands are contracted around the nucleolus and it is not possible to analyse the whole nucleus. In the single isolated bivalents it is however possible to count the number of chiasmata. The number of chiasmata range from one to four in one bivalent. The mean value is by the hybrid lower than by *A. silvestris* (table 21), but this difference is not absolutely sure, as the chiasmata number decreases rapidly during the diplotene and it is very difficult to found two preparations in a strictly comparable stage.

TABLE 21. Number of chiasmata and terminalization coefficient by F_1
A. silvestris x *A. multifida* v. *magellanica*.

Stage	No. of bival.	Total no. of XX	No. of term. XX	Total no. of XX per bival.	No. of term. XX per bival.
Diplotene	15	40	12	2,66	0,80
Diakinesis	51	91	17	1,78	0,33
Metaphase I	37	41	6	1,10	0,16

At diakinesis it is to be seen from 21 to 16 chromosomal bodies. In this stage it is easy to distinguish the uni- and bivalents. The bivalents are usually greater than the univalents and form crosses or rings. The univalents are rod- or V-shaped. Judging from the dimensions of bivalents, it is possible that most of the bivalents are formed between the longest chromosomes of the types A, B, C, D and E and that the short types of chromosomes rest mostly unpaired. This is probably connected with the fact observed by *A. silvestris* that the number of chiasmata is proportionate to the length of the pairing chromosomes. Thus the shortest chromosomes have less chance to form one chiasma and rest unpaired. In this stage the mean value of chiasmata per bivalent is much lower than by *A. silvestris*. This difference seems to be real, especially when it is remembered that by the hybrid the bivalents are formed chiefly between the longest chromosomes with the highest chiasmata number. The lower chiasmata number by the hybrid is probably in connection with the only partial homology of the pairing chromosomes. The trivalents were never observed, notwithstanding I have made special researches to find them. The configurations of chromosomes found in 20 P. M. C. analysed at diakinesis are tabulated in table 22.

TABLE 22. Configurations of chromosomes at diakinesis and metaphase by F_1
A. silvestris x *A. multifida* v. *magellanica*.

Configuration	$3_{II} + 18_I$	$4_{II} + 16_I$	$5_{II} + 14_I$	$6_{II} + 12_I$	$7_{II} + 10_I$	$8_{II} + 8_I$	Mean no. bival. p. cell.
Diakinesis No. of P.M.C.	1	9	3	3	2	2	5,1
Metaphase I No. of P.M.C.	7	17	20	20	7	—	5,04

It is evident from this table that at diakinesis it is formed from 3 to 8 bivalents in 1 P. M. C., the mean value is 5,1 which by the 12 bivalents theoretically possible form 42,5% of the possible

conjugation. The figs. 16c, d and e illustrate the different types of configurations observed at diakinesis in this hybrid.

At the metaphase I the bivalents move towards the equator and form a regular plate. The univalents usually lie at random on the spindle. In late metaphase the majority of univalents are on the edges of the metaphasal plate. Sometimes all univalents lie on the near proximity of the equatorial plate. The bivalents are in form of crosses or rings with free ends as by parental species. The number of chiasmata in one bivalent is 1 or 2, the mean value is 1,10 and the terminalization coefficient — 0,16 (table 21). The number of bivalents in different P. M. C. range from 3 to 7. The actual numbers of P. M. C. with different chromosome configurations are tabulated in table 22. The mean value of bivalents per 1 P. M. C. is 5,04 quite the same as at diakinesis. This indicates, that notwithstanding the great reduction of chiasmata frequency from diakinesis to metaphase, no separation of bivalents occurs at this period. The figs. 16f, g and h illustrate the different metaphasal plates from polar-view. The figs. 16i, j k and l illustrate side-views of metaphasal plates. In fig. 16i we have a configuration of $6_{II}+12_I$ and the all univalents lie on the periphery of the equatorial plate, in figs. 16j, k and m is a configuration of $5_{II}+14_I$ and the univalents are scattered on the whole spindle, in the fig. 16f we see the configuration of $7_{II}+10_I$.

When the anaphase begins the bivalents begin to separate and the univalents follow different courses according to their position in the cell. Those lying far away from the equator are included unsplitted to the daughter anaphasal groups. The majority of univalents which lie nearer the equator move in the late anaphase after the separation of bivalents on to the equator and divide after a short interval. The two halves follow the daughter bivalents to the poles and are included to the daughter nuclei. In 18 P. M. C. at late anaphase I have counted following numbers of dividing univalents.

No. of dividing unival.:	3	5	6	7	8	9	10	11	12	13	14
No. of P. M. C.	1	2	—	3	6	1	—	2	—		1

In average there is at anaphase 8,3 dividing univalents per 1 P. M. C. when at metaphase there are in average 13,9 univalents per 1 P. M. C. This seems to indicate that in average 5 univalents per 1 P. M. C. divide in early anaphase together with the bivalents or even, which is more probable, they are included univided to daughter chromosome groups. On the figures 17a and b we have a typical view of the anaphase I by the hybrid. We see two anaphasal

chromosome groups and on the spindle the splitted univalents. The process of univalents splitting is very slow and its beginning is very delayed in comparison with the bivalents separation. It is not rare that the univalents have not finished their splitting when in the daughter nuclei telophasal changes are to be seen. In the figures 17c, d

Fig. 17. F_1 *A. silvestris* $\times$ *A. multiifida* v. *magellanica*: a-anaphase I with univalents on the spindle, b, c, d, e-late anaphase I with dividing univalents, f-telophase I with univalent halves on the spindle, g, h, i-interkinesis with chromosomes lost in the cytoplasm and small supernumerary nucleus. ($\times 3000$).

and e we see dividing univalents when in the daughter nuclei telophasal changes are visible. On the fig. 17f we see the halves of 4 univalents passing to opposite poles. The delaying of univalents splitting and movements causes that sometimes the lagging chromosomes are not included to the daughter nuclei. In telophase and interkinesis we often see the whole chromosomes or univalents halves

left out in cytoplasm. These chromosomes are either scattered in the cytoplasm, or form small supernumerary nuclei, which undergo the telophasal and interkinetic changes. The fig. 17g illustrates a P. M. C. at interkinesis with two interkinetic nuclei and with two undivided and one half of univalents. On the fig. 17h between the two interkinetic nuclei we see three halves of univalents and on the fig. 17i besides one half of univalent it is also a small supernumerary nucleus. In the 67 P. M. C. analysed in late telophase I have found following numbers of univalents lost in cytoplasm:

No. of lost univalents	0	1-3	more than 3
No. of P. M. C.	15	47	8

At interkinesis the conditions are similar:

No. of lost chrom. or small nuclei	0	1	2	3	more than 3
No. of P. M. C.	15	8	3	3	2

Thus more than 50% of P. M. C. are abnormal with the lost chromosomes or supernumerary nuclei, the other P. M. C. have only two nuclei but often of different size. In spite of very careful searches I never found in anaphase, telophase or interkinesis the formation of restitution nucleus containing all chromosomes in one nucleus. This was to expect as I have stated in this hybrid the formation of diads after second division.

The second division shows also very numerous irregularities. In metaphase II it is often to be seen that beside the metaphasal plates the lost chromosomes are scattered in cytoplasm or even small supernumerary plates with their own spindles are formed. In 7 P. M. C. with only two plates I was able to count exactly the chromosome number in the both plates. The numbers found were as follows: in 3 P. M. C. it was 18 chromosomes in both plates, in 2 P. M. C. there were 17 and 18 chromosomes in the two plates, in 1 P. M. C. it was ca. 19 in both plates and in 1 P. M. C. it was 15 and 16 (fig. 18a). These numbers are a direct proof, that in the metaphasal plates of the second division they are beside the separated chromosomes from the bivalents, also the halves of the splitted univalents. In the first case 12 univalents have splitted; in the second — 11, in the third — 14 and in the fourth 7 univalents. The course of the anaphase II presents also some abnormalities. In numerous P. M. C. between the anaphasal chromosome groups numerous chromosomes lag on the spindle. The most probably there are the halves of the univalents which have already divided at the first division. These halves are in the second division distributed at random to the tetrad nuclei

(fig. 18d). Sometimes the second anaphase has a quite normal appearance, as in figs. 18b and c. In some P. M. C. I have observed between the anaphasal groups long and thin chromatine bridges joining the both anaphasal groups. From 40 P. M. C. at anaphase II only 21 were normal with four chromosome groups, 12 have from 1 to 6 chromosomes lagging on the spindles and 7 have beside lagging chromosomes also one or two bridges. The number of chromosomes in

Fig. 18. F_1 *A. silvestris* $\times$ *A. multifida* v. *magellanica*: a-metaphase II with 16 and 15 chromosomes in each plate, b, c-anaphase II, d-anaphase II with lagging chromosomes, e-anaphase II with 12 and 11 chromosomes in two groups, f, g, n, o-telophase II with chromosomes lost in the cytoplasm and supernumerary nuclei, h, i, j-diads, k-triad, l-tetrad, m-pentad. (a-g- $\times 3000$, h-o- $\times 1700$).

anaphase II groups range from 8 to 14. In the figs. 18d and e are illustrated the second anaphase groups with 11 and 12 chromosomes. The axis of two divisions form usually an obtuse angle but sometimes they are quite parallel. In some cases I have observed that the both spindles and anaphasal groups of the two divisions touch together or are very near and are almost fused. This seems to indicate that in some cases the two anaphasal groups fuse completely and may be the cause of diad formation observed in this hybrid. In telophase II ca

50% of P. M. C. have chromosomes out in cytoplasm or some small „micronuclei”. The figs. 18f, g, h and i illustrate the various views of the second telophase. The tetrads are very irregular. Beside normal ones with 4 microspores, I have found also the „tetrads” with 2, 3, 5, 6, 7 and 8 cells. The microspores have usually one nucleus but in some cases small supernumerary nuclei are to be found. In studying the P. M. C. of the hybrid at the tetrad stage, the following data were obtained:

No. of microspores	2	3	4	5	6	7	8
P. M. C. studied	2	3	59	30	13	2	2
Pollen calculated	4	9	236	150	78	14	16

The number of pollen grains was calculated on the assumption that every microspore gives rise to one pollen grain. Basing on this assumption the percentage of unreduced pollen grains from diads may be 0,8% in this hybrid. The percentage of giant viable pollen grains in this hybrid is about 0,3% as was described above. What is the cause of the discrepancy between the calculated and observed percentage of unreduced pollen grains it is difficult to say. All the pollen originating from other types of tetrads than the diads is most probably unviable. The figs. 18j, k and l illustrate different types of diads and the figs. 18l, m and n represent a triad, tetrad and pentad respectively.

4. F_2 *A. silvestris* $\times$ *A. multifida* v. *magellanica* = *A. Janczewskii*,
Warsaw

I have stated 48 chromosomes in somatic plates in 7 out of 8 plants forming the second generation of this cross and in one a number very near to 48. The course of meiosis in its general traits is similar by all these plants. The difference in fertility and in the percentage of good pollen among the different plants of F_2 are mainly due to some minor and of rather quantitative nature differences in chromosome conjugation by different plants.

The cytological observations have shown no essential qualitative differences in meiotic course among the various plants of F_2 . Thus the description given below is based chiefly on one F_2 plant with the culture number AJW — 7 and it holds to other sister plants too.

A. The somatic complement. In a great number of somatic metaphasal plates I have stated with no doubt 48 chromosomes

(fig. 19a). This is a twice greater number than by the parental plants which have 24 chromosomes in somatic plates. The morphological analysis of chromosomes was not performed, owing to the large number of chromosomes, but there are so conspicuously characteristic that the resemblance with the basal types distinguished by *A. silvestris* cannot escape from discovery. This is true in particular of chromosomes of the A and H types. The A type chromosomes are consi-

Fig. 19. F_2 -amphidiploid AJW-7: a-somatic metaphase, b-diplotene, c-diakinesis, with 24_{II} , d-diakinesis with 21_{II} , 1_{IV} and 2_I , e-diakinesis with 20_{II} , 1_{III} and 5_I , f-metaphase I with 24_{II} , g-metaphase I with 23_{II} and 2_I , h-metaphase I with 22_{II} , 1_{III} and 1_I , i-metaphase I with 19_{II} , 1_{IV} , 1_{III} and 3_I . ($\times 3000$).

derably longer than the others. Chromosomes of this type were found to be represented from 4 to 6 times. Special mention must also be made of the chromosomes designated H, which are just easy to identify as the A type. Besides being the shortest they are also characterized by being satellite-bearing. These chromosomes, too, are present in the number from 4 to 6. The bulk of chromosomes are similar as by triploid parent.

B. Meiosis. The great number of chromosomes and their relatively great dimensions make the analysis of meiosis rather

difficult. Already in the prophasal stages I was impressed by the unusual great dimensions of cells and nuclei. The number of nucleoli varies from 4 to 5. The leptotene and zygotene up to diplotene stages were not analysed owing to the very complicated and dense mass of chromatin threads in which it is very difficult to distinguish chromosomes (fig. 19b).

Fig. 20. F_2 -amphidiploid AJW-7: a-metaphase I with only bivalents, b-metaphase I with two univalents, c-metaphase I with one trivalent and one univalent, d-metaphase I with one quadrivalent, e-metaphase I with 1 tri-, 1 quadri- and three uni-valents, f-metaphase I with 2 quadrivalents, g-metaphase I with 1 hexa- and 2 uni-valents, h-normal anaphase I, i-anaphase I with chromosomal bridge- ($\times 3000$).

At diakinesis, in few completely analysed P. M. C., besides uni- and bivalents usually also tri- and tetravalents are seen distributed in the nuclear area. In 7 P. M. C. at late diakenesis I have stated the chromosome configurations as follows:

24_{II} in 4 P.M.C. (fig. 19c) $20_{II} + 1_{IV} + 6_I$ in 1 P.M.C.
 $21_{II} + 1_{IV} + 2_I$ in 1 P.M.C. (fig. 19d) $20_{II} + 1_{III} + 5_I$ in 1 P.M.C.

The presence of polyvalents in ca. 50% of P. M. C. was a new and striking fact. The polyvalents like the bivalents possess subter-

minal chiasmata and rarely terminal. The univalents were found only in the presence of polyvalents.

In metaphase I in P. M. C. with only bivalents a large metaphasal plate is formed. The form of bivalents and also the number and distribution of chiasmata are similar as by other *Anemone* species. Fig. 19f illustrates such a metaphasal plate from a polar view and the fig. 20a from profile view. When univalents are present they lie outside the plate scattered on the whole spindle, see figs. 19g (polar view) and 20b (side view). The polyvalents lie together with the bivalents on the equator. They are easy to distinguish especially in the profile view of metaphasal plates, as they are much longer in comparison with bivalents and their ends protrude distinctly above the compact mass of bivalents. A complete analysis of metaphasal plates is rather difficult owing to the high chromosome number and also because the majority of the bivalents and polyvalents are crowded and it is generally difficult to determine as certain whether there is a real connection between pairs lying close together or whether they are simply lying in close proximity. In 31 metaphasal plates analysed from polar view I have found following configurations:

Configuration	No. of P. M. C.	Configuration	No. of P. M. C.
24_{II}	20 (fig. 19f)	$22_{II} + 1_{III} + 1_I$	1 (fig. 19h)
$23_{II} + 2_I$	6 (fig. 19g)	$19_{II} + 2_{IV} + 2_I$	1
$22_I + 1_{IV}$	1	$19_{II} + 1_V + 1_{III} + 3_I$	1 (fig. 19i)
$20_{II} + 2_{IV}$	1		

We see that in 64% of P. M. C. normal pairing consisting of 24_{II} occurs, in ca 20% of P. M. C. also univalents occur and in the remaining 16% of P. M. C. the polyvalents were found. Owing to the very low number of examined cells these numbers seem to be not statistically defined. In scope to obtain more accurate data to the relative frequency of the different configurations, I have studied a larger number of metaphasal plates from the side view.

In the side view of metaphasal plates it is much easier to establish the number of univalents which lie on the spindle and also of polyvalents which by their greater dimensions are distinguishable from bivalents. The disadvantage of examination from the side view the plates with the high chromosome number is, that in the great majority of cases, it is impossible to distinguish and then to count all chromosomes which lie densely on the equator. It is then possible that some univalents and even some polyvalents lying on the equator in the dense mass of bivalents can escape from the

observation. By this method it is however possible to analyse a great number of metaphasal plates and to obtain data of statistical significance. I have analysed 196 metaphasal plates from the side view and I have obtained the following results:

No. of uni- and polivalents	No. of P. M. C.	No. of uni- and polivalents	No. of P. M. C.
0 _I , 0 _P	116	3 _I , 1 _{III}	3
2 _I , 0 _P	33	3 _I , 1 _{III} , 1 _{IV}	1 (fig. 20e)
4 _I , 0 _P	8	0 _I , 2 _{IV}	3 (fig. 20f)
6 _I , 0 _P	3	2 _I , 2 _{IV}	1
1 _I , 1 _{III}	14 (fig. 20c)	2 _I , 2 _{III}	2
2 _I , 1 _{IV}	4	2 _I , 1 _{VI}	1 (fig. 20g)
0 _I , 1 _{IV}	7 (fig. 20d)		

The P. M. C. without uni- and polyvalents form 60%, P. M. C. with only univalents but without polyvalents form 22%. These numbers are very near to the numbers established from a polar view. Taking into account the possible omission of some uni- and polyvalents it seems to be sure, that at last one half of P. M. C. possesses only bivalents at metaphase. The number of polyvalents in one P. M. C. varies from 1 to 3. The most frequent are tri- and tetravalents, the hexavalent was only once found in the whole material. The trivalents form usually crosses like the bivalents with one chromosome subterminally attached, the ring of three chromosomes was never found. The tetravalents form chains or rings of four chromosomes usually with subterminal chiasmata. Terminal chiasmata are very rare. The hexavalent forms a chain of six chromosomes with subterminal chiasmata.

The course of anaphase I differs according to the configuration at metaphase. The P. M. C. with only bivalents and perhaps also the P. M. C. with polyvalents but without univalents show quite normal anaphasal separation of the chromosomes (Fig. 20h). The P. M. C. with univalents present abnormal course of the anaphase. The univalents at anaphase lag on the spindle and after a long interval split equationally and the two halves move then to the opposite poles (figs. 21a, b, c, d). In some P. M. C. also bridges were found (fig. 20i). The univalents are very delayed in their splitting and often are not included to the daughter nuclei and form at interkinesis small supernumerary nuclei or lie in cytoplasm (figs. 21h, i). The number of chromosomes in the both anaphasal groups may be equal or not, in the 8 P. M. C. in which both anaphasal groups were analysed

I have found following numbers of chromosomes: in 6 P. M. C. it was 24 chromosomes in each group (fig. 21e), in 1 P. M. C. it was 27 and 26 chromosomes in each group (fig. 21f) and in 1 P. M. C. it was 25 and 24 chromosomes in each group (fig. 21g). The chromosomes in

Fig. 21. F_2 -amphidiploid AJW-7: a, b, c, d-lagging univalents at anaphase I, e-anaphasal groups with 24 chromosomes in each, f-two anaphasal groups with 26 and 27 chromosomes, g-two anaphasal groups with 24 and 25 chromosomes, h, i-interkinesis with supernumerary nuclei. ($\times 3000$).

anaphasal groups were visibly double, but in the cells with unequal chromosomal distribution and with the sum of chromosomes higher than 48 some chromosomes were seen as single strands. These single-strand chromosomes were in most probability the halves of univalents already splitted. In 135 P. M. C. at the stage of interkinesis 120 were normal with two nuclei, 10 P. M. C. possess one small supernumerary

nucleus or some lost chromosomes, 4 P. M. C. have two supernumerary nuclei and 1 P. M. C. has three supernumerary nuclei. The cells with normal 2 nuclei form 88,8% on the total analysed cells at diakinesis, this indicates that not only all cells with only bivalents at metaphase show normal chromosome separation but also about

Fig. 22. F_2 -amphidiploid AJW-7: a-metaphase II with 24 chromosomes in both plates, b-anaphase II with chromosomes lost in the cytoplasm, c-normal anaphase II, d-anaphase II with lagging chromosomes, e-tripolar anaphase II, f-telophase II with chromosomes lost in the cytoplasm g-normal telophase II, h-one of the anaphasal groups with 24 chromosomes, i-tetrad, j-tetrad with supernumerary nuclei, k, l-pentads, m-hexad. (a-h $\times 3000$, i-m $\times 1700$).

30% of P. M. C. with poly-or univalents show no loss of chromosomes at anaphase.

The second meiotic division shows also some irregularities in its course. In the great majority of P. M. C. studied we see at metaphase II two plates with 24 chromosomes in each (fig. 22a). In some P. M. C. beside the two metaphasal plates a third plate with few chromosomes is formed or some chromosomes are scattered in cytoplasm without forming his own spindle mechanism (fig. 22b). In anaphase II either the separation of chromosomes is

normal (fig. 22c) or in some cells some chromosomes are lagging on the spindles (fig. 22d). These are perhaps the halves of the splitted univalents. In cells with supernumerary plates, supernumerary spindles are also found. In one P. M. C. I have found a three-polar spindle uniting the three nuclei together (fig. 22e). In telophase II beside normal cells with four nuclei often P. M. C. occur with supernumerary small nuclei or with chromosomes lost in cytoplasm (fig. 22f). The accurate counting of chromosome number in anaphase II groups is very difficult, the fig. 22h represents one of such groups containing 24 chromosomes. In the stage of tetrads I have analysed 229 cells. They are very heterogenous. Beside normal-ones with four microspores there are also to be found pentads (figs. 22k, l), hexads (fig. 22m) and even heptads and octads. In numerous cells small nuclei forming or not their own cells are to be found. The results of „tetrads” examination are as follows:

No. of microspores	4	5	6	7	8
P. M. C. studied	165	45	15	3	1
Pollen calculated	660	225	90	21	8

The percentage of normal interkinetic P. M. C. was 88,8% and the percentage of normal tetrads is 72% what indicates that some normal appearing interkinetic cells produce after second division abnormal tetrads. The amount of normal pollen calculated on the basis of tetrad appearance would be 66%. The percent of good pollen in this plant was found to be 50,3%. Thus we see that the calculated percentage of good pollen is higher than the percentage of the viable pollen.

By comparison of the triploid and hexaploid plants we see that with the doubling of the chromosome number the course of meiosis in about 60% of P. M. C. becomes normal. In the resting 40 per cent of P. M. C. poly- and univalents were formed. Multivalent chromosome groups obviously result from chiasma formation and crossing-over following auto- and allosyndetic chromosome conjugation in F_1 . The segments of *A. silvestris* and *A. multifida* chromosomes which synapsed in the F_1 hybrids obviously synapse sometimes in the amphidiploid too. The course of meiosis is a reason for expecting the amphidiploids to form a certain percentage of nonviable gametes and not to be fully fertile.

The plants raised from the seeds obtained from Edinburgh were quite identical with the amphidiploids described above not only from the morphological point of view but also in their cytology.

Three generations of these plants were carefully cytologically examined. The great majority of examined plants have 48 chromosomes in somatic plates and some rare plants have somewhat smaller chromosome number, ranging from 42 to 47. The course of meiosis of the hexaploid plants is essentially the same as in amphidiploids described above. Only the percentage of P. M. C. without uni- and polyvalents is somewhat greater. In plants with aneuploid chromosome number 42—47 the course of meiosis resembles in general traits to the meiosis in true amphidiploid but the amount of abnormalities is increased. Space does not allow me to give a detailed description of the cytology of various plants from the Edinburgh family, but cytologically they present any essential new trait in comparison with amphidiploids described above.

VII. The pollen and seed fertility of the studied plants and their crossability

The question of amphidiploids fertility merits special attention as he presents some general theoretical significance. In the light of modern cytogenetic researches it was stated that the experimentally produced amphidiploids present very different degrees of fertility ranging from complete fertility to the nearly complete sterility. The degree of fertility found in a given amphidiploid depends directly on the conditions of chromosome pairing at meiosis. Now, on the other hand the chromosome conjugation in amphidiploid plant is conditioned by the homology between the chromosomes of the parental species, which can be studied in F_1 hybrids. The inheritance of the infertility by amphidiploids is not sufficiently studied although it presents many interesting problems together with the possible role of amphidiploids in nature. In the present part I would accumulate all data bearing on the questions of fertility from my *Anemone*-material.

Before a detailed presentation of accumulated data it seems necessary to discuss a terminological question of the use of the term amphidiploid in this paper instead of the usually employed term „allopolyploid“. In cyto-genetic literature the terms of auto- and allo-polyploids are most frequently used to denominate the intra- and inter-specific polyploids respectively. These terms were created by Kihara and Ono (1926) and determined as follows: „Unter Autopolyploidie versteht man die Verdoppelung desselben Chromosomensatzes, ...unter Allopolyploidie die durch das Zusammenkommen verschiedener Chromosomensätze auf dem Wege der Basterdierung

erfolgte Chromosomenvermehrung". The meaning of the words „desselben and verschiedener" was differently interpreted by different authors and this has caused many misunderstandings in cytogenetic literature. In the year 1936 Lilienfeld, a collaborator of Kihara, has given a more precise explanation of these terms, writing that under „desselben and verschiedener" we may understand homologous and nonhomologous genomes. The degree of genome homology can be measured by the degree of pairing between the chromosomes of two genomes in the hybrid. According to this definition autopolyploids are arisen from plants or hybrids with normal pairing and the allopolyploids from asyndetic hybrids. The auto-and allopolyploids according to Kihara-Lilienfeld definition are by no means identical with the conception of intra-and inter-specific polyploids. It is known that by interspecific hybrids it is to be found all degrees of chromosome pairing from complete lack of chromosome pairing to full pairing of all chromosomes. Allopolyploids sensu stricto would be only these polyploids that have arisen from asyndetic hybrids (naturally only in these cases in which asyndesis is not due to genetic factors but only to the lack of homology). Other interspecific polyploids that have arisen from hybrids with partial or complete chromosome pairing would be at once auto-and allopolyploids together or true autopolyploids. It seems thus to be more appropriate to distinguish between the intra and interspecific polyploids. Only the autopolyploids are synonym to the intraspecific polyploids. To the interspecific polyploids can be applied also the term amphidiploid of Navashin or more generally amphipolyploid. According to Fagerlind (1937) I am inclined to believe that the more rational is the distinction between intraspecific polyploids (or autopolyploids) and interspecific polyploids (or amphidiploids) using as the base the existing specific units of taxonomy except some special cases in which genetic investigations prove the uncorrectness of taxonomic specific differentiation. Such an example furnish Winge (1933) experiments on *Erophila* in which a hybrid between the two ecotypes of *Erophila verna* differing in their chromosome number shows complete asyndesis and the two ecotypes can be regarded as distinct species.

1. Pollen and seed fertility in *Anemone*

Pollen grains were studied in aceto-carmin preparations. Germination tests were unfortunately not performed. The difference between good and bad pollen is sharp and the border-line cases

difficult to classify are relatively rare. It seems that the great majority of pollen grains with a good appearance are really able to germinate.

The seed fertility was tested from the appearance of achenes and also by germination tests. In the table 23 the pollen and seed fertility is recorded in percentage of the total number of tested pollen grains and ovulae respectively.

TABLE 23 Pollen and seed fertility.

Name of the plant	2n	% good pollen	% good seeds	Germination power in %
<i>A. silvestris</i>	16	92,7	96,4	ca. 100
<i>A. multifida</i> v. <i>magellanica</i>	32	97,2	93,3	ca. 100
<i>A. multifida</i> v. <i>hudsoniana</i>	32	93,8	91,6	ca. 90
<i>A. multifida</i> v. <i>alba</i> (Cluj)	32	95,3	93,2	ca. 100
F ₁ <i>A. silv.</i> x <i>A. mult.</i> v. <i>magell.</i>	24	0,26	0,6	ca. 40
F ₁ <i>A. silv.</i> x <i>A. mult.</i> v. <i>hudson.</i>	24	0,12	0,32	0
F ₁ <i>A. mult.</i> v. <i>alba</i> x <i>A. silv.</i> 1	24	0,20	1,09	ca. 45
	2	24	0,24	
	3	24	0,22	
	4	24	0,18	
	5	24	0,21	
	6	24	0,31	
F ₂ <i>A. silv.</i> x <i>A. mult.</i> v. <i>magell.</i>				
AJW 1	48	71,2	70,4	ca. 85
AJW 2	48	51,1	41,4	
AJW 3	48	57,5	31,7	
AJW 4	ca.48	52,2	22,5	
AJW 5	48	78,9	74,8	
AJW 6	48	59,5	42,7	
AJW 7	48	50,3	43,0	
AJW 8	48	42,8	35,2	

The percentage of good pollen and seeds is not absolutely stable for a given plant but vary according to preparation or flower from which the test was made. The greatest variation in pollen fertility was observed by amphidiploids ranging to 10% of difference between the two tests from the same plant. This is evidently caused by the great variation in pairing behaviour of the chromosomes observed even in the different parts of the same flower by the amphidiploids. The cause of the variability of pairing capacities among the chromosomes by amphidiploids is probably of cytogenetic nature but the influence of environmental factors is also possible.

The most striking fact from the table 23 is the high fertility of F₂ plants in comparison with nearly sterile F₁ plants. The fertility of the second generation of this amphidiploid was tested on the

plants raised from the selfed plants AJW3, AJW5 and AJW6, and is represented on the table 24.

TABLE 24. Pollen and seed fertility of the second generation of *A. Janczewskii*, Warsaw.

No. of plant	% good pollen	% good seeds	No. of plant	% good pollen	% good seeds
AJW3 (1)	76,8	67,3	AJW6 (8)	80,3	83,9
AJW3 (2)	66,5	50,0	AJW6 (9)	33,5	56,8
AJW3 (3)	62,8	69,3	AJW6 (10)	74,4	69,8
AJW3 (4)	87,0	41,5	AJW6 (11)	82,8	58,8
AJW3 (5)	53,6	50,1	AJW6 (12)	76,7	46,0
AJW3 (6)	61,4	57,9	AJW6 (13)	81,1	63,7
AJW3 (7)	40,0	56,1	AJW6 (14)	60,0	54,6
AJW5 (1)	65,5	64,8	AJW6 (15)	61,1	78,4
AJW5 (2)	86,6	81,0	AJW6 (16)	59,2	63,2
AJW5 (3)	64,0	64,8	AJW6 (17)	68,0	84,3
AJW5 (4)	70,2	73,2	AJW6 (18)	46,5	59,2
AJW6 (1)	52,5	53,3	AJW6 (19)	51,2	32,0
AJW6 (2)	89,8	65,3	AJW6 (20)	80,9	48,5
AJW6 (3)	75,1	70,2	AJW6 (21)	35,9	77,6
AJW6 (4)	66,2	68,5	AJW6 (22)	68,0	50,5
AJW6 (5)	70,0	80,6	AJW6 (23)	72,2	68,5
AJW6 (6)	80,5	76,1	AJW6 (24)	75,2	75,4
AJW6 (7)	52,6	83,3			

The comparison of the tables 23 and 24 shows that the II generation of the amphidiploid is more fertile than the I, both in pollen and seeds. The average fertility of the first generation is 58,7% for pollen and 48,4% for seeds but for the second generation the corresponding values are for AJW3 family 64,0% and 61,7%, for AJW5 family 71,6% and 70,9% and for the AJW6 family 66,4% and 61,5%. The range of variation of the pollen and seed fertility is enlarged in the second generation as compared with the first. The values of minimal and maximal fertility in both generations are as follows: maximum of pollen fertility in I generation is 78,9 in II gen. 89,8% minimum " " " " I " " 42,8 " II " 35,9% maximum " seed " " I " " 74,8 " II " 84,3% minimum " " " " I " " 22,5 " II " 32,0%

Unfortunately the chromosome numbers of the II generation plants were not studied. I have also investigated the fertility of some plants of three generations of *A. Janczewskii* from Edinburgh and of two generations of *A. Janczewskii* from Freiburg. As the time of

origin of these plants is unknown the first generation raised by me, I have denoted N and the following generations by N+1 and N+2. Unfortunately the fertility of the N-generation of *A. Janczewskii* Edinburgh was not fully investigated. One plant with 69,6% of good pollen and 51,0% of good seeds was selfed and gave the N+1 generation. 18 plants were investigated cytologically and in the respect of their fertility and the results are tabulated in the table 25.

TABLE 25. Pollen and seed fertility of the N+1 generation of *A. Janczewskii* Edinburgh.

No. of plant	2n	Pollen fertility	Seed fertility	No. of plant	2n	Pollen fertility	Seed fertility
N+1,1	42	23,0%	37,5%	N+1,10	48	80,1%	78,8%
N+1,1	48	88,9%	71,4%	N+1,11	48	76,0%	86,2%
N+1,1	48	65,3%	43,6%	N+1,12	48	64,8%	56,5%
N+1,1	48	75,5%	63,2%	N+1,13	48	83,3%	61,7%
N+1,1	48	90,5%	84,6%	N+1,14	48	73,6%	56,8%
N+1,1	±47	not flowered		N+1,15	48	94,4%	83,3%
N+1,1	47	49,3%	68,3%	N+1,16	48	88,8%	78,1%
N+1,1	46	63,6%	40,0%	N+1,17	48	82,7%	75,5%
N+1,1	48	82,1%	80,6%	N+1,18	48	65,6%	62,5%

The N+2 generation was obtained by selfing the N+1, 13 plant and designed AN+2, by selfing N+1, 9 plant designed DN+2 and by selfing N+1, 15 plant designed EN+2. The fertility of the N+2 generation is given in table 26.

TABLE 26. Pollen and seed fertility of the N+2 generation of *A. Janczewskii* Edinburgh.

No. of plant	2n	Pollen fertility	Seed fertility	No. of plant	2n	Pollen fertility	Seed fertility
AN+2,1	48	85,7%	87,7%	AN+2,14	48	71,7%	58,3%
AN+2,2	±44	63,5%	74,6%	AN+2,15	48	56,6%	68,9%
AN+2,3	48	81,0%	71,9%	AN+2,16	48	84,3%	79,3%
AN+2,4	45	22,1%	49,8%	DN+2,1	48	86,2%	53,0%
AN+2,5	48	63,9%	80,0%	DN+2,2	48	82,6%	61,8%
AN+2,6	46	71,7%	69,3%	DN+2,3	48	88,1%	62,3%
AN+2,7	48	87,0%	83,3%	DN+2,4	48	91,3%	61,4%
AN+2,8	48	53,5%	52,9%	EN+2,1	48	83,9%	68,1%
AN+2,9	48	87,9%	71,8%	EN+2,2	48	79,1%	68,2%
AN+2,10	48	80,2%	80,6%	EN+2,3	48	88,1%	78,9%
AN+2,11	48	87,2%	62,7%	EN+2,4	48	69,5%	66,3%
AN+2,12	48	82,0%	69,4%	EN+2,5	48	74,6%	66,1%
AN+2,13	48	76,0%	61,3%	EN+2,6	48	68,9%	67,4%

The N generation of *A. Janczewskii* Freiburg was composed of only 1 plant with $2n=48$ and with 73,0% of good pollen and 47,5% of good seeds. From selfing of this plant I have obtained the N₁+1 generation the fertility of which is represented in table nr 27.

TABLE 27. Fertility of N₁+1 generation of *A. Janczewskii* Freiburg.

No. of plant	2n	Pollen fertility	Seed fertility	No. of plant	2n	Pollen fertility	Seed fertility
N + 1,1	44	46,5%	57,5%	N + 1,11	48	72,3%	67,1%
N + 1,2	48	not flowered		N + 1,12	±48	42,1%	47,0%
N + 1,3	48	56,0%	59,5%	N + 1,13	48	89,6%	63,2%
N + 1,4	46	55,9%	39,6%	N + 1,14	48	not flowered	
N + 1,5	47	56,3%	58,9%	N + 1,15	48	80,9%	66,0%
N + 1,6	48	60,0%	59,4%	N + 1,16	48	72,5%	63,3%
N + 1,7	46	55,6%	45,3%	N + 1,17	48	77,5%	74,2%
N + 1,8	48	72,2%	63,6%	N + 1,18	48	75,1%	59,9%
N + 1,9	48	73,2%	73,4%	N + 1,19	48	67,5%	61,1%
N + 1,10	48	83,3%	60,4%				

From these tables following conclusions concerning the fertility of *A. Janczewskii* can be drawn:

1. The fertility of seeds is in average lower than the fertility of pollen.

2. The fertility of the first and second generations of amphidiploids obtained from the cross *A. silvestris* x *A. multifida* v. *magellanica* is lower than the fertility of N+1 and N+2 generations of *A. Janczewskii* from Edinburgh and Freiburg. The average fertility is:

for F ₂	plants	pollen 58,7%	seeds 48,4%
„ F ₃	„	„ 67,3%	„ 64,7%
„ N+1, Edinburgh	„	„ 79,4%	„ 70,2%
„ AN+2,	„	„ 76,7%	„ 72,1%
„ N+1, Freiburg	„	„ 73,4%	„ 64,0%

3. Plants with aneuploid chromosome numbers from 42 to 47 form in average 16—21% of the total number of polyploids in each generation. They are less fertile than the true amphidiploids with $2n=48$. Their average fertility is:

for N+1 Edinburgh	pollen 43,3%	seeds 48,6%
„ AN+2 „	pollen 54,2%	„ 64,5%
„ N+1 Freiburg	„ 51,3%	„ 49,6%

4. In average the fertility of N+1 and N+2 generations of *A. Janczewskii* Edinburgh is equal. However in the descent of any plant they appear plants more fertile than maternal plant. It seems not impossible that by proper and sufficiently long selection of most fertile plants some highly fertile lines could be obtained.

The facts that the seed fertility is lower than the pollen fertility, that not all seemingly good seeds germinate and that they exist shrunken seeds with aborted embryos strongly suggest that by amphidiploids act not only gametic but also zygotic sterility too.

2. The results of interspecific crosses

The results of crosses between different *Anemone* species differing in their chromosome numbers are represented in table 28.

TABLE 28. The results of interspecific crosses.

Cross		No. of good seeds	No. of bad seeds	% of good seeds
<i>A. silvestris</i> x <i>A. multifida v. magell.</i>	2n x 4n	15	181	7,6
<i>A. silvestris</i> x <i>A. mult. v. alba Cluj</i>	2n x 4n	180	164	52,3
<i>A. multifida v. magell.</i> x <i>A. silvestris</i>	4n x 2n	25	120	17,2
<i>A. mult. v. alba Cluj</i> x <i>A. silvestris</i>	4n x 2n	180	22	89,1
<i>A. Jancz.</i> DN+2 x <i>A. mult. v. magell.</i>	6n x 4n	10	260	3,7
<i>A. mult. v. magell.</i> x <i>A. Jancz.</i> DN+2,3	4n x 6n	60	240	20,0
<i>A. mult. v. alba</i> x <i>A. Jancz.</i> N+1,17 Frbg.	4n x 6n	165	121	57,6
<i>A. silvestris</i> x <i>A. Jancz.</i> AN+2,3	2n x 6n	0	286	0,0
<i>A. Jancz.</i> AN+2,3 x <i>A. silvestris</i>	6n x 2n	2	260	0,76

From this table it is evident that the crosses with the maternal species with higher chromosome number generally give better results than the reciprocal crosses. This is especially striking in the crosses of diploid and tetraploid species. *A. multifida v. alba* gives better results than the variety *magellanica* when crossed with *A. silvestris*. The new hexaploid species *A. Janczewskii* shows incompatibility in crosses with parental species. Only the tetraploid *A. multifida* pollinated with pollen from hexaploid *A. Janczewskii* gives from 20% to 50% of seemingly good seeds. The hybrid plants between *A. Janczewskii* and *A. silvestris* forms only 1% of good seeds and the hybrid with *A. multifida* is somewhat more fertile and forms 6,6% of good seeds.

The germination power of seeds obtained from interspecific crosses is variable but generally very low. The percentage of germinated seeds varies in different cases from 0% to a maximum of 55%. The germination period is often extended to two or even three month,

the young plants often die in cotyledon stage and are very difficult to cultivate. In the rather bad culture conditions that I have to my disposition I have never more raised to maturity than 10% of sown seeds. This seems to indicate that the great cytogenetic unbalance of interspecific hybrids causes strong elimination of hybrids during their development.

Greater incompatibility in the crosses $2n \times 4n$ than in the reciprocal ones $4n \times 2n$ was often observed in different genera e. g. in wheats (Watkins 1932), *Galeopsis* (Müntzing 1933) and others. Müntzing (1933) has attempted to explain the cause of the disturbed seed development by alterations in the development of three different tissues with different chromosome number which are in intimate contact: embryo, endosperm and the surrounding somatic tissues of the mother plant. In normal cases i. e. in non-hybrid plants the relation between the chromosome numbers of these tissues is 2:3:2, in the cross $4n \times 2n$ this relation is 3:5:4 and in the cross $2n \times 4n$ this relation is 3:4:2. To this fact Müntzing attributes the cause of the differential incompatibility by reciprocal crosses. The more distant is the chromosome number relation of the three tissues from the normal, the more disturbed are the interrelations of the tissues and the result is the poor development or complete abortion of the seeds.

Incompatibility may be caused not only by zygotic sterility but also by checked pollen tube growth. The differences in success of reciprocal crosses may be due to differences of pollen germination on foreign stigma, to different pollen tube growth rates and to differences of fertilization capacities. The physiological interaction between the maternal tissues and the germinating pollen grains may be very different in reciprocal crosses. Recently Schlosser (1936) in an ingenious experiment has demonstrated that in special cases the difficulty to intercross the polyploids is mainly due to the difference in osmotic pressure of maternal tissue and of pollen tube.

VIII. Discussion

The first experimentally and cytologically proved example of new species formation through the doubling of the chromosome number in a sterile species hybrid was *Primula kewensis* (Digby 1912, Pellew a Durham 1916). In the year 1917, Winge on theoretical grounds has advanced the hypothesis, that by sterile species hybrids the doubling of chromosome number must always have as conse-

quence the total restoration of their fertility. He has supposed that on this way new species can arise in nature. The numerous experimental works carried by different authors have totally confirmed Winge's hypothesis. In the year 1932 Winge has reported already 24 examples supporting his hypothesis. Through the last years the number of analogical cases is greatly enlarged, up to 1939 I have found in literature about 100 cases describing the doubling of the chromosome number in species hybrids. The apparition of amphidiploids was described in various genera of flowering plants and especially numerous cases are known in genera *Nicotiana*, *Brassica*, *Triticum* and *Aegilops*.

In 65 cases describing amphidiploids arising spontaneously in experiments:

43 or 66,1% have arisen in F_2 from the union of unreduced gametes,

24 or 31,5% have arisen in F_1 by somatic doubling of the chromosome number,

4 or 5,2% have arisen from parthenogenetic egg development,

and 3 or 3,9% have arisen through „triploid” stage.

Thus we see that two-thirds of amphidiploids have arisen through the union of unreduced gametes produced by the F_1 hybrids. This is obvious in connection with the generally observed profound disturbances in maturation divisions in partially sterile species hybrids, which lead to unreduced gametes production.

In the table nr. 28 are represented the pairing conditions in 19 F_1 hybrids together with the fertility degree of corresponding amphidiploids. It must be stated that in most cases the data from literature about the fertility and chromosome pairing in hybrids and corresponding amphidiploids are incomplete. The estimation of an average percentage of conjugation in F_1 hybrids needs statistical studies as the conjugation is often variable. The relation between the percentage of conjugation in F_1 and the percentage of seed fertility of the first generation of amphidiploids is represented on the fig. nr 23.

From the table 28 and the fig. 23 it is evident that between the percentage of conjugation in F_1 hybrids and the fertility of amphidiploids it exists an inverse correlation. The function seems to be rather parabolic than a linear but owing to the insufficient number of examined cases and to insufficient exactness of estimation more reaching deductions are dangerous.

TABLE 29. Fertility of amphidiploids together with the conjugation in F_1 .

C r o s s	2n	F_1 -Conjugation a. %	Fertility of amphidipl.		A u t h o r i t y
			seed	pollen	
<i>Raphanus</i> x <i>Brassica</i>	18	18 ₁ = 0%	100%	100%	Karpezenko, 1927a, b
<i>Nicotiana multibracteata</i> x <i>N. suaveolens</i>	40	40 ₁ = 0%	100%	100%	Kostoff, 1937b
<i>Erophila verna</i> n. 8 x <i>E. ver.</i>	47	47 ₁ = 0%	quite normal		Winge 1933
<i>Madia nutans</i> x <i>M. Ramii</i>	17	17 ₁ = 0%	"		Clausen et al. 1936
<i>Aegilops ovata</i> x <i>Triticum dicoccum</i>	28	0-4 _{II} = ca. 14%	42%	84%	Taylor & Leighty 1931
<i>Triticum dicoccoides</i> x <i>Aegilops ovata</i>	28	0-5 _{II} = ca. 17%	42.8%		Kihara & Katayama 1931
<i>Triticum turgidum</i> x <i>Tr. villosum</i>	21	1-4 _{II} = ca. 23%	29.7%		Tschermak 1930, 33 v. Berg 1934a, b, Sando 1935
<i>Aegilops triuncialis</i> x <i>Tr. dicoccum</i>	28	4-5 _{II} = ca. 30%	25%		Oehler 1936, Sorokina 1937c
<i>Aegilops triuncialis</i> x <i>Triticum durum</i>	28	1-8 _{II} = ca. 30%	25%		Oehler 1936
<i>Aegilops caudata</i> x <i>Triticum dicoccum</i>	21	0-7 _{II} = ca. 33%	27.6%		Oehler 1934
<i>Raphanus</i> x <i>Brassica</i>	18	1-5 _{II} = ca. 33%	24%	ca. 60%	Howard 1938
<i>Anemone silvestris</i> x <i>A. multifida</i>	24	3-8 _{II} = ca. 41.5%	8.4%	58.7%	Gajewski 1946
<i>Nicotiana glauca</i> x <i>N. Langsdorffii</i>	21	1-9 _{II} = ca. 45.7%	18%	51%	Kostoff 1938b
<i>Aegilops speltoides</i> x <i>Ae. unbelulata</i>	14	2-5 _{II} = ca. 50%	23.5%	57.2%	Kihara 1937
<i>Nicotiana rustica</i> x <i>N. paniculata</i>	36	12 _{II} = 66%	32%		Sears 1939
<i>N. silvestris</i>	36	12 _{II} = 66%			Singleton 1932
<i>Solanum nigrum</i> x <i>S. luteum</i>	60	24 _{II} = 80%	15.1%	98.2%	Lamerts 1931, 1933a, b
<i>Aquilegia chrysantha</i> x <i>A. flabellata</i>	14	7 _{II} = 100%	25%	35%	Rybin 1929
<i>Crepis rubra</i> x <i>C. foetida</i>	10	5 _{II} = 100%	ca. 14%	25%	Ternowsky 1937
			2.7%	34.5%	Jorgensen 1928
					Skalińska 1935
					Poolle 1931, 1932

From the shape of the curve in fig. 23 it is to be seen that even little degree of pairing in F_1 among few chromosomes causes a relatively great lowering of the amphidiploids fertility, the further increase in the pairing degree has less influence on the amphidiploids fertility. The differences in individual behaviour of various amphidiploids are great, they seem to depend on various agents such as the number of chromosomes, size of chromosomes and the number of chiasmata, which have great influence on the mechanism of dif-

Fig. 23. The relation between the chromosome conjugation in F_1 and the amphidiploid fertility.

ferential affinity. The amphidiploids with small chromosomes which form few chiasmata possess smaller number of polyvalents and are more fertile. The mode of apparition of the amphidiploids has here great significance. The case of *Raphanobrassica* which was investigated by Karpezenko (1927a, b) and Howard (1938) demonstrates direct correlation between pairing in F_1 and fertility. In the same cross Karpezenko has used the forms giving only univalents in F_1 and Howard the forms giving ca 33% of pairing in F_1 . The fertility of corresponding amphidiploids is lowered from about 100% to only 24%.

What are the causes of the decrease of amphidiploids fertility by the increase of pairing degree in F_1 hybrids? No doubt, we have to do by the amphidiploid sterility with gametic and zygotic sterility

together. Many facts seem to indicate that the gametic sterility alone is insufficient to account for amphidiploids sterility. The percentage of seeds sterility by amphidiploids is always considerably lower than the pollen sterility. There is no reason to believe that the sterility of female gametes is greater than the sterility of male gametes and even, on the contrary, it is known that many hybrids are more fertile in eggs than in the pollen. By many amphidiploids the zygotic sterility was observed directly. The observations indicate that in ovulae the fertilization has occurred but the development of the embryos was soon stopped (Skalińska 1935, Howard 1938), also seemingly good seeds of amphidiploids don't germinate (Skalińska 1935, Karpezenko 1937). The causes of amphidiploid infertility can be various, they depend firstly on the mode of amphidiploids apparition and on the character of pairing in F_1 and in amphidiploids themselves.

Amphidiploids according to their mode of origin can be divided in two groups: I — origin by somatic doubling in F_1 and II — origin by the union of two unreduced gametes formed by F_1 plant. Each of these two groups can be subdivided according to the mode of pairing in F_1 in two types: a) — these that have arisen from asyndetic hybrids and b) — these that have arisen from hybrids with complete or partial pairing. This question was recently discussed by Howard (1938) and this author concludes that the amphidiploids of the Ia group should be just as fertile as a normal diploid plant. The amphidiploids of the group IIa should also be quite fertile except the rare cases when amphidiploids are formed with the constitution of $n+1$ chromosomes of one species and $n-1$ chromosomes of the second species. Such uncomplete amphidiploids can arise as a result of meiotic irregularities before diad formation. The fertility of amphidiploids of the group Ib depends on the pairing behaviour at meiosis. If only bivalents are formed and the pairing due to differential affinity is entirely intraspecific and the multivalents are lacking, then the amphidiploids should be quite fertile. But, if interspecific conjugation occurs or if multivalents are formed, then the amphidiploid should be only partially fertile. The amphidiploids of the group IIb are always only partially fertile. In F_1 the pairing of only partially homologous chromosomes leads to the production of diads with unequal segmental constitution. Assuming that partially homologous chromosomes are built from homologous and nonhomologous segments and if the chiasma and crossing-over is formed between the nonhomologous segments of two chromosomes and their centro-

meres then the diads are formed with chromosomes showing deficiency and reduplication of small segments. The amphidiploids arisen from the union of such diads are not true amphidiploids as they possess some reduplicated and some lacking segments and are only partially fertile. Also, by the conjugation of partially homologous chromosomes in amphidiploids, gametes should arise with reduplicated or lacking segments in some chromosomes. The union of such gametes would in certain conditions produce unviable zygotes. In following generations of such amphidiploids the irregularities of this type can be eliminated through the gradual elimination of gametes and zygotes with unviable segmental constitution. This would account for the observed higher fertility of latter amphidiploids generations. The increase of amphidiploids fertility in latter generations was observed by Sando 1935 (amph. *Triticum turgidum* x *villosum*), Katayama 1935 (*Triticum dicoccoides* x *Aegilops ovata*), Oehler 1936 (*Aegilops triuncialis* x *Tr. durum*), Howard 1938 (*Raphanobrassica*), Kostoff 1938b (*Nicotiana Vavilovii*) Poole 1932 (*Crepis rubra* x *foetida*). In following generations of these amphidiploids, notwithstanding higher average fertility, some less fertile or even quite sterile plants arise indicating a segregation of some sterility factors. The increase of fertility is correlated with a decrease of meiotic disturbances. (Kostoff 1938b, Katayama 1935). It is possible that the recovery of fertility of amphidiploids is not only due to chromosomal factors but also to the action of some extra-chromosomal factors, such as the harmonizing of the interaction between the genomes and the cytoplasm or between the two genomes. These ideas are dealt by Navashin (1935), Michaelis (1937), Wettstein (1928), Kostoff (1938b) and others. It is obvious that the genome of an amphidiploid could not be regarded as an arithmetic sum of two genomes of parental species. This is a new species with his own new physiological equilibrium. The discussion of this topic needs a more profound development of the investigations on the physiology of meiosis.

91,75% of amphidiploids obtained experimentally show more or less distinct cross-sterility or hybrid sterility when crossed to parental species. The barriers are mainly caused by quantitative differences in the chromosome numbers between amphidiploids and their parents but in some cases by the differences in physiological properties.

The process of amphidiploidity has greatly influenced the evolution of many genera of flowering plants. It seems that in some

genera such as *Nicotiana*, *Triticum*, *Brassica* and some others the species of amphidiploid origin are especially common.

Goodspeed (1934) points out, that in *Nicotiana* it is possible to obtain from every species hybrid an amphidiploid when only a sufficiently large number of individuals are grown. He believes that all American *Nicotiana* species with 48 chromosomes are amphidiploids. Goodspeed, Clausen, Kostoff and others have brought a conclusive evidence that *N. tabacum* and *N. rustica* with 48 chromosomes represent amphidiploids from the crosses between *N. silvestris* x *N. tomentosiformis* and among related species to *N. paniculata* and *N. undulata* respectively. The very vaste investigations on wheat species especially these of the Kihara school demonstrate that the hexaploid wheat species are probably amphidiploids arisen from the hybrids between diploid and tetraploid wheat species. For *Brassica* Morinaga, Karpezenko, U and others have accumulated many interesting data showing that the species with higher chromosome numbers have originated by amphidiploidy from the species with lower chromosome numbers. For the cultivated plants undoubtedly the amphipolyploidy has played a very decisive role in the formation of new species. The cultivation of different species together and the selection of bigger individuals by mankind trough many centuries have created many times propitious conditions for the origin of new amphidiploid species.

But also in nature there exist species of amphidiploid origin. The synthesis of *Galeopsis tetrahit* by Müntzing (1930a, b), *Salix cinerea* by Nilsson (1931), Hakansson (1938) and *Brassica Napus* by U (1935) are the best examples of the origin of new species known from nature through the process of amphidiploidy. For many others species many conclusive data have been accumulated showing their origin through hybridity and the doubling of the chromosome number as e. g. for *Aegilops triuncialis* (Sorokina 1937), *Aesculus carnea* (Skovsted 1929), *Galium Schultesii* (Fagerlind 1937), *Iris versicolor* (Anderson 1936), *Narcissus dubius* (Fernandes 1937), *Phleum pratense* (6x) (Gregor & Sansome 1930), *Pentstemon neotericus* (Clausen 1933), *Prunus domestica* (Rybin 1936), *Rosa Wilsoni* (Blackburn & Harrison 1924), *Rubus maximus* (Rozanova 1934, 1938) and *Spartina Townsendii* (Huskins 1930). Huskins has demonstrated that *Spartina Townsendii* arose probably in the nineteenth century on the southern coast of England and had so great expansion power that has overrun some of the territory previously occupied by *Spartina stricta* and *S. alterniflora*, his putative

ancestors. This example shows that some new amphidiploid species arisen in nature have sufficiently great capacities in the struggle of life or even surpass their ancestors.

The amphidiploidity is only a part of a more large topic of the role of polyploids in the evolution of plant kingdom. The polyploids form about one half of the species of flowering plants with known chromosome number. In northern parts of Europe the polyploids seem to form the majority of the here existing vegetation (Tischler 1935, 1937). The process of polyploidity has played and plays probably now-to-day a great role in the reoccupation by polyploids of the grounds left by the last great glaciation. Many intra-specific polyploid races exist among plants of northern hemisphere with special physiology, ecology, or even distribution and represent the first steps to new species formation (Müntzing 1936, Hagerup 1932).

The experimental work on polyploidity and mutation forms a new era in evolution problem. The previous work on evolution was of merely speculative nature and after a long period of theoretical researches the mechanics of species origin was unknown. Dobzhansky in his recent book „Genetics and the origin of species”, points out that polyploidity is only one of the numerous ways on which new species arose. He names this way „a cataclysmic origin of species” which is opposite to the very long and slow way of species origin through the gradual accumulation of minute mutational changes.

IX. Summary

1. Attempts have been made to repeat Janczewski's crosses of *Anemone multifida* ($n=16$) with *A. silvestris* ($n=8$). From this cross Janczewski has obtained fertile and constant progeny which was probably of amphidiploid origin.

2. The morphological features of the two parental species are described.

3. A highly sterile, triploid ($n=12$) hybrids were obtained and their biometric description together with the parental species is given.

4. From few good achenes produced by triploid hybrids hexaploid F_2 plants with $2n=48$ were obtained.

5. The description of hexaploids is given and their morphology is compared with the parental species and the triploid ancestor.

The hexaploids have larger cells, stomata, sepals, pollen-grains and seeds and have broader and thicker leaves than the triploids.

6. Hexaploid plants identical with the F_2 amphidiploids were obtained from Botanical Gardens of Edinburgh, Freiburg, Stäfa and Dublin. It is believed that they have arisen in similar way from the hybrids between *A. silvestris* and *A. multifida*. They are described as new species and named *Anemone Janczewskii*.

7. The correlations between chromosome numbers and the size and shape of cells and organs in the whole *Anemone* material are discussed.

8. The cytological features of the parental species both of somatic and meiotic divisions are described. *A. multifida* has twice the same somatic chromosomes complement as by *A. silvestris*. The course of meiosis by the two parental species is quite normal.

9. The meiotic division in P. M. C. of F_1 hybrids shows many irregularities. At M I 3 to 8 bivalents are formed and the rest of chromosomes remain unpaired. The univalents lag on the spindle and divide equationally at late anaphase or pass undivided to the poles. Many univalents are lost in cytoplasm. At the tetrad stage diads were observed probably as the result of homeotypic spindles fusion.

10. The F_1 hybrids form about 0,3% of viable pollen grains. The average diameter of the latter is as large as that of the amphidiploids which suggests that they are unreduced.

11. The amphidiploids with $2n=48$ obtained in F_2 of the mentioned cross have abnormal course of meiosis in P. M. C. At MI beside P. M. C. with only 24 bivalents also numerous cells exist containing a variable number of uni-, tri- and tetravalents. In the latter many irregularities in the chromosome disjunction and distribution were observed leading to the deficient gametes production.

12. Similar meiotic features were observed by hexaploid plants from Edinburgh and also by aneuploids with $2n=42-47$, obtained from the progeny of selfed plants of *A. Janczewskii* from Edinburgh.

13. The pollen and seed fertility of the investigated plants was studied. The parental species are quite fertile. The F_1 hybrids are highly sterile they produce 0,12—0,31% of good pollen grains and 0,1—3,2% of good seeds. The germination power of these seeds is low. The amphidiploids obtained in F_2 are only partially fertile, they produce 42,8—78,9% of good pollen and 31,7—74,8% of good

seeds. The fertility of amphidiploids from Edinburgh is somewhat higher, the average fertility of $N+1$ generation is for pollen grains 79,4⁰/₁₀₀ and for seeds 70,2⁰/₁₀₀. The fertility of aneuploids is distinctly lower. In the progeny of selfed amphidiploids, plants arose with higher and lower fertility than the parental plant.

14. The crossability between the amphidiploid and the parental species is minime and the resulting hybrids are highly sterile. The reciprocal crosses in *Anemone* give unequal results, the crosses when the maternal species possesses higher chromosome number give better results than the reciprocal ones.

Botanical Garden, University of Warsaw.

References

- Anderson E., (1936). „The species problem in *Iris*“. Ann. Missouri Bot. Gard., 23, 57—509.
- Becker G., (1932). „Experimentelle Analyse der Genom- und Plasmonwirkung bei Moosen. III. Osmotischer Wert heteroploiden Pflanzen“. Z. indukt. Abstamm. u. Vererb. Lehre, 60, 17—38.
- Berg K. H. von, (1934 a). „Cytologische Untersuchungen an *Triticum turgidovillosum* und seinen Eltern“. Z. indukt. Abstamm. u. Vererb. Lehre, 67, 342—373.
- (1934 b). „Cytologische Untersuchungen an den Bastarden des *Triticum turgidovillosum* und an einer F_1 *Triticum turgidum* $\times$ *villosum*. III. Teil.“ Z. indukt. Abstamm. u. Vererb. Lehre, 68, 94—126.
- Blackburn K. B. & Harrison J. W. H., (1924). „Genetical and cytological studies in hybrid roses. I. The origin of a fertile hexaploid form in the *Pimpinellifoliae-Villosae* crosses“. Brit. J. Exp. Biol. 1, 557—570.
- Clausen J., (1933). „Cytological evidence for the hybrid origin of *Pentstemon neotericus* Keck.“ Hereditas, Lund, 18, 65—76.
- Clausen J., Keck D. D. & Hiesey W. M., (1936). „Experimental taxonomy“. Carneg. Inst. of Wash. Ann. Rep. of Div. Plant. Biol. for 1935—36, 208—214.
- Digby L., (1912). „The cytology of *Primula kewensis* and other related *Primula* hybrids“. Ann. of Bot. 26, 357—388.
- Fagerlind F., (1937). „Embryologische, zytologische und bestäubungsexperimentelle Studien in der Familie Rubiaceae nebst Bemerkungen über einige Polyploiditätsprobleme“. Acta Horti Bergiani, 11, 108—470.
- Fernandes A., (1937). „Sur l'origine du *Narcissus dubius*“. Bol. Soc. Brot. 12, 93—116.
- Goodspeed T. H., (1934). „*Nicotiana* phylaxis in the light of chromosome number, morphology and behaviour“. Univ. Calif. Publ. Bot. 17, 369—398.
- Gregor J. W. & Sansome F. W., (1930). „Experiments on the genetics of wild populations. II. *Phleum pratense* L. and the hybrid *P. pratense* L. $\times$ *P. alpinum* L.“. J. Genet. 22, 373—387.
- Hagerup O., (1932). „Über Polyploidie in Beziehung zu Klima, Oekologie und Phylogenie“. Hereditas, Lund, 16, 19—40.

- Hakansson A., (1938). „Zytologische Studien an Salix-Bastarden“. Hereditas, Lund, 24, 1—32.
- Howard H. W., (1938). „The fertility of amphidiploids from the cross *Raphanus sativus* × *Brassica oleracea*“. J. Genet. 36, 239—273.
- Huskins C. L., (1931). „The origin of *Spartina Townsendii*“. Genetica, 12, 531—538.
- Janczewski E., (1889). „Les hybrides du genre *Anemone*. I. II“. Bull. intern. Acad. Sci. Cracovie, 1889, 228—230.
- (1892). „Les hybrides du genre *Anemone* III“. Bull. intern. Acad. Sci. Cracovie, 1892, 228—230.
- Jorgensen C. A., (1928). „The experimental formation of heteroploid plants in the genus *Solanum*“. J. Genet. 19, 133—211.
- Karpetchenko G. D., (1927 a). „The production of polyploid gametes in hybrids“. Hereditas, Lund, 9, 349—368.
- (1927 b). „Polyploid hybrids of *Raphanus sativus* L. × *Brassica oleracea* L.“. Bull. appl. Bot. Gen. Pl. Breed. 17, 305—394.
- Katayama Y., (1935). „Further investigations on synthetised octoploid *Aegiloticum*“. J. Coll. Agric. Tokyo Imp. Univ., 13, no. 4.
- Kihara H. a. Katayama Y., (1931). „Zur Entstehungsweise eines neuen konstanten oktoploiden *Aegilotriticum*“. Cytologia, 2, 234—255.
- Kihara H. a. Ono T., (1926). „Chromosomenzahlen und systematische Gruppierung der *Rumex*-Arten“. Z. Zellforsch. mikr. Anat., 4, 475—481.
- Kostoff D., (1937). „Studies on polyploid plants. XVII. *Nicotiana multivalvis* ($2n=48$) × *Nicotiana suaveolens* ($2n=32$) amphidiploid ($2n=80$)“. C. R. Acad. Sci. U. R. S. S., 14, 215—217.
- (1938 a). „Directed heritable variations conditioned by euploid chromosome alterations“. J. Genet., 36, 447—468.
- (1938 b). „Studies on polyploid plants. XXI. Cytogenetic behaviour of the allopolyploid hybrids *Nicotiana glauca* Grah. × *Nicotiana Langsdorffii* Weimm. and their evolutionary significance“. J. Genet. 37, 129—209.
- Kostoff D. a. Kendall J., (1937). „Studies on polyploid plants III. Cytogenetics of tetraploid tomatoes“. Gartenbauwiss., 9, 20—44.
- Lammerts W. E., (1931). „Interspecific hybridisation in *Nicotiana*. XII. The amphidiploid *rustica paniculata* hybrid; its origin and cytogenetic behaviour“. Genetics, 16, 191—211.
- (1933 a). „The *Nicotiana rustica-paniculata* amphidiploid derivatives“. Cytologia, 4, 46—51.
- (1933 b). „An experimentally produced secondary polyploid in the genus *Nicotiana*“. Cytologia, 5, 38—45.
- Langlet O., (1927). „Beiträge zur Zytologie der Ranunculaceen“. Svensk Bot. Tidsk. 21, 1—15.
- Larier L. H. W., (1932). „Chromosome variation and behaviour in *Ranunculus*“. J. Genet., 26.
- Lawrence W. J. C., (1936). „The origin of new forms in *Delphinium*“. Genetica, 18, 101—115.
- Lilienfeld F. A., (1936). „Karyologische und genetische Studien an *Fragaria*. I. Ist *Fragaria elatior* eine autotetraploide Pflanze?“. Jap. J. Bot. 7, 119—149.
- Michaelis P., (1937). „Untersuchungen zum Problem der Plasmavererbung. Sammelreferat“. Protoplasma, 27, 284—289.

- Miczyński K., (1892). „Mieszance zawileców (*Anemone* L.) opracowane pod względem anatomicznym“. Rozpr. Wydz. mat.-przr. Akad. Um. Kraków, 24, 1—32.
- Moffet A. A., (1932). „Chromosome studies in *Anemone*. I. A new type of chiasma behaviour“. *Cytologia*, 3.
- Muntzing A., (1930). „Über Chromosomenvermehrung in *Galeopsis*-Kreuzungen und ihre phylogenetische Bedeutung“. *Hereditas*, Lund, 14, 153—172.
- (1933). „Hybrid incompatibility and the origin of polyploidy“. *Hereditas*, Lund, 18, 33—55.
- (1936). „The evolutionary significance of autopolyploidy“. *Hereditas*, Lund, 21, 263—378.
- Navashin M., (1934). „Chromosome alterations caused by hybridisation and their bearing upon certain general genetic problems, *Cytologia*, 5, 169—203.
- Nilsson H., (1931). „Über das Entstehen eines ganz cinerea-ähnlichen Typus aus dem Bastard *Salix viminalis* × *caprea*“. *Hereditas*, Lund, 15.
- Oehler E., (1934). „Untersuchungen an der neuen konstanten additiven *Aegilops* Weizenbastarden“. *Züchter*, 6, 263—270.
- (1936). „Untersuchungen an einen neuen konstant-intermediären additiven *Aegilops*-Weizenbastard“. *Züchter*, 8, 29—33.
- Pellaw C. & Durham F. M., (1916). „The genetic behaviour of the hybrid *Primula kewensis* and its allies“. *J. Genet.* 5.
- Poole C. F., (1931). „The interspecific hybrid *Crepis rubra* × *foetida* and some of its derivatives. I.“. *Univ. Calif. Pub. Agr. Sci.*, 6, 169—200.
- (1932). „The interspecific hybrid *Crepis rubra* × *foetida* and some of its derivatives. II.“. Two selfed generations from amphidiploid hybrid“. *Univ. Calif. Pub. Agr. Sci.*, 6, 231—255.
- Propach H., (1939). „Was ist eigentlich unsere Rittersporne“. *Gartenschönh.* 1939.
- Renner O., (1929). „Arbastarde bei Pflanzen“. *Hdbch. d. VererbWiss.* Lief. 7, 1—161.
- Rosanowa M. A., (1934). „Modes of form genesis in the genus *Rubus*“. *Bot. Ž. U. R. S. S.* 19, 376—384.
- (1938). „O polymorfnom *wosnikovenii* vidow.“. *C. R. Acad. Sci. U. R. S. S.*, 18, 681—683.
- Rosenthal C., (1936). „Chromosomenstudien an *Pulsatilla*“. *Jahrb. f. wiss. Bot.*, 83, 809.
- Rybin W. A., (1929). „Über einem allotetraploiden Bastard von *Nicotiana Tabacum* × *N. sylvestris*“. *Ber. Dtsch. Bot. Ges.*, 47, 385—394.
- (1936). „Spontane und experimentell erzeugte Bastarde zwischen Schwarzdorn und Kirschlordele und das Abstammungsproblem der Kulturpfleume“. *Planta*, 25, 22—58.
- Sando W. J., (1935). „Intergeneric hybrids of *Triticum* and *Secale* with *Haynaldia villosa*“. *J. Agric. Res. Wash.*, 51, 759—800.
- Sax K. a. H. J., (1937). „Stomata size and distribution in diploid and polyploid plants“. *J. Arnold Arb.*, 18, 164—172.
- Schlösser L. A., (1936). „Befruchtungsschwierigkeiten bei autopolyploiden und ihre Überwindung“. *Züchter*, 8, 295—301.
- Sears E. R., (1939). „Amphidiploids in the *Triticinae* induced by colchicine“. *J. Hered.*, 30, 38—43.
- Singleton W. R., (1932). „Cytogenetic behaviour of fertile tetraploid hybrids of *Nicotiana rustica* and *paniculata*“. *Genetics*, 17, 510—544.

- Skalińska M., (1935). „Cytogenetic investigations of an allotetraploid *Aquilegia*“. Bull. Acad. Sci. e. Lett. Cracovie. Ser. B., 33—63.
- Skovsted A., (1929). „Cytological investigations of the genus *Aesculus* L.“. Hereditas, Lund, 12, 64—70.
- Sorokina O. N., (1937). „New *Aegilops*-wheat amphidiploids“. Bull. appl. Bot. Gen. Pl. Breed. Ser. 2, no. 7, 161—173.
- Taylor J. W. a. Leighty C. E., (1931). „Inheritance in a „constant“ hybrid between *Aegilops ovata* and *Triticum dicoccum*“. J. Agric. Res. Wash. 43, 661—679.
- Ternovsky M. F., (1936). „Amphidiploid shoots of the F_1 hybrid *Nicotiana tabacum* $\times$ *syvestris*“. C. R. Acad. Sci. U. R. S. S., 15, 151—152.
- Tischler G., (1935). „Die Bedeutung der Polyploidie für die Verbreitung der Angiospermen“. Bot. Jahrb., 67, 1—36.
- (1937). „Die Halogenflora der Nordsee in Lichte zytologischer Forschung“. Cytologia, Fujii Jub. I, 162—170.
- Tschermak E., (1930). „Neue Beobachtungen am fertilen Artbastard *Triticum turgidovillosum*“. Ber. dtsch. Bot. Ges., 48, 400—407.
- (1933). „Weitere Studien am fertilen, konstanten Artbastard *Triticum turgidovillosum* und seinem Verwandten“. I. Teil. Z. indukt. Abstamm. und VererbLehre, 66, 180—219.
- U. N. (1935). „Genome-analysis in *Brassica* with special reference to the experimental formation of *B. napus* and the peculiar mode of fertilization“. Jap. J. Bot., 7, 389—452.
- Ulbrich E., (1906). „Über die systematische Gliederung und geographische Verbreitung der Gattung *Anemone*“. Engler Bot. Jahrb., 37, 172—334.
- Watkins A. E., (1932). „Hybrid sterility and incompatibility“. J. Genet., 25, 125—169.
- Wettstein F. v., (1928). „Morphologie und Physiologie des Formwechels der Moose II.“. Biblioth. Genet., 10, 1—216.
- (1938). „Experimentelle Untersuchungen zum Artbildungsproblem. I. Zellgrößenregulation und Fertilvermögen einer polyploiden *Bryum*- Sippe“. Z. indukt. Abstamm. u. VererbLehre., 74.
- Winge O., (1917). „The chromosomes. Their numbers and general importance“. C. R. Trav. Lab. Carlsberg. 13, 131—275.
- (1932). „On the origin of constant species hybrids“. Svensk Bot. Tidskr. 26, 107—122.
- (1933). „A case of amphidiploidy within the collective species *Erophila verna*“. Hereditas, Lund, 18, 181—191.

Anemone silvestris (right), *A. multifida* v. *magellanica* (left and their F_1 hybrid (middle). Note striking hybrid heterosis.

The triploid F_1 hybrid (middle), hexaploid F_2 plant (left) and *A. Janczewskii* from Edinburgh (right).