Allium test — some questions

MAŁGORZATA WIERZBICKA, DANUTA ANTOSIEWICZ

Laboratory of Electron Microscopy, Institute of Botany, University of
Warsaw, Krakowskie Przedmieście 26/28, 00-927 Warsaw, Poland

(Received: June 10, 1987. Accepted: September 28, 1987)

Abstract

The paper concerns some problems connected with the experiments with the use of the *Allium* test. The variation degree of the material analysed is also presented. For the sake of standard experiments using the *Allium* test, it is suggested to grow onion bulbs in deionized water or in Hoagland nutrient. Taking as an example the lead compounds, the paper proves that the kind of water which has been used to prepare the analyzed solution affects significantly the toxicity level of the tested compound. This may be the reason why the results of experiments made by various researchers are incomparable. The differences in metabolism between individual roots are studied as well, both in the control and the experimental materials. Eventually, it is suggested that the roots of each bulb should be treated individually in all *Allium* test experiments. The presented approach largely increases accuracy and repeatability of the *Allium* test results.

Key words: *Allium cepa* L., *Allium* test, mitotic activity, root tip, lead, lead in apoplast, variation in test material

INTRODUCTION

The *Allium* test, just like other plant tests, is often used to detect potential environmental mutagens that can threaten human health (Grant and Zura 1982). In spite of big differences in organization and metabolism between plant and animal cells, it is possible to extrapolate the results obtained for animal organisms (Grant 1978). Great similarities in the kind and frequency of chromosomal aberrations between plant and animal cells have repeatedly been

found and reported (Nilan 1978, Serres 1978, Fiskesjö 1982). Also similar chemical transformations of compounds in both cell types have been observed (Antosiewicz 1984, 1987, 1988, Menn and Still 1977, Plewa 1978, Scott et al. 1978, Spano and Takahashi 1981).

While using a test it is important to know the variation range of the experimental material. It has been already described how big can be the differences in the localization of lead salts both between individual tissues of root tip, and between consecutive ground meristematic tissue layers (Wierzbicka 1987a, b). It is even of greater importance as ground meristematic tissue cells are those analysed in squashed slides for the *Allium* test. Thus it is essential to have such an experimental procedure which would make the material the most homogenous. Although the *Allium* test has been widely used in various studies, serious problems concerning the material non-homogeneity have not been discussed so far.

The paper aims to present the difficulties and problems arisen in the course of experiments with the *Allium* test, as well as to suggest how to solve them.

MATERIAL AND METHODS

Allium cepa L. bulbs were prepared as described by Wierzbicka (1987c) in order to obtain bigger number of roots of the same length. The bulbs were grown in black containers of 320 cm³ at room temperature, in tap distilled deionized water, or 1/8 Hoagland nutrient solution (Peterson et al. 1978) supplemented with A-Z microelements.

During growing after they had reached the length of 3 mm onion adventitious roots, were constantly aerated till the end of the experiment. To prevent any appearance of fungi or microorganisms, all liquids in the containers and the containers themselves were changed each 24 h. Just before that all liquids were aerated for 24 h.

The initial material for the *Allium* test consisted of adventitious roots, each 3 ± 1 cm long. To start their incubation some bulbs with roots were transferred either to lead salt solutions (pH 6.5) or to ledakrin solutions. Others remained in water or nutrient as controls.

In the course of all experiments, because of the 24 hour rythm, the incubation was started at 8 a.m. Thus, in Table 1 and Figs. 15, 16 the initial (zero) time falls on 8 a.m., 6 h time — on 2 p.m. and 12 h time — on 8 p.m. etc.

Randomly chosen roots were cut off after the appropriate incubation times. Then the breadth of root tips (at the hight of 1 mm) was measured in a microscope with an eyepiece with a microscale. After that the roots were fixed in alcohol-acetic acid (3:1) and rinsed in 96% alcohol, and then kept in 80% alcohol.

Permanent squashed slides were made from 2 mm long root tips on dry ice, after maceration and staining with acetic orcein — 1 n HCl (9:1). Each slide prepared in this way was used to calculate (after each treatment time there were at least 5 slides):

1. Dividing cells per each 1000 meristematic cells in 4 random fields with 250 meristematic cells each.
2. The number of disturbances in each mitotic phase per each 50 prophase, 50 metaphases, 20 anaphases and 50 telophases.
3. Interphase cells with micronuclei (Degrassi and Rizzoni's micronucleus test, 1982) per the total of 1000 meristematic cells in 4 random fields with 250 meristematic cells each.

Autoradiographic studies of the nucleic acids and proteins synthesis level were carried out with labelled precursors such as ^3H -thymidine (111 kBq cm^{-3}), ^3H -leucine (296 kBq cm^{-3}) and ^3H -uridine (370 kBq cm^{-3}). For this purpose 10 roots were cut off at the base from 5 onions (2 roots each) at one hour before the end of the incubation time. Then the incubation liquid in which the roots had been grown was poured into 3 groups of 20 cm^3 vessels, a labelled precursor added into each and the cut off roots placed back in the liquid. After one hour the roots were taken out, fixed in alcohol-acetic acid (3:1), rinsed in 96% alcohol, and then kept in 80% alcohol. Then squashed slides were prepared according to the procedure described above. The slides were covered with Ar-10 Kodak photographic emulsion and exposed for 14 days at 4°C .

The autoradiographs were used to calculate (after each treatment time there were usually 5 autoradiographs):

1. The number of labels above each of 60 nuclei and the number of cells with labels per 1000 interphase cells in 4 random fields in each ^3H -thymidine-treated autoradiograph.
2. The number of labels above each cell of 1000 interphase cells in 4 random fields in each ^3H -leucine-treated autoradiograph.
3. The number of labels above each nucleoli in 100 interphase cells in 4 random fields in each ^3H -uridine-treated autoradiograph.

Histochemical analysis of lead localization in roots was carried out by rhodizonate method (Wierzbicka 1987b). Thus, the whole roots were immersed in the 0.2% solution of sodium rhodizonate just after the incubation. After 1 h root colour was observed. In the lead presence they stained red.

Autoradiographic analysis of lead localization was undertaken after adding to the incubation liquids 3.77 kBq cm^{-3} of lead isotope ($^{210}\text{Pb}(\text{NO}_3)_2$, of $0.7 \text{ MBq mg}^{-1} \text{ Pb}^{2+}$ specific activity). Additionally, the incubation liquids contained $2.5 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ from $\text{Pb}(\text{NO}_3)_2$ in water and $15 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ from $\text{Pb}(\text{NO}_3)_2$ in Hoagland nutrient (without PO_4^{-2}). The incubation time was 24 h. The method had been thoroughly described by Wierzbicka (1987a). The material was the roots growing from onion bulbs, contrary to

some other isotope experiments where the roots were cut off. Semithin, longitudinal sections from root tips were covered with AR-10 Kodak photographic emulsion. The exposure time was 12 days at 4°C.

Ultrastructural studies of the lead localization were carried out by classical technique of electron microscopy. The roots were incubated for 1-24 h in $2.5 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ from PbCl_2 and $\text{Pb}(\text{NO}_3)_2$ in nutrient solution, then their tips were fixed in 2% glutaraldehyde and postfixed in osmium tetroxide (the both fixatives prepared in 0.1 M cacodylate buffer). The technique applied had been described by Wierzbicka (1987a, b).

The analysis of lead content in liquids in the course of incubation was carried out by dithizone method (Błok 1955) after the roots had been treated with 1.0; 2.5; $3.0 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ from $\text{Pb}(\text{NO}_3)_2$ and PbCl_2 for 1, 3, 6, 12 and 24 hours. These lead solutions were prepared with deionized water.

The results obtained were analysed statistically with t-Student test.

RESULTS AND DISCUSSION

GROWTH MEDIUM FOR ONION ROOTS

Tap water has been recommended to grow onion roots in (Grant 1982). Considering the varying chemical composition of the water depending even on the time it flows in the pipes (Fiskesjö 1981) and having in mind the antagonism of ions the use of tap water seems to include in the experiment a factor which is both variable and uncontrollable. This is the cause why it is impossible to repeat the same experimental conditions.

Water distilled in metal distillers, which are mostly used in laboratories, has very low pH (about 4-5). That is why in our experiments we grew onion in deionized water (pH 6-7). The results obtained were satisfactory, the numbers of disturbances of cell divisions was small, the value of mitotic index high (Table 1), and the result of micronucleus test was very good ($\bar{x} = 0.05\%$). Such water has been already used to grow onion test seedlings (Mäkinen 1963).

Table 1
Mitotic activity and metaphase disturbance in the course of onion root growth in deionized water
(mean values of 6 roots)

Time (h)	0	1	3	6	12	24	30	36	48	54	60	72	78	84	96
Mitotic index (%)	10.5	9.8	9.6	10.0	10.2	10.6	11.0	10.5	10.3	8.5	6.8	11.0	8.8	9.3	9.2
Metaphase disturbance (%)	0.5	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.3	0.6	0.0	0.0	0.0	0.0

However, it appears that in long experiments (5-14 days) onions are better grown in 1/8 Hoagland nutrient solution supplemented with A-Z microelements. The growth of roots in deionized water and in the nutrient solution is compared in Fig. 1. Then the number of disturbances of cell divisions is low as well, and the average value of mitotic index ranges from 7.4 to 10.6.

Fig. 1. Average increase of root length in the course of onion growing in deionized water (1) and 1/8 Hoagland nutrient (2). Initial root length 3 cm

DIFFERENT WATERS AND CHEMICAL COMPOUND ACTIVITY

The *Allium* test is used to find how different chemical compounds influence cell divisions in onion root tips. Yet it has turned out that the activity of the analysed compound highly varies with the kind of water (tap, distilled or deionized), or nutrient used to prepare test solution. The strongest effect of the tested compound appeared when it had been added to deionized water, and this effect was the weakest when the compound was given in nutrient. To illustrate this phenomenon the concentration ranges of lead salts, prepared in deionized water and nutrient were compared in Table 2. The concentrations of Pb^{2+} were chosen so as to obtain their analogous reaction on onion roots. It was found that the effect of lead salts is at least five fold weaker in the case of nutrient than in that of deionized water (Table 2). From analogous observations of lead added to nutrient and tap water it was inferred that the effect of lead salts in both solutions was similar, whereas the effect of lead salts in

distilled water was a little weaker than when the salts were added to deionized water. Such a relation was explained due to the studies on lead localization in roots.

The total lead content in live non-fixed roots was studied by rhodizonate method. Basing on root staining intensity with lead rhodizonate it was found that the total lead content in roots was higher after treating them for 24 h with $15 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ in nutrient (Fig. 2) than when roots were treated for 24 h with $2.5 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ in deionized water (Fig. 3). The result is surprising because the physiological reaction of roots of the both lead doses was similar (Table 2).

Table 2

The comparison between the effect of lead salts on *Allium cepa* L. roots, those prepared in deionized water and those prepared in 1/8 Hoagland nutrient (without PO_4^{2-})

The effect of Pb^{2+} in 24 h treated roots	Pb^{2+} (mg dm^{-3})	
	$\text{H}_2\text{O} +$ either PbCl_2 or $\text{Pb}(\text{NO}_3)_2$	nutrient + either PbCl_2 or $\text{Pb}(\text{NO}_3)_2$
Root growth slightly inhibited, mitotic activity unchanged, cell divisions usually undisturbed	1.0	5.0
Root growth strongly but reversibly inhibited, c-mitoses, t-tumors	2.5	15.0
Lethal action on most roots	3.0	20.0

Further studies on the lead localization in root tissues were carried out by autoradiographic method (with ^{210}Pb). Considerably higher lead content was found in root cell walls after administering $15 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ in nutrient than after $2.5 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ in deionized water (Figs. 7-10).

These observations were confirmed by the studies on lead localization in ultrastructures. There were much more lead deposits both on the root surface (Figs. 11, 12) and in cell walls (Figs. 13, 14) in nutrient than in water.

The present results show that different lead content in roots after adding lead to nutrient and water is caused by the accumulation of lead in root apoplast. Lead content in apoplast depends directly on its concentration outside a root. This is a non-toxic lead pool for root cells (Wierzbicka 1984). The lead that has entered root symplast is the only lead which is toxic. Its amount depends on the presence of antagonistic ions in the surrounding.

Thus, the differences in the analysed compound activity, depending on the kind of water used in preparing the tested solution, are caused by different numbers of ions antagonistic to the compound analysed in the tested solution.

Figs. 2, 3. Lead detection in live roots by the rhodizonate method. $3\times$. Fig. 2 — Control root (C) and root incubated for 24 h in $15\text{ mg dm}^{-3}\text{ Pb}^{2+}$ from $\text{Pb}(\text{NO}_3)_2$ supplied in nutrient. Fig. 3 —

Root incubated for 24 h in $2.5\text{ mg dm}^{-3}\text{ Pb}^{2+}$ from $\text{Pb}(\text{NO}_3)_2$ supplied in water

Fig. 4. Root tip covered with an additional structure — "a hood" (arrows). After 24 h incubation in $2.5\text{ mg dm}^{-3}\text{ Pb}^{2+}$ from PbCl_2 and 48 h postincubation in water, the root was treated with sodium rhodizonate. The light coloured root fragment grew in the course of postincubation. The dark colour that made the hood more visible results from the lead presence in the incubation period. $3\times$

Fig. 5. Detached hood stained with acetic orcein. An unilayer structure, its walls do not undergo maceration in the course of standard preparation of macerated squashed slides stained with acetic orcein with 1N HCl (9:1). $10\times$

Fig. 6. Hood cell stained with acetic orcein. $250\times$

In the investigations when the *Allium* test was applied, such a relation may lead on to many misunderstandings, especially when various works are compared. Sometimes, the information concerning the water used for the solution seems too obvious for the author to include in his paper. Also evaluation of the toxicity level of environment contaminants may be wrong, as under the natural conditions these pollutants are never the only compounds plants have contact with. On the other hand, the sensitivity of the *Allium* test can increase when deionized water is used for tested solution. It is of big importance when the compound analysed is hardly soluble in water, or when little of it is available.

A SMALL AND A BIG CONTAINER PITFALL

A rough examination was made of how fast onion roots took up Pb^{2+} administered in deionized water in the form PbCl_2 or $\text{Pb}(\text{NO}_3)_2$. Concluding from a very sensitive colour reaction between lead and dithizone it was found out that lead had been taken up by onion roots totally or almost totally from the incubation containers as soon as after 6 to 12 hours. Testing ledakrine with the isotope method Antosiewicz (1987) had in her experiment a similar timing of the drug uptake by onion roots. This had not been expected especially as the amount of incubation liquid per one bulb seemed to be relatively big i.e. 320 cm^3 . Hence, we think that the apriori assumption on the excess amount of the compound analysed in the incubation liquids, which is often met elsewhere, is not always true. Roots of onion bulbs are capable of taking up the analysed compound fully from the solution in the first incubation hours. Moreover, it was observed that lead content in water solutions decreased only due to water aeration, which had been caused by the

PLATE II

Figs. 7, 8. Autoradiographs of ground meristematic tissue. The longitudinal section of the root tip, 24 h incubation in $2.5 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ from $\text{Pb}(\text{NO}_3)_2$ with $3.77 \text{ kBq cm}^{-3} \text{ }^{210}\text{Pb}$ supplied in water. 200x

Fig. 7 — Section at 3 mm of the root height. Fig. 8 — Section at 1 mm of the root height

Figs. 9, 10. Autoradiographs of ground meristematic tissue. The longitudinal section of the root tip. Lead isotope labels along cell walls, 24 h incubation in $15 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ from $\text{Pb}(\text{NO}_3)_2$ with $3.77 \text{ kBq cm}^{-3} \text{ }^{210}\text{Pb}$ supplied in nutrient. 200x. Fig. 9 — Section at 3 mm of the root height.

Fig. 10 — Section at 1 mm of the root height

PLATE III

Figs. 11-14. Electrographs of 24 h lead salt treated roots. RS — root surface, CW — cell wall, Pb — lead deposits. Fig. 11 — Small lead deposits in and on surface wall of cap cell, incubation in $2.5 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ from PbCl_2 supplied in water. 27 000x. Fig. 12 — Large lead deposits in and on surface wall of cap cell, incubation in $15 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ from $\text{Pb}(\text{NO}_3)_2$ supplied in nutrient. 21 000x. Fig. 13 — Small lead deposits in walls of ground meristematic cells, incubation in $2.5 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ from $\text{Pb}(\text{NO}_3)_2$ supplied in water. 34 000x. Fig. 14 — Large lead deposits in walls of ground meristematic cells, incubation in $15 \text{ mg dm}^{-3} \text{ Pb}^{2+}$ from PbCl_2 supplied in nutrient.

27 000x

precipitation of insoluble lead carbonate. This should be taken into account when the components of solutions analysed can give insoluble compounds with carbonate acid.

If during onion root treatment the analysed compound is totally taken up from incubation liquid, it may cause considerable differences between reactions among roots of individual bulbs. This was illustrated in Wierzbicka's work (1986) by the ultrastructural root reaction to the same lead concentration. The bigger amount of the root fresh mass in one bulb the faster the concentration of the analysed compound decreases in the incubation liquid, what results in big differences in root reaction of particular bulbs during the experiment. In other words the amount of the analysed compound per one root fresh mass unit determines the root reaction to the analysed compound.

Practically then, the volume of an incubation container is of the first importance. Many results in our experiments seemed totally inexplicable before this simple relation was found. Yet having this relation in mind, this problem can be easily solved by using always the same amount of analysed liquid per one bulb.

However, it is much more difficult to overtake the other obstacle i.e. the difference in root fresh mass between individual bulbs. Thus, it was attempted to obtain root systems of a similar size (Wierzbicka 1987c). Due to the preliminary onion bulb preparation, the differences between particular root systems were considerably diminished. Nevertheless, these differences still do occur, especially between consecutive experimental series (Table 3). Eventually, total root mass per one onion ($3\text{ cm} \pm 1\text{ cm}$ length of root) ranged from 414 to 2000 mg. In order to decrease such big differences each bulb was left with only 20 roots, each 3 cm long. It enabled diminishing the extremes of root mass (Table 3), yet not to a required extent. The main reason for this fact lay in the 2.5 fold differences in the root breadth. Eventually the experimentators decided that leaving the same number of roots of a bulb was not enough to overcome the problems resulting from the differences in the weight of individual root systems. As cutting roots off is not unharmed to plants, this procedure was discontinued in further experiments. Therefore in practice the following procedure is recommended:

1. Before the test (when onion roots are 3 cm long) select the bulbs with roots of similar breadth, length and number. Bulbs with the thinnest and the thickest roots should not be considered.

2. Increase the number of onion bulbs in one experiment treatment to e.g. 8-10.

3. In the course of the experiment observe macroscopically the growth and appearance of the roots, individually for each bulb. The darkening of root tips, root transparency and lack of root turgor always indicate lethal action of the analysed compound. Measuring the longest root of each bulb every 12 or 24 h enable evaluating the inhibition degree of root growth. These data are very

valuable when the results of consecutive series are compared and already in the course of the experiment they usually enable estimating the size differences in reaction to the analysed compound between individual bulbs.

All the problems described or mentioned in this chapter occur only when the whole of the analysed compound is taken up by the onion roots before the incubation time is over. To exclude such a situation at least rough studies on

Table 3
Variation in root weight of individual bulbs applied as the initial material for the *Allium* test (mean values of 10 bulbs)

Experiment	Mean weight of all roots of a bulb (mg)	Mean weight of 20 roots (mg)
1	620 (414-1010)	452 (348-706)
2	469 (284-711)	339 (216-571)
3	1383 (900-2000)	
4	1416 (807-1682)	

the dynamics of the uptake of the analysed compound by onion roots under given experimental conditions are necessary. It would be advisable to make experiments with the *Allium* test in the continuous flow of incubation liquids, what enables keeping the concentration of the analysed compound at a fixed level.

VARIETY IN CONTROL MATERIAL

In the studies with any test, it is very important to know the variation range within the control material. In the analyses with the *Allium* test in conditions favourable for the root growth big variation within the control material was observed regardless of the fact whether nutrient or deionized water were used. E.g. Fig. 15, which illustrates this phenomenon, shows all individual results obtained in the analysis of mitotic activity during 96 h of root growth. Similar differences were observed between individual tips in autoradiographic studies of the synthesis levels of nucleic acids and protein in ^3H -thymidine, ^3H -leucine and ^3H -uridine treated roots (Fig. 16).

All the presented graphs depict the analysis of one or two tips from one bulb out of three to seven in one experimental time. Thus, the question arises whether the observed variation results from the differences between roots of different bulbs or between roots of one bulb only. In order to answer it 8 bulbs, each with about 20 roots of the same length (about 3 cm) left were selected. Next, five bulbs were grown for 72 h whereas the remaining three — for 84 h. Then the length of the roots was measured, squashed slides were made and mitotic index was calculated for all the roots.

Fig. 15. Mitotic activity in control roots grown in deionized water. The graph shows all individual results of mitotic activity of one root tip each (●)

Fig. 16. Label number in autoradiographs in control roots grown in deionized water after being treated with ^3H -thymidine, ^3H -leucine, ^3H -uridine. The graph shows individual results of one root tip each (●) and arithmetic mean (x)

Figure 17 presents mitotic activity in all the roots of individual bulbs. It illustrates both big differences between roots of individual bulbs and between single roots of one bulb. Bulb *c* was statistically different (verified with t-Student test at $\alpha = 0.05$) from bulbs *a* and *b*, whereas bulb *h* – from bulbs *d-g*. No correlation between root length and mitotic index value ($r = 0.16$) was found. It confirms earlier results (Zielewicz 1971) that the value of mitotic index stabilizes in onion roots at 2-4 cm length and remains constant in the course of further growth.

Fig. 17. Mitotic activity in all onion bulb roots. The bulbs were initially left with around 20 roots, 3 cm long each, then onions *a-e* were grown in deionized water for 72 h and onions *f-h* for 84 h. The graph shows individual results of mitotic activity of one root tip each (●) and arithmetic mean (x)

The experiment indicates that high diversity in the *Allium cepa* test material results from differences among individual roots of one bulb as well as from those among roots of individual bulbs. But these differences are not associated with different length of particular roots. The variation presented should be considered as extremely high possibly because of relatively long root growth in deionized water. But even regardless of the handicaps of the experimental method used here, this variation is too high anyway. Although the roots look similar they seem to differ physiologically and functionally.

Thorough analysis of root morphology showed that tips of some of them were additionally covered with one-layered "hood" (Figs. 4,5). Its cells are live and closely packed (Fig. 6). It occurs independently of the root cap and is only loosely attached to the root tip which is why it detaches easily in the course of fixation. Perhaps this is the reason why it has not been described so far. Although the role of "the hood" is not yet known we think it worth mentioning as it may be an anatomical reflection of the root variation within one onion bulb.

SUGGESTED MODEL OF THE EXPERIMENT

The variation described in control and experimental material taken into account, the experiment may be carried out in two different ways. The first one lies in the limitation of root number in the experiment and individual treatment of each bulb, the other one contrarily — in the increase in root number analysed in each experimental group, then considered (as it has been done as far, Wierzbicka 1988a,b) as a composed sample.

In our opinion the choice of either way should enable the experimenter to make use of the essential advantage of the *Allium* test, which lies in the opportunity to grow 40 genetically identical roots from one bulb. Therefore, the recommended procedure is as follows:

Each bulb is considered individually. In 3-5 replicates of the experiment, after given incubation period 5 roots from each bulb are randomly chosen for further analysis. For example: 3-5 control bulbs and 3-5 experimental bulbs (i.e. treated with one concentration of the analysed compound). After 6, 12 and 24 hours, 5 roots are cut off from each bulb. Next, each experimental bulb is transferred for 24-48 h to the liquid devoid of the analysed compound to check the recovery ability. After 24-48 h 5 roots per each bulb are collected.

Individual treatment of each onion bulb is the best experimental procedure which through the statistical analysis enables illustrating most thoroughly the effect of the analysed compound on *Allium cepa* L. cell division in root tips. The disadvantage of such a technique lies in the limited number of experimental points, as only 40 roots per each bulb can be used at maximum (Wierzbicka 1987c). Moreover, cutting a root off in the course of the experiment is not indifferent to remaining roots. That is why it is so crucial that a similar number of roots were cut off control and experimental bulbs, what in consequence means that the number of experimental and control bulbs should be the same.

Summing up: what is the answer to the question whether the variation within the *Allium* test material disqualifies the application of onion roots in test studies? In our opinion the *Allium* test is very good while the variation described is typical of any biological material. We think all researches working experimentally have in some time encountered such a high variation range in

their studies. Only by thorough analysis of test material, appropriate experimental procedure and statistical analysis of the results can we generalize with a low error risk.

Acknowledgment

The experiment was financed by Centralny Problem Badań Podstawowych (CPBP) No 05.02.

REFERENCES

- Antosiewicz D., 1984. Binding of 1-nitro-9-(3-dimethylaminopropyl-amino)-acridine to the DNA of the apical meristem cells of adventitious onion (*Allium cepa* L.) roots. Acta Soc. Bot. Pol. 53: 217-222.
- Antosiewicz D., 1987. An evaluation of the suitability of the onion test for studying the biological activity of potential antitumor drugs on the example of ledakrin. Acta Soc. Bot. Pol. 56: 543-555.
- Antosiewicz D., 1988. An analysis of the cell cycle in *Allium cepa* L. root meristem under the influence of ledakrin. Folia Histochem. Cytobiol. (in press).
- Błok N., 1955. Jakościowa analiza chemiczna. PWN, Warszawa.
- Degrassi F., Rizzoni M., 1982. Micronucleus test in *Vicia faba* root tips to detect mutagen damage in fresh-water pollution. Mutat. Res. 97: 19-33.
- Fiskesjö G., 1981. *Allium* test on copper in drinking water. Vatten 37: 232-240.
- Fiskesjö G., 1982. Evaluation of short-term for toxicity and mutagenicity with special reference to mercury and selenium. Institute of Genetics, University of Lund, Sweden.
- Grant W. F., 1978. Chromosome aberrations in plants as a monitoring system. Envir. Hlth. Persp. 27: 37-43.
- Grant W. F., 1982. Chromosome aberration assays in *Allium*. Mutat. Res. 99: 273-291.
- Grant W. F., Zura K. D., 1982. Plants as sensitive *in situ* detectors of atmospheric mutagens. In: Mutagenicity: new horizons in genetic toxicology. Heddle J. A. (ed.). Academic Press, New York, pp. 407-433.
- Mäkinen Y., 1963. The mitotic cycle in *Allium cepa* with special reference to the diurnal periodicity and to the seedling aberrations. Ann. Bot. Soc. "Vanamo" 34: 3-60.
- Menn J. J., Still G. G., 1977. Metabolism of insecticides and herbicides in higher plants. CRC Crit. Rev. Toxic. 1: 1-21.
- Nilan R. A., 1978. Potential of plant genetic system for monitoring and screening mutagens. Envir. Hlth. Persp. 27: 181-196.
- Peterson C. A., Peterson R. L., Robards A. W., 1978. A correlated histochemical and ultrastructural study of the epidermis and hypodermis of onion roots. Protoplasma 96: 1-21.
- Plewa M. J., 1978. Activation of chemicals into mutagens by green plants: a preliminary discussion. Envir. Hlth. Persp. 27: 45-50.
- Scott B. R., Sparrow A. H., Schwemmer S. S., Schairer L. A., 1978. Plant metabolic activation of 1,2-dibromoethane (EDB) to a mutagen of greater potency. Mutat. Res. 49: 203-212.
- Serres F. J., 1978. Introduction utilization of higher plant systems as monitors of environmental mutagens. Envir. Hlth. Persp. 27: 3-6.
- Spanó M. A., Takahashi C. S., 1981. Genetic effects of nifurtimox on *Drosophila melanogaster*, wistar rats and *Allium cepa* L. Rev. Brasil Genet. 3: 357-366.
- Wierzbicka M., 1984. Ultrastructural location of lead in the cell walls of *Allium cepa* L. roots. Post. Biol. Kom. 11: 509-512.

- Wierzbicka M., 1986. The effect of lead on the ultrastructure changes in the root tip of onion — *Allium cepa* L. *Folia Histochem. Cytobiol.* 24: 340-341.
- Wierzbicka M., 1987a. Lead accumulation on its translocation barriers in roots of *Allium cepa* L. — autoradiographic and ultrastructural studies. *Plant Cell Environ.* 10: 17-26.
- Wierzbicka M., 1987b. Lead translocation and localization in *Allium cepa* L. roots. *Can. J. Bot.* 63: 1857-1860.
- Wierzbicka M., 1987c. An improved method of preparing onion bulbs for the *Allium* test. *Acta Soc. Bot. Pol.* 56: 43-53.
- Wierzbicka M., 1988a. Mitotic disturbances induced by low doses of inorganic lead. *Caryologia* (in press).
- Wierzbicka M., 1988b. Disturbances in cytokinesis caused by inorganic lead. *Env. Exp. Botany* (in press).
- Zielewicz E., 1971. The course of mitotic activation of *Allium cepa* L. adventitious roots. M. A. Thesis, Zakład Anatomii i Cytologii Roślin, Uniwersytet Warszawski (in Polish).

Niektóre problemy testu Allium

Streszczenie

W pracy omówiono niektóre problemy związane z prowadzeniem doświadczeń z wykorzystaniem testu *Allium*, oraz przedstawiono różnorodność materiału testowego. Zaproponowano, aby podczas standardowych prac z zastosowaniem testu *Allium* prowadzić hodowlę cebul w wodzie dejonizowanej lub na pożywce Hoaglanda. Wykazano, na przykładzie związków ołowiu, iż rodzaj wody użytej do przygotowania roztworu ma istotny wpływ na toksyczność badanego związku i może być przyczyną nieporównywalności wyników uzyskanych przez różnych badaczy. W pracy przeanalizowano różnice w metabolizmie poszczególnych korzeni i to zarówno w materiale kontrolnym, jak i w doświadczalnym. Ostatecznie zaproponowano, aby w doświadczeniach testowych każdą cebulę traktować indywidualnie. Przedstawione podejście w dużym stopniu zwiększa dokładność i powtarzalność wyników w teście *Allium*.