POLSKIE TOWARZYSTWO BOTANICZNE

MONOGRAPHIAE
BOTANICAE

Vol. 67, 1985

Redaktor: *R. Olaczek*

WARSZAWA

Wydano z pomocą finansową
Polskiej Akademii Nauk

KOMITET REDAKCYJNY

R. Olaczek (Łódź), E. Sidorkiewicz (Warszawa)

RADA REDAKCYJNA

W. Matuszkiewicz (Warszawa), Z. Podbielkowski (Warszawa), E. Sidorkiewicz (Warszawa),
R. Olaczek (Łódź), H. Piotrowska (Gdańsk), K. Rostański (Katowice), W. Żukowski (Poznań)

ADRES REDAKCJI: Al. Ujazdowskie 4, 00-478 Warszawa

© Copyright by Państwowe Wydawnictwo Naukowe, Warszawa 1985

ISBN 83-01-04485-3
ISSN 0077-0655

SPIS TREŚCI

J. Misiewicz: Investigations on the synanthropic flora of Polish sea harbours — Badania nad florą synantropijną polskich portów morskich	5
H. Tomaszewicz, S. Kłosowski: Roślinność wodna i szuwarowa jezior Pojezierza Sejneńskiego — Aquatic and rush vegetation of the Sejny Lakeland lakes	69

INVESTIGATIONS ON THE SYNANTHROPIC FLORA OF POLISH SEA HARBOURS

Badania nad florą synantropijną polskich portów morskich

JÓZEF MISIEWICZ

Botanical Department, Institute of Biology
Pedagogical School, Słupsk

1. ABSTRACT

The results of studies on the synanthropic flora of all Polish sea harbours are reported. The first chapters describe in outline the physiographic conditions, trade contacts and the history of research on the synanthropic flora of Polish sea harbours. The detailed part (chapter 6) gives an analysis of the synanthropic flora with reference to its state in the particular harbours, its biological durability, the proportion of apophytes and anthropophytes, the origin of anthropophytes, the flora of the particular harbour habitats, the state of the synanthropic flora of harbours as compared with the flora of some chosen areas. In the end chapter the results are summarised and conclusions are drawn.

2. INTRODUCTION

Sea harbours are places where human activity is very diversified and intensive. Here are the crossroads of water and land routes, and home and foreign goods from other countries and continents are trans-shipped. Seeds and fruits of numerous plants are brought here to the grain elevators, warehouses, rice pearling mills, oil mills etc. Numerous diaspores are also carried involuntarily particularly with mass goods. In old times many foreign plant species arrived with balast earth unloaded in the ports from sailing ships carrying goods. In this way the activity of ports created and still

creates favourable conditions for the intrusion of very numerous diaspores of various plant species.

As the result of intensive human activity there arise in harbours diversified habitats, some of which are not found in other regions, such as for instance breakwaters, and the frequent destruction of the vegetal cover exposes wide spaces on which the newcomers can settle.

The synanthropic vegetation arising in this way develops usually without human intention and the harbour areas become very good gates for penetration of many new plant species which thrive here and even many a time exhibit a tendency to spreading. A better knowledge of the synanthropic flora of harbours, particularly of its composition, migration routes and rate as well as the factors decisive for its development, beside being of theoretical interest are of considerable practical importance. If only to mention the appropriate maintenance of stacking yards, port railway tracks, fixing of sandy waste land etc.

Investigations on sea harbour flora were conducted in Poland until recently only in the harbour areas of Gdańsk and Gdynia (Schwarz 1967) and partly in Szczecin (Ćwikliński 1970, 1974). Neither does the available foreign literature supply numerous elaborations concerning the flora of harbours (e.g. Lehman 1895, Blomquist 1917, Blom 1927, 1933, 1936, 1939, 1961, Saaristo-Taubert 1963). A somewhat different problem is the flora of river ports tackled among others by Hupke (1933, 1935), Runge (1956, 1965), Stricker (1960, 1961) and in Poland Szotkowski (1972), and Rostański and Szotkowski (1973).

The present work is the result of special research on the synanthropic flora of all Polish sea harbours. The chief aim of the research was to establish the size, composition and origin of the plant populations of ten ports with a different history and function and of different importance, situated on territories developing extremely fast and along various lines in connection with advancing industrialisation and urbanisation. Systematic investigations were undertaken, in the period 1971-1979 in the harbour of Gdańsk and 1973-1979 in Gdynia. Herbarium materials in the number of more than 5000 sheets are deposited in the Department of Botany of the Normal School in Słupsk.

In the course of preparation of this study I profited of the help and advice of many persons to which I wish to express my deep gratitude. Above all I am indebted to Prof. Dr habilit. Waldemar Żukowski for suggesting the theme and his interest in the course of the work, for making available to me some unique references and for checking the correctness of determination of some critical plant species. My thanks go also to Doc. Dr habilit. Krzysztof Rostański for his valuable advice during the search for *Oenothera* in the harbours of Gdynia, Gdańsk, Darłowo and Ustka in the years 1974 and 1975 and also for checking some of identifications of plant species including the collections of *Oenothera*. Particular thanks go to the earlier investigators of the flora of Polish sea harbours that

is Dr Zofia Schwarz and Doc. Dr habilit. Eugeniusz Ćwikliński for information on the distribution of rare species on harbour territories and for checking of the determinations of part of the species. I wish to thank Doc. Dr habilit. Ryszard Sowa for information concerning the list of ephemerophytes and dr Karol Latowski for his help in comparison of critical plant species with herbarium material in the Department of Plant Taxonomy of the A. Mickiewicz University in Poznań. I also thank my son Adam for his unswerving aid in field work, particularly in completion of the herbarium.

I am also indebted to the Direction of the marine boards in Szczecin, Słupsk and Gdynia and the Direction of the relevant ports for making the studies possible, supplying information concerning the lay-out of the port territory, economic activity etc.

In the period 1975-1979 the work was supported by a grant of the Committee on Botany of the Polish Academy of Sciences for which I am grateful.

3. OUTLINE OF PHYSIOGRAPHIC CONDITIONS

3.1. TERRITORY INVESTIGATED

The studies were extended to the trade and fishery harbours on the Polish sea coast (Fig. 1).

Fig. 1. Situation of Polish sea-harbours against background of geobotanical regions: 1 – Brzeg Bałtyku; 2 – Pobrzeże Bałtyckie; 3 – Nizina Szczecińska; 4 – Żuławy Wisły; 5 – Pojezierze Pomorskie

The term harbour refers to the area administered by the marine board at the time of the investigations. The harbour complex Szczecin-Świnoujście is treated as two separate harbours, mainly on account of the large distance separating them (65 km).

In some cases the investigations were extended to territory adjacent to

the harbours: in Szczecin the area of the railway Port Centralny station, in Świnoujście the railway premises with the station Świnoujście-Odra and the region of the ferry landing place, in Ustka the square adjacent on the NW-side used sometimes as a temporary storage yard, in Łeba the area neighbouring with the harbour channel, in Władysławowo the railway premises with the station Władysławowo-Port, in Hel the area of the railway branch line to the port and the vegetation belt developing along the fence surrounding the port on the northern side, in Gdańsk the whole island (Ostrów) used at present by the repair docks and the sites with species reported in the literature (Schwarz 1967) from areas which presently are not included in the harbour as for instance Brzeźno, partly Składy, Młyniska, Przeróbka, Wisłoujście and others.

The investigations were conducted above all on land anthropogenic habitats, less frequently on seminatural ones which may still be found in harbours, such as fragments of dunes, coastal coniferous forest etc. In view of the necessity of applying different methods to aquatic habitats, rivers and channels were excluded, the study was limited there to recording of species occurring on the banks.

In the characteristic of the harbours only concise information is given concerning their situation, surface area, economic role and trade contacts. More extensive information on the history of these ports, the successive steps of their development, facilities on the particular quays, development of trade relations etc. may be found in special publications („Port Gdański” 1929, „Port Gdynia” 1936, Grodek 1948, Kowalenko 1950, Brocki 1959, Andruszkiewicz 1960, Mikołajski 1960, Andruszkiewicz et al. 1961, Heliński 1962, Szopowski 1962, Lesiński 1965, Koselnik 1965, Zaremba and Orlińska 1965, Magiera 1966, Hoffa 1968, „Locja Bałtyku, Część południowa. Wybrzeże polskie” 1973, Dziędziul 1975).

Szczecin, owing to its situation, is at the same time a sea and river port. The surface area of the harbour is 1760 ha including 1215 ha of land. The total length of quays in 1970 was 17.5 km. The surface area of open storage yards was jointly in the port complex of Szczecin-Świnoujście in 1970 41.6 ha, including for mass goods 14.3 ha, timber 18.4 ha and general cargo 8.9 ha (Zaremba and Orlińska 1965, Dziędziul 1975). Not all port areas are managed, there are still many unprotected quays and places treated as waste areas. In the spatial lay-out of the port several regions with different functions can be distinguished such as the port for mass trans-shipment, for general cargo and the industrial port. The Szczecin port has regular relations with all major ports on the Baltic in northern and north-western Europe and in the Mediterranean region. It is, moreover, the home port of many ocean lines.

Świnoujście — lies at the outlet of the river Świna into the Pomeranian

bay. The area of the port includes an about 4.5-km segment of the river with the terrains surrounding it and the water spaces on both sides. The total surface area of the storage yards in 1967 was about 6 ha, and the length of the quays 2487 m (Dziedziul 1975). The Świnoujście port is being intensively extended and is becoming a transshipping base in the first place of mass goods for high tonnage ships. The port is also an intervention port for Szczecin where high draft ships with Szczecin as destination are unloaded and those going out to sea from Szczecin are fully loaded. Moreover, Świnoujście is an important high sea fishery base.

Kołobrzeg — an industrial, trade and also fishery harbour situated at the mouth of the river Parsęta. Its surface area is 10.94 ha (Hoffa 1968). The port has grain elevators, general cargo storehouses and a fish preserving plant.

Darłowo — a fishing and industrial port, situated at the mouth of the river Wieprza. It has large grain elevators, general cargo storehouses and a fish preserving plant. In the immediate neighbourhood on the southern side there is an open space for timber storage.

Ustka — an industrial, trade and also fishery harbour at the mouth of the river Słupia. It has grain elevators, general cargo storehouses and a fish preserving plant.

Łeba — a small fishery harbour at the mouth of the river Łeba without any grain elevators or railway facilities.

Władysławowo — a large fishery port at the base of the Hel Peninsula, built in the interwar period. It is connected by branch railway lines with the general railway network and neighbours with the railway station.

Hel — a fishery and industrial harbour situated at the eastern end of the Hel Peninsula.

Gdynia — is one of largest, youngest and most modern ports on the Baltic. It was built in the period 1922-1935 („Port Gdynia” 1936). Basic data concerning its state at the end of 1960 may be found in the publication of Andruszkiewicz et al. (1961). The total surface area of the port is 854 ha, the length of the quays about 16 km, the length of the breakwaters 2928 m, the length of the port railway tracks 204 km, the general surface area of the storage yards 21.65 ha. The port includes grain elevators, an oil mill, a rice pearling mill, many storehouses and storage yards. The Gdynia port has regular relations with most great ports of the world. Transshipping includes mass goods (coal, coke, ores, apatites, sulphur), agricultural products, mineral raw materials, semiproducts and industrial products. The port is also the site of trans-shipment of goods in sea container turnover.

Gdańsk — is a typical natural port of sea and river character connected with the network of the Polish waterways of the Vistula catchment. The

surface area of the port is 1010 ha, the land part comprising 740 ha. The quay length is about 13 km, those exploited 8.2 km („Locja Bałtyku” 1973). The length of the railway tracks in the port is 142 km and the storage yards occupy about 200 ha („Port Gdański” 1929). The port has regular relations with most great ports of the world. Trans-shipments include mass goods, industrial and agricultural products, mineral raw materials and semiproducts.

3.2. CLIMATE

The climate of the southern Baltic coast on which the ports in point lie has many individual characters, therefore, it creates different conditions for plant development and affects particularly the course of phenological phenomena.

As base for characterising the climate of the sea coast served meteorological data from eight localities (no data from Gdynia and Władysławowo). According to Romer's classification (1949) based on statistical calculations of 11 climate elements, Szczecin lies within the climatic range of the Central Polish Lowland zone, whereas all the remaining harbours lie within the range of the Baltic climate.

The influence of the sea is strong within both climatic ranges, but a marked tendency towards continental climate is noticeable as we advance in eastern direction (Romer 1949, Prawdzic 1961, 1962).

As regards temperatures (Table 1) the climate of the coast does not show major differences: not very high annual temperatures (7.5° - 8.3° C) and small annual temperature amplitudes (17.7° - 19.2° C). A retardation of nearly all seasons is noticeable, the summer is cool, the autumn long and mild and the winter not very severe. The warmest part of the coast is the Szczecin-Świnoujście region where the mean annual temperature is 8.3° C (Świnoujście 8.1° C). Winter starts here latest in Poland so that plant vegetation lasts sometimes to January. The higher values of the temperature amplitudes in Szczecin and Gdańsk as compared with other localities (Table 1) indicate a smaller influence of the Baltic on account of their situation.

The remaining data concerning thermic relations show a similar course, namely, the number of days with high frost and frost or ground frost and of hot and very hot days increases in eastern direction (Romer 1949, Mońiak 1961, Prawdzic 1961). These phenomena indicate a declining influence of oceanic, with a simultaneous increase of the effect of continental climate.

Precipitation on the sea coast (Table 2) does not show such a regularity as do thermic conditions (Wiszniewski 1953). The mean annual sum of precipitation varies from 528 in Hel to 667 in Ustka. The precipitation

Table I
Mean air temperature for the period 1881-1930 (Kołobrzeg, Ustka, Łeba and Hel — mean for period 1948-1959)

Meteorological station	A.s.l.	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII	Year
Szczecin	26	-0.9	-0.1	3.0	7.5	12.9	16.2	18.3	16.9	13.6	8.5	3.5	0.5	8.3
Świnoujście	6	-0.6	-0.1	2.5	6.6	11.4	15.2	17.4	16.7	13.8	8.7	3.9	0.9	8.1
Kołobrzeg	4	-0.4	0.9	2.0	6.0	10.3	14.2	16.8	16.4	13.6	9.1	4.2	1.7	7.8
Darłowo	6	-1.1	-0.7	1.5	5.5	10.2	13.9	16.6	16.1	13.2	8.3	3.7	0.0	7.5
Ustka	6	-0.5	-1.0	1.3	5.6	10.1	14.2	16.7	16.6	13.3	8.9	4.3	1.7	7.6
Łeba	3	-0.6	-1.1	1.2	5.5	9.9	14.0	16.6	16.6	13.3	8.9	4.2	1.6	7.5
Hel	7	-0.6	-0.5	1.4	5.1	9.9	14.1	17.2	16.8	14.0	9.2	4.3	1.0	7.7
Gdańsk — Nowy Port	5	-1.6	-1.1	1.6	6.1	10.9	14.8	17.5	16.6	13.5	8.1	3.2	-0.1	7.5

Table 2

Mean sums of precipitation for the period 1891-1930 (Ustka 1948-1959) (Wiszniewski 1953, Prawdzic 1962)

Meteorological station	A.s.l.	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII	Year
Szczecin	20	44	31	36	37	44	52	77	64	46	43	39	48	561
Swinoujście	3	49	35	39	43	47	55	78	76	51	54	46	53	628
Kołobrzeg	2	41	29	36	37	43	52	70	75	58	53	44	49	587
Darłowo	6	42	30	36	37	41	54	65	75	69	64	59	55	627
Ustka	6	58	42	31	40	37	50	94	73	88	56	47	51	667
Łeba	3	43	29	30	35	41	49	67	76	75	64	56	54	619
Hel	7	30	24	28	34	41	51	64	74	56	43	43	40	528
Gdańsk — Nowy Port	5	31	25	32	35	46	56	68	68	56	43	47	39	546

distribution in the course of the year is not very favourable — in autumn rainfall is much more abundant than in spring and the amount in the vegetation period is not high. The cause of these wide differences in the amount of precipitation in the particular seasons is the distribution of winds with prevalence of the western ones.

3.3. HABITAT CONDITIONS IN HARBOUR AREAS

The situation of most ports around the mouths of rivers decides of some characters of the habitats and is conditioned by natural factors such as the substrate, the water regime and climate. The primary habitat on which the harbour areas formed were usually swamps and peat bogs or dune sands. For instance the Szczecin harbour lies in a river valley lined with alluvial formations to a depth of 25 m. On sand and gravel lies a 5-9 m thick peat layer. In the course of construction of the port an earth layer of 4 m³ per 1 m² was spread over the peat (Mikołajski 1960). The harbours of Świnoujście, Kołobrzeg, Darłowo, Ustka, Łeba and most of the quays of the Gdańsk port arose on similar territory (Mieczysławski 1946). The harbours of Władysławowo and Hel, on the other hand, are built on dunes. The Gdynia port is also artificial, it arose, however, on a partly marshy site where the port docks were dug out („Port Gdynia” 1936).

The construction of the Gdańsk Northern Port was conducted under different conditions (Siuta 1973a, b). Here, within the harbour area there are soils silted up from the sea bottom with mechanical composition of loose sand, thus very poor in assimilable nutrient components. Moreover, there are soils which developed from dune sands, among them slightly clayey sands passing into shallow loose sand. On small areas muck soil may be found. The above mentioned outset soils were recultivated by adding 100 m³/ha (an about 10-cm layer) of sewage deposits from the city of Gdańsk, slag, waste betonite and organic-mineral silt from the area on which the refinery in Przejazdów was built, containing about 20 per cent of organic matter.

Wide spaces of the harbour area were covered with material which makes plant development difficult (slag, gravel, sand, crushed stone, sett, cut stone, concrete, sometimes sulphur and coal dust or coarser pieces of coal). Larger areas covered with foreign material are classified to artificial habitats, in contrast to natural and seminatural ones developed on local formations (Kornaś 1972, Ćwikliński 1974).

Beside the different origin, mechanical and chemical composition of the substrate, there are many other factors decisive for plant development such as: trampling, traffic, wide temperature variations (especially on stony or concrete substrate) illumination and humidity conditions lubricant dropping,

fertilisation with dust of transloaded mineral fertilisers etc. Moreover, mechanical and chemical weed control, mainly on railway tracks, less frequently in storage yards plays an important role.

Thus, the habitats of synanthropic plants in the studied harbours are greatly differentiated as regards mechanical composition, edaphic conditions, humidity, angle of slope, thermic and light conditions and, above all, human activity. The vegetal cover is frequently destroyed by excavations, reconstruction of roads, removal or adding of earth, liquidation of old and laying out of new storage yards, levelling of dumping grounds, moving and removal of the vegetation etc. Under these conditions classification of habitats in harbours is rather difficult, above all owing to the lack of a distinct boundary between them and the frequent changes in their utilisation in connection with intensive development. Therefore, the kind of habitats distinguished and the data on their flora (Table 14) are only of general character. A short characteristic of these habitats is given below.

1. Storage yards – are mostly artificial habitats. Their surface is covered with concrete, slag, gravel or stone chippings. Sometimes the substrate consists of dune sand or natural soil.

2. Railway premises – the railway tracks are artificial habitats, but on the slopes of embankements seminatural ones may be found. In the Łeba harbour there are no railway premises.

3. Wasteland – is the most diversified group of harbour habitats. Included here are unmanaged areas, dumping grounds, building grounds, belts of soil along enclosures, around cottages, along roads etc.

4. Edges of docks, channels and segments of rivers – within the harbour territory have various slopes and mostly have a concrete or cut stone revetement, less frequently they are protected with brushwood.

5. Breakwaters – are the most typical for harbours artificial habitat with a specific combination of ecological factors. The material consists of concrete and cut stone. The joint length of the breakwaters in all the examined harbours is about 8 km. These habitats are exposed to very intensive trampling, spraying and inundation with sea water contaminated with lubricants.

They are fertilised by bird excrements and exhibit wide temperature variations.

6. Streets, squares, parkings, paths – are artificial habitats paved with sett, covered with asphalt, concrete, slag or gravel.

7. Lawns and flower beds, small gardens – belong to artificial habitats. The soil has mostly been brought from elsewhere. In recent years the surface area of these habitats increases rapidly, mainly owing to liquidation of wasteland. The problems connected with introducing greenery into the Northern Port in Gdańsk are discussed by Siuta (1974).

8. Young forest plantations – are seminatural or artificial habitats. They are relicts of what was once riverside brushwood and fragments of coastal

coniferous forest, and so is the scrub along fences, walls, railway premises, storage yards etc.

9. Dunes and sandy wastes in their region – belong to natural or seminatural habitats. When enriched in foreign components as for example coal dust they should be considered as artificial. They usually remain as a protective zone, sometimes, however, they constitute the main harbour area (e.g. Władysławowo, Hel, Northern Port in Gdańsk), less frequently they are utilised as storage places for various materials.

4. TRADE RELATIONS OF POLISH SEA HARBOURS

The sea harbours, on account of their convenient situation at the mouth of rivers, have long been important economic centres in which handicraft flourished, and above all sea and land trade developed (Kowalenko 1950).

The increasing economic requirements caused the harbours to extend continuously from immemorial times, but a more intensive development is noted as late as the 14th century. The trade contacts with the ports of western and northern Europe become more and more lively (Andruszkiewicz 1950, Zbierski 1962). This already organised exchange of goods in which cities belonging to the Hansa play an ever more important role leads to an intensive development of the shipping trade and most of the existing harbours on the Baltic sea. Some of the latter, as for instance Świnoujście, Władysławowo and Gdynia were founded much later and in the remaining ports new quays were added for exploitation at various periods (Szopowski 1962, Dziedziul 1975).

The trade turnover varied in dependence on the economic needs. For long periods the type of exchange was established according to the agricultural and forest character of the Pomerania province and the Vistula and Odra catchments. Important items in export were above all grain and timber, also starch, flour, oils and sugar. The import consisted chiefly of various ores, cotton, southern fruits, rice, herrings, and industrial products (Andruszkiewicz 1960).

A characteristic feature of the trade harbours was in general the prevalence of export over import (Andruszkiewicz 1960). Owing to this, many ships arrived empty or loaded with ballast earth which was unloaded in the port in specially marked out areas. These were the site of mass appearance of new species, particularly ephemerophytes (Schwarz 1967). An exception was the port of Szczecin where import prevailed over export. The main bulk of goods consisted of mineral coal brought from England, iron ore and stone from Sweden. Besides, cellulose was imported, soy bean and linseed (Heliński 1962, Dziedziul 1975).

Up to World War I trade was carried on mainly with European

countries lying on the Baltic and North Sea coasts. Trials were undertaken to start ocean lines without the intermediary of west European ports, for instance in mid 19th century raw material was imported for the Szczecin refinery directly from South America and India (Dziedziul 1975).

In the interwar period regular ocean lines were established with many countries on various continents, owing to the construction of a modern port

Table 3

Cargo in Polish harbours in 1000 tons (Rocznik statystyczny 1978)

Cargo groups	Years		
	1975	1976	1977
Total	54 046	60 775	63 773
Coal and coke	24 096	25 184	25 774
Ore	5 511	5 126	6 294
Cereals	4 118	6 182	5 634
Timber	894	943	871
Rock-oil	4 394	7 054	7 993
Other goods	7 200	7 269	7 999
General cargo	7 833	9 017	9 238
Gdańsk:	18 554	22 939	25 188
Coal and coke	9 417	10 439	10 775
Ore	1 641	1 255	1 344
Cereals	929	1 392	1 004
Timber	388	478	451
Rock-oil	1 394	3 749	5 337
Other goods	2 911	3 512	4 249
General cargo	1 879	2 114	2 028
Gdynia:	12 750	13 206	13 454
Coal and coke	3 722	3 258	3 322
Ore	1 024	863	1 404
Cereals	2 036	2 772	2 699
Timber	1	—	—
Rock-oil	1 856	1 924	1 572
Other goods	698	586	617
General cargo	3 413	3 803	3 840
Szczecin:	22 481	24 259	24 661
Coal and coke	10 957	11 487	11 647
Ore	2 846	3 008	3 546
Cereals	1 076	1 904	1 742
Timber	482	447	400
Rock-oil	1 144	1 381	1 084
Other goods	3 559	3 118	3 093
General cargo	2 417	2 914	3 149

in Gdynia. The trade contacts of other ports were also extended, particularly of Gdańsk and Szczecin (Brocki 1959, Andruszkiewicz 1960, Magiera 1966).

After World War II the trade turnover in the Polish sea ports increased consistently, their characteristic feature remaining the predominance of export over import. The main goods included in export are coal, coke, timber and sulphur and in import various ores, southern fruits, grain and general cargo. Transit shipments constitute a large proportion (Andruszkiewicz 1960) (Table 3).

In the postwar period it is the large trade ports that develop most intensively (Gdynia, Gdańsk, Szczecin-Świnoujście), many regular ocean lines arise, particularly from the Gdynia port to America, Africa, Asia and Australia. A number of ocean lines were also established from Gdańsk and the port complex Szczecin-Świnoujście (Andruszkiewicz 1960). The remaining harbours, in view of their small turnover, modest facilities and less extensive range of activity serve mainly as fishery and trade ports. Among the smaller ports only Kołobrzeg develops a wider trade activity (Rocznik Statystyczny 1978).

The sea ports have good railway connections with the inland by railway lines, motor roads and some also by inland waterways.

The here presented trade contacts of Polish sea harbours with other countries of the world influence the synanthropic flora as regards the species composition and the geographical character on the territory occupied by these ports. These problems will be considered in the further part of this paper.

5. HISTORY OF THE RESEARCH ON SYNANTHROPIC FLORA OF POLISH SEA HARBOURS

The synanthropic, and particularly adventitious flora of the harbours on the southern Baltic coast has long been an object of interest. To the best known cities from the floristic point of view belongs no doubt Gdańsk where investigations were carried out regularly from the beginning of the 19th century (Schwarz 1967). The synanthropic flora of the harbour area of this city aroused the interest of Klinsmann (1836, 1843, 1865), C. Klinggraeff (1848, 1854, 1866), Bail (1880), H. Klinggraeff (1880), Helm (1881), Lutzow (1892), Abromeit (1898-1940), Hilbert (1910), Preuss (1910, 1928), Herweg (1914), Wangerin (1916, 1921, 1925, 1930-1931, 1937), Kalkreuth (1941), Hansen (1959) and others. Valuable information concerning the synanthropic flora of the Gdańsk harbour is moreover, contained in numerous floristic notes and preserved old herbaries quoted by Schwarz (1967). The most penetrating studies of this flora, however,

involving the harbour area of Gdańsk and Gdynia were carried out in the period 1949-1965 by Schwarz (1961, 1967).

Beside the above named two harbours, Szczecin is also well known as regards floristics. The adventitious flora of this city including that of the harbour areas has been described by Holzfuss (1916, 1920, 1936, 1937, 1941). Numerous sites with synanthropic species recorded before 1945 were published by Scheuermann in 1956. After 1945 some synanthropic species have been dealt with by Czubiński (1950), Kownas (1958, 1959), Kownas and Piskorski (1958), Kownas and Sienicka (1962), Celiński (1964), Radomski and Jasnowska (1964, 1965) and Balcerkiewicz (1971). Most comprehensive information on the synanthropic flora of Szczecin has been supplied, however, owing to the special studies of Ćwikliński (1965, 1967, 1968, 1970, 1971a, b, 1974) who also mentions the sites of some species in the harbour areas. Investigations of the synanthropic flora of the remaining harbours were mainly connected with studies on the vegetal cover of Pomerania which belongs to the relatively well known regions in this respect (Czubiński 1950). Many valuable remarks concerning synanthropic species may be found in descriptions of the flora of this region (Schmidt 1840, Marsson 1869, Abromeit 1898-1940, Ascherson and Graebner 1898-1899, Müller 1904, Hegi 1908-1960). A list of bibliographic data on the flora of Pomerania, published up to 1925 is given by Leick (1926) and later by Czubiński (1950) and Sulma (1959).

Some information on the synanthropic species on the area of Świnoujście is supplied, beside the above mentioned flora records, by the papers of Piotrowska (1957, 1966b) and Urbański (1968). Many valuable observations concerning the distribution of ruderal species in Świnoujście, including the harbour area, may be found in the publication of Piotrowska (1966a) devoted to vascular plants of the Wolin and south-eastern part of Uznam islands. A more extensive characteristic of the synanthropic flora of Świnoujście is given in another elaboration (Misiewicz et al. 1979).

Up-to-date reports on the synanthropic flora of the harbours and cities of the middle part of the coast were very modest. Several mentions on the adventitious flora of these cities and harbour areas may be found in the papers of Schoeps (1931), Holzfuss (1933, 1936, 1937), Krauze (1934), and at a later period, Żukowski (1960, 1961), Nowiński (1962, 1965) and Żukowski and Piaszyk (1971). Fragmentary but important information on the flora of Kołobrzeg, Darłowo and Ustka is also supplied by the old herbarium prepared by the Słupsk florists in the period 1897-1943 (Misiewicz 1977c).

The harbours of Władysławowo and Hel have not been examined floristically until lately.

The present elaboration, the result of nine-years' investigations in the period 1971-1979, made it possible to gain a good knowledge of the present state of the synanthropic flora of all Polish sea harbours, and allowed at the same time to follow the changes in the quantitative and qualitative composition of this flora. Some data accumulated in the period 1971-1975 have already been published in earlier papers of the author (Misiewicz 1976, 1977a, 1978a, b). In an extensive publication of 1976 a floristic review is given comprising 1014 plant species with reference to: the geographical-historical group according to Kornaś (1968a), the frequency of their occurrence, their origin from natural communities (in the case of apophytes), their primary homeland (in the case of anthropophytes). A characteristic is also given of 22 associations and 8 ruderal communities of harbours on the basis of 307 phytosociological records.

In the period 1976-1979 and partly also in 1980 floristic studies were continued in harbours. Their result is the finding of new sites of many species in various harbours and the detection of 41 species new to synanthropic habitats. Thus, the list of the synanthropic flora of sea harbours has increased to 1055 species. Lack of space in the present elaboration does not allow a detailed index of the sites recorded after 1975. Therefore, only new species for the synanthropic flora of Polish sea harbours, detected in the period 1976-1979 are specified. These are among apophytes: 1) in Szczecin: *Equisetum maximum*, *Rumex sanguineus* (also in Gdańsk), *Filipendula hexapetala*, *Epilobium adnatum* (also in Gdańsk), *Ajuga genevensis*, *Ajuga reptans* (also in Gdańsk), *Galeopsis bifida* (also in Darłowo and Gdańsk), *Vincetoxicum officinale* and *Hieracium racemosum*, 2) in Świnoujście: *Genista tinctoria*, *Veronica dillenii*; 3) in Ustka: *Potentilla collina*, *Aira caryophyllea* and *Aira praecox*; 4) in Gdańsk: *Matteucia struthiopteris*, *Populus nigra*, *Impatiens noli-tangere*, *Salvia pratensis*, *Senecio silvaticus*, *Carex vesicaria* and *Agropyron intermedium*.

In the synanthropic flora of Polish sea harbours anthropophytes constitute an interesting group. Twenty more of them were found at present than in 1975 (Misiewicz 1976).

From among ergasiophytes, that is transiently growing wild crop plants were noted: 1) in Szczecin and Gdańsk: *Picea excelsa*, *Padus serotina*; 2) in Szczecin and Świnoujście: *Amaranthus caudatus*; 3) in Szczecin: *Colutea arborescens*; 4) in Świnoujście: *Chaenomeles japonica*; 5) in Gdańsk: *Potentilla fruticosa*, *Acer tataricum*, *Thladiantha dubia*, *Inula helenium* and *Rudbeckia hirta*.

New archaeophytes: *Nigella arvensis* — Szczecin, *Erysimum repandum* — Gdańsk, *Gagea arvensis* — Darłowo.

New epocophytes: *Vicia pannonica* — Gdańsk.

New holoagriophytes: *Mimulus guttatus* — Darłowo.

More attention should be devoted to new ephemerophytes, that are species accidentally brought from distant places. Here belong:

1. *Atriplex roseum* L. — an east-Mediterranean species detected in Szczecin and Gdańsk in the region of grain elevators.
2. *Hirschfeldia incana* (L.) Lagreze-Fossat. — a Mediterranean-Iranian-Turanian species. Some few specimens were found in Szczecin and Gdynia, mainly in the region of the grain elevators.
3. *Ambrosia trifida* L. — a north-American species. It occurs rarely in the grain elevator region and on railway premises in large harbours such as Szczecin, Gdynia, Gdańsk.
4. *Andropogon halepensis* (L.) Brot. (= *Sorghum halepense* (L.) Pers.) — an east-Mediterranean species appearing in some years on railway premises and in the region of grain elevators in several ports: Szczecin, Darłowo, Gdynia and Gdańsk.
5. *Setaria faberi* Hermann. Asiatic species, new for the synanthropic flora of Poland, known from numerous sites in Czechoslovakia (Hejny et al. 1973), recorded in the region of grain elevators in Gdynia, and in 1980 was found by the author together with dr V. Jehlik also in Darłowo.

On the margin of the of adventitious sphemic species, particularly those new for the synanthropic flora of Poland, should by mentioned single specimens of *Solanum sisymbriifolium* noted in the period 1976-1979 in the harbours of Szczecin, Gdynia and Gdańsk, mainly in the region of grain elevators and on railway premises.

As regards the taxonomic classification of specimens of the genus *Chenopodium* reported in an earlier paper (Misiewicz 1976) as *Chenopodium carinatum* R.Br. (Dostal 1950), after careful analysis of the collected specimens in the years 1976-1979 and inspection of herbary material it was decided to revise and classify these specimens to *Chenopodium pumilio* R.Br. The causes and cases of confusing these two closely related taxons from Australia are explained by Lhotská and Hejný (1979). These authors also give a precise description of *Ch. pumilio* R.Br. and its distribution the world over and in Czechoslovakia. In the Gdańsk harbour *Ch. pumilio* R.Br. — expands on a wider and wider range, becoming a permanent component of ruderal communities.

6. ANALYSIS OF SYNANTHROPIC FLORA

6.1. STATE OF FLORA

A total of 1055 species of vascular plants were recorded on the area of the Polish sea harbours in the period 1825-1980, belonging to 87 families and 457 genera (Table 4). As regards numbers this flora is exceptionally

rich, it exceeds in this respect the flora of the city of Szczecin which not long ago exhibited 874 species (Ćwikliński 1970) and the most numerous until lately synanthropic flora of Gdańsk and its environs consisting of 1006 species (Schwarz 1967). The recorded 1055 species in the investigated harbours constitute as much as 67.0 per cent of the entire flora of Western Pomerania which includes 1575 species (Czubiński 1950).

Table 4

Families with the largest number of species

Family	Polish sea-harbours 1055 species			Flora of Poland 2250 species	
	total species	%	anthro- pophytes	total species	%
<i>Compositae</i>	140	13.3	77	265	11.8
<i>Gramineae</i>	110	10.4	47	160	7.1
<i>Cruciferae</i>	74	7.0	53	95	4.2
<i>Papilionaceae</i>	76	7.2	37	97	4.3
<i>Rosaceae</i>	48	4.5	19	159	7.1
<i>Caryophyllaceae</i>	47	4.5	11	100	4.4
<i>Scrophulariaceae</i>	40	3.8	23	98	4.4
<i>Labiatae</i>	41	3.9	16	77	3.4
<i>Umbelliferae</i>	37	3.5	14	71	3.2
<i>Chenopodiaceae</i>	36	3.4	31	33	1.5
<i>Polygonaceae</i>	32	3.0	13	31	1.4
<i>Ranunculaceae</i>	24	2.3	8	74	3.3
<i>Boraginaceae</i>	19	1.8	12	31	1.4
<i>Solanaceae</i>	17	1.6	16	11	0.5
<i>Cyperaceae</i>	17	1.6	—	125	5.6
<i>Oenotheraceae</i>	17	1.6	7	22	1.0
<i>Salicaceae</i>	15	1.4	3	31	1.4
<i>Geraniaceae</i>	13	1.2	6	14	0.6
<i>Juncaceae</i>	13	1.2	1	33	1.5
<i>Papaveraceae</i>	13	1.2	12	16	0.7
<i>Liliaceae</i>	12	1.1	8	43	1.9
<i>Euphorbiaceae</i>	11	1.0	8	19	0.8

Most numerously represented families are listed in Table 4. In many cases the number of species found on the harbour areas is but slightly lower than that of species in the flora of Poland, and as regards the family *Solanaceae* and *Chenopodiaceae*, it even exceeds this number. This is due above all to the large proportion of anthropophytes frequently penetrating into the harbours.

All the studied harbours have only 198 species in common (Table 5). Most of them are common plants occurring in abundance on various harbour

Table 5

Species common for a definite number of harbours

Number of harbours	Number of common species
10	198
9	48
8	53
7	41
6	54
5	52
4	71
3	101
2	142
1	295

an.

habitats. Here belong: *Equisetum arvense*, *Polygonum aviculare*, *Rumex acetosa*, *Rumex crispus*, *Chenopodium album*, *Amaranthus retroflexus*, *Stellaria media*, *Berteroa incana*, *Capsella bursa-pastoris*, *Trifolium repens*, *Plantago maior*, *Artemisia vulgaris*, *Cirsium arvense*, *Poa annua*, *Agropyron repens* and others. As frequent may also be considered species appearing in 6-9 harbours (jointly 196 species). In the majority of harbours, that is six to ten a total of 394 common species have been noted. This number is close to the total number of species recorded in small harbours (Table 7).

The number of species found in five or less harbours increases from 52 to 295 occurring only in one of the harbours. These are mostly rare or very rare species which usually are decisive for the flora of the particular harbours (Figs. 2-21). This problem will be considered in a further chapter (Table 6).

In the latest investigations species new for the flora of Poland have been recorded: *Chenopodium pumilio*, *Solanum sisymbriifolium* and *Setaria faberi* as well as numerous species new to the synanthropic flora of the particular cities and the province of Pomerania.

6.2. FLORA OF THE PARTICULAR HARBOURS

The number of species recorded in the harbours varies within the limits of 254 in the Władysławowo harbour to 856 in the Gdańsk harbour (Table 7). These wide differences reflect the different conditions associated with the age of the harbours, the history of their development, their surface area and above all their function and the habitat conditions for the newly arriving species.

The number of species recorded in the period of the most recent investigations (Table 7) do not show such wide variations and may serve

Table 6

Species occurring in only one of the examined harbours

Name of harbour	Total of species	Apophytes							Anthropophytes					
		in forest and shrub communities	in riverside and water communities	in xerothermic grassland communities	in dune and sand communities	in meadow communities	in salt-marsh communities	undetermined	ephemero-phytes	ergasiophytes	archaeophytes	epoecophytes	holoagro-phytes	hemiagro-phytes
Szczecin	37	7	1	3	1	1	—	—	12	8	1	3	—	—
Świnoujście	9	1	—	—	2	—	—	—	1	2	—	2	1	—
Kołobrzeg	8	1	2	—	—	—	—	—	—	4	—	1	—	—
Darłowo	20	8	3	—	—	—	—	—	1	4	3	—	1	—
Ustka	11	1	—	3	3	1	—	—	—	1	—	2	—	—
Łeba	9	6	—	—	1	—	—	—	—	2	—	—	—	—
Władysławowo	1	—	—	—	1	—	—	—	—	—	—	—	—	—
Hel	5	—	2	—	—	—	—	—	—	3	—	—	—	—
Gdynia	8	1	—	—	—	—	—	—	3	4	—	—	—	—
Gdańsk	187	18	12	13	3	6	6	4	77	23	15	10	—	—
Total	295	43	20	19	11	8	6	4	94	51	19	18	2	—

Changes in the synanthropic flora of Polish sea harbours in the period 1825-1979

† — for Szczecin species given by Ćwikliński in 1971a and 1974 have been included

as basis for evaluation of the present contribution of harbours in the introduction of new plants.

In the period 1971-1979 in all the ten examined harbours a total of 892 species were found. They were most numerous in the Gdańsk harbour (697). Schwarz (1967) mentions from the area of this harbour 399 species. Among the plants found in the course of the present investigations 94 species are new for the synanthropic flora of Gdańsk and its environs (Table 8). Beside apophytes and ergasiophytes, noteworthy are particularly

Table 8

New species in the synanthropic flora of Gdańsk and Gdynia in the years 1973-1979

Groups of synanthropic flora	New species in harbour		New species in Gdańsk and its environs (Schwarz 1967)
	Gdańsk	Gdynia	
Apophytes:			
Forest and shrub communities	69	43	26
Meadow communities	28	38	6
Xerothermic grassland communities	29	16	6
Riverside and water communities	30	16	9
Dune and sand communities	11	12	4
Salt-marsh communities	—	—	—
Undefined	3	3	1
Total	170	126	52
Anthropophytes:			
Archaeophytes	15	40	2
Ergasiophytes	57	38	22
Epoecophytes	16	25	9
Ephemeroxytes	9	14	9
Hemiagriophytes	6	1	—
Holoagriophytes	—	—	—
Total	103	117	42
Total	273	243	94

new epoecophytes: *Amaranthus chlorostachys*, *A. blitoides*, *Atriplex oblongifolium*, *Corispermum hyssopifolium*, *Oenothera syrticola* and *O. hoelscheri*, as well as the ephemerophytes: *Chenopodium pumilio*, *Hibiscus trionum*, *Abutilion theophrasti*, *Oenothera octolineata*, *Trifolium resupinatum*, *Solanum sisymbriifolium* and *Ambrosia trifida*. Moreover, sites with 22 species reported in the literatura before 1949, but not found by Schwarz (1967) were noted. Among anthropophytes belong here: *Ranunculus arvensis*, *Nigella damascena*, *Fagopyrum tataricum*, *Chenopodium urbicum*, *Coronopus didymus*, *Lepidium sativum*, *Rorippa austriaca*, *Cerastium tomentosum*, *Carduus nutans*, *Ambrosia artemisiifolia*, *Centaurea solstitialis* and *Alopecurus myosuroides*; from among apophytes:

Thalictrum flavum, *Polygonum hydropiper*, *Spergularia rubra*, *Stachys recta*, *Androsace septentrionalis*, *Holcus mollis* and *Koeleria glauca* (Fig. 2).

In the Gdańsk harbour there are as many as 189 species not recorded in other harbours (Table 6). Among them species earlier mentioned in the literature (Schwarz 1967) and found no more at present prevail. The examples quoted are evidence of a high contribution of the Gdańsk harbour in the introduction of new species.

Second as regards abundance of species comes the Szczecin harbour (634). Extensive areas (1760 ha), differentiated habitats, numerous trade

Fig. 2. Distribution of some new species in the Gdańsk harbour found in the years 1973-1979: 1 – *Chenopodium pumilio*, 2 – *Atriplex oblongifolium*, 3 – *Amaranthus chlorostachys*, 4 – *Amaranthus blitoides*, 5 – *Trifolium resupinatum*, 6 – *Soja hispida*, 7 – *Oenothera octolineata*, 8 – *Oenothera syrticola*, 9 – *Hibiscus trionum*; 10 – *Abutilon theophrasti*, 11 – *Solanum sisymbriifolium*, 12 – *Ambrosia trifida*

contacts and diverse trans-shipments are the factors which decide here of the composition and abundance of the flora. As compared with the material collected by Ćwikliński (1970, 1974) 24 new species were recorded in this harbour new for the synanthropic flora of Szczecin: anthropophytes – *Amaranthus chlorostachys*, *Nigella arvensis*, *Lepidium sativum*, *Sisymbrium wolgensse*, *Hirschfeldia incana*, *Trifolium resupinatum*, *Solanum sisymbriifolium*, *Anthriscus vulgaris*, *Anthemis austriaca*, *Carthamus tinctorius*, *Avena fatua* and *Acorus calamus*. Moreover, rare species of anthropophytes mentioned in the literature before 1945 and not noted since were detected here (Ćwikliński 1970): *Atriplex roseum*, *Eruca sativa*, *Geranium pyrenaicum*, *Astragalus cicer*, *Abutilion theophrasti*, *Andropogon sorghum*, *Bromus squarrosus* and *Alopecurus myosuroides* (Fig. 3).

The remaining harbours as regards the diversity of the flora and their role in introducing new species, may be divided into two groups. To the first belong trade and fishery ports characterised by a larger number of species, from 374 in Ustka to 469 in Darłowo. The remaining harbours with a much smaller amount of species are mainly used for fishery purposes (Table 7).

The abundant flora in the Darłowo harbour is due to its situation in the area of large grain elevators, the diversity of habitats and the frequent intrusion of man into the life of the developing plant communities, associated with the harbour extension. Apophytes dominate, however, in the composition of the flora (64.6%). This may be explained by the immediate vicinity of natural meadow, dune and brushwood communities.

The floristic list of the Gdynia harbour comprises 461 species. The port is relatively new, but owing to numerous trade contacts with many ports of the world and the diversity of trans-shipments, it plays an important role in the introduction of new species. The conditions for their development are, however, worse because the predominant habitats are railway premises and storage yards covered with slag, gravel or concrete slabs. These conditions are reflected in the prevalence of short-lived species (53.8%) and the highest proportion of anthropophytes in the composition of the flora (47.5%). As compared with the records of Schwarz (1967) 243 new species were noted in the Gdynia harbour (Table 8). Noteworthy are also here new anthropophytes: *Chenopodium botrys*, *Amaranthus blitoides*, *A. chlorostachys*, *Sisymbrium wolgensse*, *Solanum sisymbriifolium*, *S. rostratum*, *Abutilion theophrasti*, *Oenothera hoelscheri*, *Ambrosia trifida*, *Chrysanthemum maximum*, *Setaria faberi* and *Cynodon dactylon*. In the Gdynia harbour a number of rare anthropophytes were found which had once been noted in Gdańsk but their occurrence had not been confirmed by Schwarz (1967) after 1949: *Fagopyrum tataricum*, *Atriplex tataricum*, *Cerastium tomentosum*, *Stachys recta*, *Malva pusilla*, *Carduus nutans*, *Ambrosia artemisiifolia* and *Alopecurus myosuroides*.

Fig. 3. Distribution of some of new species in the Szczecin harbour found in the years 1971-1979: 1 – *Chenopodium botrys*, 2 – *Amaranthus chlorostachys*, 3 – *Sisymbrium wolgensse*, 4 – *Eruca sativa*, 5 – *Abutilon theophrasti*, 6 – *Geranium pyrenaicum*, 7 – *Anthriscus vulgaris*, 8 – *Solanum sisymbriifolium*, 9 – *Carthamus tinctorius*, 10 – *Ambrosia trifida*

To regions particularly rich in new species in the Gdynia harbour belong the Indyjskie and Bojowników o Pokój quays along which are situated grain elevators, fat processing plants, rice pearling mills and storehouses for cotton. Very numerous species were also found in the region of mass trans-shipment on the Węglowe Pier (Fig. 4a, b).

The Świnoujście harbour also exhibits a large number of species (457). As compared with an earlier elaboration (Misiewicz 1976) in the period 1976-1979 as many as 70 new species were found in this harbour. This is due to the intensive enlargement of the port, increasing trade contacts and numbers of trans-shipments. A factor restricting the development of plants in the harbour is the aridity of habitats situated on old dune sites.

In the Świnoujście harbour only nine new species were recorded: *Spergula vernalis*, *Armeniacia vulgaris*, *Sorbus aria*, *Genista tinctoria*, *Chaenomeles japonica*, *Aster salignus*, *Echinops sphaerocephalus*, *Lactuca tatarica* and *Vulpia bromoides*. *Lactuca tatarica* found for the first time in Świnoujście by Urbański (1958) spreads to natural dune communities. It is, therefore, no doubt a holoagriophyte which in the present investigations was also found on building premises in the coal port.

Particularly noteworthy are the species new to the flora of the island of Uznam and Wolin and recorded in the Świnoujście harbour, but not mentioned by Piotrowska (1966a): *Polygonum sachalinense*, *Galeopsis angustifolia*, *Erigeron annuus*, *E. ramosus*, *Trifolium resupinatum* and *Phalaris canariensis*. Worth noting are the numerous sites with *Potentilla intermedia*, a species known from earlier publications (Holzfuss 1916, 1937, after Ruthem 1892). Its occurrence in this region was put in doubt by Piotrowska (1966a). The abundant appearance of *Plantago maritima* on the eastern break-water should be noted.

In the Kołobrzeg harbour a total of 441 species were recorded. The main factors affecting the composition of the flora in this harbour are: the situation of grain elevators, trans-shipment of general cargo and some mass goods as well as the rich flora of the city (Müller 1904, Żukowski 1960, 1961). To species found only in this harbour belong: *Lunaria annua*, *Gypsophila muralis*, *Fragaria ananassa*, *Nicotiana rustica*, *Tulipa silvestris*, *Glyceria aquatica* and *Anthoxanthum aristatum*. Noteworthy are numerous specimens of *Reseda luteola* which is also common in the city. On some few sites in the northwestern part of the country *Rumex confertus* is known (Trzcińska-Tacik 1963, Ćwikliński 1970, 1974) and *Anthoxanthum aristatum* (Misiewicz 1970, 1977b), it was also detected in the Kołobrzeg harbour.

A total of 374 species occur in the Ustka harbour. The factor decisive for the rich flora of this small harbour is the presence in it of grain elevators, whereas the large area covered with a permanent blanket and the prevalence of sandy sites limit the possibility of development of the newcomers. In the Ustka harbour 11 species were noted not found in other harbours: among anthropophytes: *Sedum spurium*, *Tropaeolum maius* and *Oenothera chicaginesis*, and among rare apophytes: *Linaria odora* occurring in a sandy storage yard and *Aira praecox* and *A. caryophyllea*.

The remaining three harbours, Łeba, Władysławowo and Hel which

mainly serve as fishery ports have a lower number of synanthropic species (Table 7), mainly because of the absence of grain elevators, and in the case of Łeba also on account of the lack of railway tracks. The large area of the port covered with concrete limits the sites favourable to plants. This situation is reflected in a prevalence of native species of synanthropic flora (Table 7). In the Hel harbour an important factor for the composition of the flora is the spreading of peat and sowing of lawns on the site of sandy wasteland. In the first year after this the flora was highly differentiated. Beside species typical for coastal dunes such as: *Elymus arenarius*, *Honckenya peploides*, *Lathyrus maritimus* and *Ammophila arenaria*, there also grew species of moist habitats such as *Hydrocotyle vulgaris*, *Cirsium palustre*, *Deschampsia caespitosa* and others.

In the harbours of Łeba, Władysławowo and Hel some species not recorded in other harbours were detected, but these are mainly apophytes and, from among anthropophytes, ergasiophytes. The site with *Eryngium maritimum* persisting in Władysławowo for some years is worth noting here.

6.3. BIOLOGICAL DURABILITY OF SPECIES

Perennial plants constituting 52.1 per cent in the floristic composition of the harbours examined predominate slightly (Table 9). Among annual plants the majority are anthropophytes (68.5%), and among perennials apophytes (71.8%). Comparison of the particular harbours shows, however, rather wide differences. In those where secondary habitats dominate the proportion of annual species is higher. In the case of Gdynia it amounts to as much as 53.8, in Gdańsk 50.2 and in Świnoujście 50.1 per cent. In Ustka and Władysławowo the annual plants constitute a high proportion (49.2 and 48.0%, respectively). This may be attributed to intensive human activity. Perennial plants grow above all on relatively durable habitats, for instance along fences, walls, on lawns, slopes of railway banks, forest plantations etc. On storage yards, building premises, refuse heaps, fresh earth and in small gardens, on the contrary, annual plants dominate.

Fig. 4. Distribution of some of new species in the Gdynia harbour found in the years 1973-1979:

- A) 1 – *Chenopodium botrys*, 2 – *Amaranthus chlorostachys*, 3 – *Sisymbrium wolgensse*, 4 – *Oenothera hoelscheri*, 5 – *Abutilon theophrasti*, 6 – *Solanum rostratum*, 7 – *Solanum sisymbriifolium*
 B) 1 – *Ambrosia artemisiifolia*, 2 – *Ambrosia trifida*, 3 – *Ambrosia psilostachya*, 4 – *Iva xanthiifolia*, 5 – *Cynodon dactylon*, 6 – *Setaria faberi*, 7 – *Alopecurus myosuroides*

Table 9

Permanence of species of synanthropic flora in Polish sea harbours

Name of harbour	Number of species total	Annual and biennial species						Perennial species					
		apophytes		anthropophytes		total		apophytes		anthropophytes		total	
		number	%	number	%	number	%	number	%	number	%	number	%
Total	1055	159	31.5	346	68.5	505	47.9	395	71.8	155	28.2	550	52.1
Szczecin	634	99	33.0	201	67.0	300	47.3	263	78.7	71	21.3	334	52.7
Świnoujście	457	88	38.4	141	61.6	229	50.1	176	77.2	52	22.8	228	49.9
Kołobrzeg	441	76	38.2	123	61.8	199	45.1	192	79.3	50	20.7	242	54.9
Darłowo	469	84	39.1	131	60.9	215	45.8	219	86.2	35	13.8	254	54.2
Ustka	374	70	38.0	114	62.0	184	49.2	156	82.1	34	17.9	190	50.8
Łeba	323	57	45.6	68	54.4	125	38.7	175	88.4	23	11.6	198	61.3
Władysławowo	254	49	40.2	73	59.8	122	48.0	123	93.2	9	6.8	132	52.0
Hel	331	61	39.4	94	60.6	155	46.8	153	86.9	23	13.1	176	53.2
Gdynia	461	78	31.5	170	68.5	248	53.8	164	77.0	49	23.0	213	46.2
Gdańsk	856	128	29.8	302	70.2	430	50.2	328	77.0	98	23.0	426	49.8

6.4. ANALYSIS OF APOPHYTES

Analysis of the synanthropic flora of the studied harbours was based on the classification of Thellung, modified and simplified by Kornaś (1968a). Crop plants which transiently turned wild were classified as ergasiophytes.

Among the 1055 species of synanthropic flora in the harbours there are 554 apophytes (52.5%) (Fig. 5). As apophytes were classified native

Fig. 5. The ratio of apophytes to the anthropophytes in the synanthropic flora of Polish sea harbours

species found on synanthropic sites. The large proportion of this group of plants is due to the presence in the immediate neighbourhood and in the area of the harbour of seminatural habitats (e.g. forest plantations, rivers, dunes, coastal coniferous forest, meadows etc.). Many of these species penetrate into synanthropic habitats. Some apophytes, particularly the meadow ones and those of moist habitats, penetrated recently into the harbours on sandy sites in connection with laying out of lawns, for instance in Świnoujście, Hel and the Northern Port in Gdańsk.

According to their aptitude to natural plant communities, apophytes were divided into forest, brushwood, meadow, riverside and aquatic, of xerothermic swards, dune and sand, and salt marsh ones (Table 10).

Forest and brushwood apophytes are represented by 187 species (33.8%). They mainly appear in forest plantations, along fences, round cottages, on railway premises and storage yards, where they find convenient conditions for growth, most similar to natural ones. Many of these plants constitute a permanent component of harbour vegetation and are common, frequently

Table 10
Participation of apophytes in the synanthropic flora of Polish sea harbours

Name of harbour	Total apo-phytes	Apophytes													
		in forest and shrub communities		in meadow communities		in riverside and water communities		in xerothermic grassland		in dune and sand communities		in salt-marsh communities		indefinite	
		num-ber	%	num-ber	%	num-ber	%	num-ber	%	num-ber	%	num-ber	%	num-ber	%
Total	554	187	33.8	115	20.8	88	15.9	74	13.4	52	9.4	18	3.3	20	3.6
Szczecin	362	119	32.9	93	25.7	57	15.8	51	14.1	27	7.5	4	1.1	15	4.1
Świnoujście	264	74	28.0	73	27.7	29	11.0	37	10.2	33	12.5	9	3.4	9	3.4
Kołobrzeg	268	77	28.7	85	31.7	45	16.8	31	11.6	16	6.0	3	1.1	11	4.1
Darłowo	303	104	34.3	87	28.7	49	16.2	23	7.6	25	8.3	3	1.0	12	4.0
Ustka	226	71	31.4	63	27.9	16	7.1	34	15.0	33	14.6	—	—	9	4.0
Łeba	232	90	38.8	67	28.9	22	9.5	22	9.5	24	10.3	—	—	7	3.0
Władysławowo	172	40	23.3	63	36.5	14	8.1	23	13.4	23	13.4	1	0.6	8	4.7
Hel	214	68	31.8	63	29.4	31	14.5	23	10.7	20	9.3	—	—	9	4.2
Gdynia	242	78	32.2	70	28.9	23	9.5	33	13.6	27	11.2	1	0.4	10	4.1
Gdańsk	456	148	32.5	109	23.9	68	14.9	60	13.2	39	8.6	17	3.7	15	3.3

occurring in masses as for instance *Aegopodium podagraria*, *Glechoma hederacea*, *Stellaria media*, *Anthriscus silvestris* or *Artemisia vulgaris*. To rare forest apophytes belong in harbour areas among others *Equisetum hiemale* and *Lathyrus silvester*.

In the harbours 115 species (20.8%) meadow apophytes were found, among them the following ones are common: *Equisetum arvense*, *Rumex acetosa*, *Cerastium vulgatum*, *Ranunculus acer*, *Trifolium repens*, *T. pratense*, *Lotus corniculatus*, *Daucus carota*, *Veronica chamaedrys*, *Plantago lanceolata*, *Galium mollugo*, *Bellis perennis*, *Achillea millefolium*, *Taraxacum officinale*, *Dactylis glomerata*, *Poa annua* and others.

Riverside and aquatic apophytes constitute 15.9 per cent (88 species) of the native flora on synanthropic sites in the harbours. Frequently recorded are among others: *Rumex crispus*, *Polygonum amphibium*, *P. persicaria*, *P. tomentosum*, *Sagina procumbens*, *Rorippa palustris*, *Potentilla anserina*, *Mentha arvensis*, *Bidens tripartitus*, *Juncus bufonius*, *Poa palustris*, *Agropyron repens*.

Apophytes of xerothermic swards are found in 74 species (13.3%) mainly on railway premises in the harbours. To this group belong *Silene inflata*, *Arenaria serpyllifolia*, *Potentilla argentea*, *Sedum acre*, *Medicago lupulina*, *Pimpinella saxifraga*, *Centaurea rhenana*, *Poa compressa* and others.

Apophytes of dunes and sands in the number of 52 species (9.4%) are an important component of the vegetation, particularly in harbours which arose on the site of coastal dunes. Frequent are here *Ammophila arenaria*, *Elymus arenarius*, *Rumex acetosella*, *Erophila verna*, *Corynephorus canescens* and others, and rare for instance *Linaria odora*, *Eryngium maritimum*, *Filago arvensis*, *Gypsophila muralis*, *Plantago indica*, *Chondrilla juncea*, *Aira praecox* and *A. caryophylla*.

Salt marsh apophytes are represented by as few as 18 species (3.3%). *Puccinella distans* is the only common one, mainly on storage yards, railway premises and breakwaters. Moreover, from among the species of this group lately the following were recorded: *Atriplex litorale*, *Spergularia salina*, *Trifolium fragiferum*, *Archangelica litoralis*, *Glaux maritima*, *Centaureum vulgare*, *Aster tripolium* and *Plantago maritima*, which is very abundant on the eastern breakwater in Świnoujście.

For 20 species of apophytes (3.6%) their apurtenance to natural communities could not be established.

The proportion of the particular groups of apophytes differs in each of the examined harbours (Table 10). This results mainly from the different habitat conditions. In the Władysławowo and Kołobrzeg harbours since forest and brushwood are lacking, the dominant group consists of meadow species, whereas in Ustka, Władysławowo and Świnoujście species growing on dunes and in sandy swards are abundant, with simultaneous scarce

occurrence of species of moist habitats. Most numerous salt marsh species occur in the Gdańsk (17) and Świnoujście (9) harbours, whereas in Ustka, Łeba and Hel representatives of this group were not noted.

6.5. ANALYSIS OF ANTHROPOPHYTES

The group of anthropophytes, that is plants of foreign origin, comprises 501 species, thus 47.5 per cent of the whole synanthropic flora of the harbours. Here belong plants brought in prehistoric or early historic times, that means before the 15th century (archaeophytes) or in more recent times — koenophytes (ephemerophytes, epocophytes, agriophytes). There is also a considerable per cent of crop plants which have become wild here (ergasiophytes). The proportion of the particular groups of anthropophytes in the particular harbours is shown in Table 11.

Ergasiophytes constitute the most numerous group among anthropophytes (146 species — 29.1%). To ergasiophytes were classified contemporary crop plant species transiently introduces into synanthropic habitats or species turning wild in regions where they used to be cultivated. The group of ergasiophytes consists mainly of cereals, ornamental plants, park trees and shrubs and vegetable species. The main site of their occurrence are frequently habitats radically transformed by man as for instance refuse heaps in the region of grain elevators, railway premises, roadsides, building premises etc. Ergasiophytes usually occur singly or transiently, but sometimes also in masses, occupying large areas, as for instance *Triticum vulgare* and other cereals in the neighbourhood of grain elevators.

The second group as regards abundance consists of archaeophytes (137 species — 27.3%). They are present on all the specified habitats, constituting the dominating group beside meadow apophytes. To frequent ones belong: *Sisymbrium officinale*, *S. altissimum*, *Descurainia sophia*, *Lepidium ruderalis*, *Erysimum cheiranthoides*, *Sinapis arvensis*, *Raphanus raphanistrum*, *Capsella bursa-pastoris*, *Malva neglecta*, *M. silvestris*, *Vicia hirsuta*, *Erodium cicutarium*, *Convolvulus arvensis*, *Echium vulgare*, *Ballota nigra*, *Linaria vulgaris*, *Tripleurospermum inodorum*, *Senecio vulgaris*, *Carduus acanthoides*, *Bromus tectorum* and others.

As rare anthropophytes on harbour areas may be listed: *Chenopodium hybridum*, *Ch. murale*, *Ch. bonus-Henricus*, *Atriplex nitens*, *Portulaca oleracea*, *Mercurialis annua*, *Neslia paniculata*, *Nigella arvensis*, *Ranunculus arvensis*, *Vaccaria pyramidata*, *Malva pusilla*, *Conium maculatum*, *Lappula myosotis*, *Asperugo procumbens*, *Chrysanthemum segetum*, *Alopecurus myosuroides*, *Avena fatua*, *Bromus secalinus* and others.

Earlier sites with the recorded following archaeophytes (Schwarz 1967, Ćwikliński 1970) have not been confirmed in recent studies (1971-1979):

Table 11
Participation of anthropophytes in synanthropic flora of Polish sea harbours

Name of harbour	total	Anthropophytes											
		ergasiophytes		archaeophytes		ephemerophytes		epoecophytes		hemiagriophytes		holoagriophytes	
		number	%	number	%	number	%	number	%	number	%	number	%
Total	501	146	29.1	137	27.3	122	24.4	83	16.6	9	1.8	4	0.8
Szczecin	272	73	26.8	104	38.2	28	10.3	58	21.3	8	2.9	1	0.4
Świnoujście	193	45	23.3	90	46.6	8	4.1	42	21.8	5	2.6	3	1.6
Kołobrzeg	173	48	27.7	83	48.0	—	—	34	19.7	6	3.5	2	1.2
Darłowo	166	41	24.7	93	56.0	4	2.4	24	14.5	3	1.8	1	0.6
Ustka	148	41	27.7	76	51.3	3	2.0	25	16.9	3	2.0	—	—
Łeba	94	18	19.8	59	61.5	—	—	11	12.1	3	3.3	—	—
Władysławowo	82	13	15.9	49	59.8	—	—	18	21.9	2	2.4	—	—
Hel	117	26	22.2	65	55.5	1	0.9	23	19.7	2	1.7	—	—
Gdynia	219	58	26.5	92	42.0	21	9.6	45	20.5	3	1.4	—	—
Gdańsk	400	98	24.5	126	31.5	97	24.3	68	17.0	9	2.3	2	0.5

Chenopodium opulifolium, *Ch. vulvaria*, *Euphorbia exigua*, *Adonis aestivalis*, *Fumaria vaillantii*, *Coronopus procumbens*, *Bupleurum rotundifolium*, *Veronica opaca*, *Stachys arvensis*, *Sherardia arvensis*, *Avena strigosa* and *Lolium remotum*. Probably these species died out, maybe, some of them may still be detected.

Most labile is the group of ephemerophytes transiently introduced into harbours from distant regions. In all the harbours jointly 122 species (24.4%) were recorded in the period 1825-1979. The considerable number of ephemerophytes shows the important role of ports in their introduction, it is also the result of many years of studies on the synanthropic flora in the Gdańsk harbour (cf. Schwarz 1967), where most of these species were observed only transiently. In recent years (1971-1979) the following ephemerophytes were recorded in the harbours: *Atriplex roseum*, *Coronopus didymus*, *Sisymbrium orientale*, *Hirschfeldia incana*, *Astragalus cicer*, *Soja hispida*, *Oenothera octolineata*, *Hibiscus trionum*, *Abutilon theophrasti*, *Anthriscus vulgaris*, *Solanum rostratum*, *S. alatum*, *Ambrosia trifida*, *A. artemisiifolia*, *Artemisia annua*, *Carthamus tinctorius*, *Centaurea solstitialis*, *Bromus japonicus*, *B. squarrosus*, *Cynodon dactylon*, *Phalaris canariensis* and *Sorghum halepense*. Three proved to be new to the flora of Poland: *Chenopodium pumilio*, *Solanum sisymbriifolium* and *Setaria faberi*. Most of the enumerated ephemerophytes were detected in large trade harbours, that is Gdańsk, Gdynia and Szczecin, although in Darłowo for instance some of these species were observed: *Sisymbrium orientale*, *Sorghum halepense* and *Setaria faberi*.

Epoecophytes to which 83 species were classified constitute 16.6 per cent among foreign plants. Some of them grow on various habitats in the harbours and are very common as for instance: *Amaranthus retroflexus*, *Diplotaxis muralis*, *D. tenuifolia*, *Sisymbrium altissimum*, *Melilotus albus*, *M. officinalis*, *Veronica persica*, *Erigeron canadensis*, *Galinsoga parviflora*, *G. ciliata*, *Senecio vernalis*, *Lactuca serriola*.

Rare epoecophytes in harbours are: *Rumex confertus*, *Polygonum cuspidatum*, *P. sachalinense* (Fig. 6), *Kochia scoparia*, *Amaranthus chlorostachys*, *A. albus*, *A. blitoides*, *Silene conica*, *Rorippa austriaca*, *Erucastrum gallicum*, *Cardaria draba*, *Bunias orientalis*, *Potentilla recta*, *Iva xanthiifolia*, *Anthoxanthum aristatum* and others. Some of them as for instance *Sisymbrium wolgensense* and *Acroptilon repens* were recorded in the harbours only as late as the most recent investigation (1971-1979) and occupy small areas.

Agriophytes, that is species established in seminatural (hemiagriophytes) and natural (holoagriophytes) communities are represented jointly by 13 species (2.6%). The low number of agriophytes results from the fact that secondary habitats which are frequently transformed by man and on which natural communities do not occur were chiefly the subject of observation.

As hemiagriophytes were classified: *Lupinus polyphyllus*, *Impatiens parviflora*, *Sambucus racemosa*, *Bidens melanocarpus*, *Erigeron annuus*, *Helianthus*

tuberosus, *Solidago canadensis*, *S. serotina* and *Juncus macer*, and as holoagriophytes: *Lactuca tatarica*, *Mimulus guttatus*, *Elodea canadensis* and *Acorus calamus*.

The numerical proportions of the particular groups of anthropophytes in the particular harbours (Table 11) are similar to the data for the harbours as a whole only in the case of the Gdańsk harbour. In the remaining ones there is a marked dominance of archaeophytes. Next come ergasiophytes, epocophytes, ephemerophytes, hemiagriophytes and holoagriophytes. Similar numerical and percentual proportions within anthropophytes are found in the synanthropic flora of other regions, for instance towns and boroughs of the Łódź district (Sowa 1971), river ports of the upper Odra (Szotkowski 1972) and Legnica, Lubin and Polkowice (Anioł-Kwiatkowska 1974). The much higher number of ephemerophytes, archaeophytes, epocophytes and ergasiophytes in sea harbours indicates the role of the latter in the introduction of new species. The low number of hemiagriophytes and holoagriophytes, on the other hand, is due to the small area with natural and seminatural communities in the harbours.

6.6. GEOGRAPHICAL AND GENETIC ANTHROPOPHYTE SPECTRUM

Most numerous represented are south-European and west-Asiatic species (32.5%) and those of southern Europe (30.5%). From among extra-European countries, those from America constitute 14.4 per cent, Asiatic ones 12.4 per cent of the total number of anthropophytes. The contribution of species from other regions is very low, however, noteworthy is the appearance of a representative of Australian flora, *Chenopodium pumilio* so far was not represented in the synanthropic flora of Poland. The proportions of species originating from various regions of the world, which were recorded in the particular harbours are close to the data concerning the flora of the harbours as a whole (Kornaś 1972). (Table 12).

Analysis of the origins of the particular groups of anthropophytes indicates wide differences between them (Table 13). Among archaeophytes south-European and west-Asiatic species prevail (69%) and south-European ones (47%), whereas for obvious reasons American species are absent. In the group of epocophytes, beside south-European and west-Asiatic species (26%) and south-European ones (24%), American species (22%) are numerous represented. No less differentiated is the group of ergasiophytes in which south-European species (41%) and Asiatic ones (40%) prevail. The widest differences as regards origin are noted among ephemerophytes (Table 13). They reflect the numerous trade contacts of Polish sea ports with many regions of the world on all continents.

Centres of origin of the anthropophytes in the synanthropic flora of Polish sea harbours

Name of harbour	Total anthro- pophytes	Country of origin															
		Southern Europe and western Asia		Southern Europe		America		Asia		Northern Western and Central Europe		Africa		Australia		unknown country of origin	
		num- ber	%	num- ber	%	num- ber	%	num- ber	%	num- ber	%	num- ber	%	num- ber	%	num- ber	%
Total	501	163	32.5	153	30.5	72	14.4	62	12.4	26	5.2	2	0.4	1	0.2	22	4.4
Szczecin	272	89	32.7	66	24.3	51	18.8	44	16.2	9	3.3	—	—	—	—	13	4.8
Świnoujście	193	67	34.7	49	25.4	34	17.6	28	14.5	6	3.1	—	—	—	—	9	4.7
Kołobrzeg	173	66	38.2	43	24.8	26	15.0	22	12.7	5	2.9	—	—	—	—	11	6.4
Darłowo	166	66	39.8	43	25.9	19	11.4	22	13.2	5	3.0	1	0.6	—	—	10	6.0
Ustka	148	50	33.8	41	27.7	24	16.2	21	14.2	4	2.7	—	—	—	—	8	5.4
Łeba	91	41	45.1	18	19.8	11	12.1	10	11.0	4	4.4	—	—	—	—	7	7.7
Władysławowo	82	33	40.2	20	24.4	8	9.8	14	17.1	1	1.2	—	—	—	—	6	7.3
Hel	117	47	40.2	31	26.5	17	14.5	14	12.0	2	1.7	—	—	—	—	6	5.1
Gdynia	219	78	35.6	46	21.0	35	16.0	36	16.4	10	4.6	1	0.6	—	—	13	5.9
Gdańsk	400	142	35.5	115	28.8	56	14.0	43	10.8	25	6.3	—	—	1	0.3	17	4.3

Table 13

Geographical spectrum of anthropophyte synanthropic flora of Polish sea harbours

Country of origin	Total species	Ergasio-phytes	Archaeo-phytes	Epheme-rophytes	Epoeco-phytes	Hemia-griophy-tes	Holo-agrio-phytes
Southern Europe and Western Asia	163	18	69	49	26	—	1
Southern Europe	153	41	47	40	24	1	—
America	72	29	—	13	22	6	2
Asia	62	40	8	7	5	1	1
Northern, Western and Central Europe	26	10	4	9	2	1	—
Africa	2	—	—	2	—	—	—
Australia	1	—	—	—	—	—	—
Unknown country of origin	22	8	9	1	4	—	—
Total	501	146	137	122	83	9	4

6.7. FLORA OF HARBOUR HABITATS

The numerical results concerning the flora of the particular habitats (Tables 14 and 15) are compiled only on the basis of the author's own records, thus, they concern 892 plant species. The flora of the balast heaps in the old Gdańsk port is discussed by Schwarz (1967).

The richest habitat, similarly as in the river ports of the lower Odra (Sztokowski 1972) is waste land. Here a total of 686 species has been noted. Waste land is characterised by highly differentiated habitat conditions and in this connection also composition of flora. The vegetal cover is frequently destroyed here, and owing to this newcomers are able to settle here, as for instance in the Gdynia harbour *Solanum rostratum* and *Sisymbrium wolgensse*. Very many species were recorded on refuse heaps in the region of grain elevators. For instance in Szczecin close to the elevator „Ewa”: *Anthriscus vulgaris*, *Solanum sisymbriifolium* and *Alopecurus myosuroides*; in Darłowo: *Geranium dissectum*, *Trifolium resupinatum*, *Chrysanthemum segetum*, *Setaria faberi* and *Sorghum halepense*. In Gdańsk at the Zbożowe Quay: *Mercurialis annua*, *Iva xanthiifolia* and others; in Gdynia: *Abutilon theophrasti*, *Vaccaria pyramidata*, *Solanum sisymbriifolium*, *Artemisia annua* and *Alopecurus myosuroides*.

On railway premises a total of 664 species were found. Anthropophytes constitute 45.7 per cent here. As compared with other railway premises of Poland (Table 15) this habitat only ranks second after the flora of the

Table 14

Participation of species on particular sites in Polish sea harbours in the years 1971-1979

Name of harbour	Type of habitat									Species occurring in only one place								
	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9
Total	664	582	686	289	166	326	372	273	84	29	16	23	23	8	2	13	34	—
Szczecin	444	373	426	199	—	246	194	226	—	26	2	14	13	—	2	—	18	—
Świnoujście	376	198	308	76	58	134	127	87	55	4	2	3	8	4	—	2	4	—
Kołobrzeg	123	382	239	137	46	67	36	74	36	1	3	11	18	—	—	—	1	—
Darłowo	127	235	394	187	238	82	59	132	14	2	1	12	18	6	—	2	20	—
Ustka	186	149	254	46	142	121	143	134	30	—	—	9	—	4	—	1	2	—
Łeba	—	—	226	136	88	64	127	130	11	—	—	4	4	1	—	—	23	—
Władysławowo	145	92	129	18	56	58	72	—	26	—	—	1	—	4	—	—	—	—
Hel	136	272	182	6	—	62	182	116	8	—	2	6	—	1	—	8	4	—
Gdynia	409	384	152	59	—	214	154	165	—	18	2	6	—	—	—	1	5	—
Gdańsk	547	522	554	195	74	275	296	212	—	23	9	18	16	—	—	4	29	—

1 — railway premises; 2 — storage yards; 3 — wasteland; 4 — edges of docks, and channels; 5 — dunes and sandy wastes; 6 — streets, sidewalks, parkings; 7 — lawns, flowerbeds, small gardens; 8 — forest plantations; 9 — breakwaters.

Table 15

Participation of apophytes and anthropophytes in the synanthropic flora of selected railway premises of Poland

Railway premises	Total species	Apophytes		Anthropophytes	
		number	%	number	%
Railway premises in Polish sea-harbours	664	354	54.3	298	45.7
Railway premises in Szczecin district (Ćwikliński 1974)	741	476	64.3	265	35.7
Railway premises in river-harbours: Koźle, Gliwice, Opole (Szotkowski 1972)	512	293	57.2	215	41.9
Railway tracks of East Pomerania coast (Lejmbach et al. 1975)	448	324	72.3	124	27.3
Railway stations in Opole (Michalak 1970)	431	225	52.1	204	47.9
Railway stations in Cracow (Kornaś et al. 1959)	399	242	60.6	157	39.4
Railway station in Kluczbork (Sendek 1971)	381	212	55.6	169	44.6
Railway premises in Czechowice-Dziedzice (Zajac i Zajac 1975)	297	233	66.8	74	33.2

railway premises of the Szczecin district (Ćwikliński 1974) where apophytes are more numerous. The richness of the flora on railway premises is due above all to the frequent carrying of diaspores with the transported goods, particularly grain, oleraceous plant seeds, general cargo and mass goods. On the other hand, the unfavourable conditions prevailing usually on railway premises restrict plant development (Ćwikliński 1974). Railway premises play an important part in the spread of new species, for instance in the Gdańsk harbour during the latest investigations the following species were found: *Solanum sisymbriifolium*, *Oenothera syrticola*, *Hibiscus trionum*, *Artemisia annua* and *Alopecurus myosuroides*, and in the Gdynia harbour: *Sisymbrium wolgensse*, *Cynodon dactylon*, *Soja hispida*, *Artemisia annua*, *Vaccaria pyramidata* and *Alopecurus myosuroides*.

Storage yards also exhibit a rich flora, 582 species have been noted here. The factor decisive for these habitat conditions is the introduction of numerous diaspores during trans-shipment and storage of goods. Storage yards represent a wide diversity of habitat conditions as regards the surface (gravel, slag, concrete slabs), the kind of usage (storage, trampling, traffic), temperature and light conditions. Moreover, the substrate frequently changes according to the chemical and mechanical components of the goods (mineral fertilisers, coal dust, sulphur, lubricants etc.). In these difficult conditions only some plant species are able to develop. Therefore, storage yards are often bare and the competition factor is not as strong here as in other habitats.

Fig. 6. Distribution of *Rumex confertus* (a), *Polygonum sachalinense* (b), and *Polygonum cuspidatum* (c) in Polish sea harbours: a, b, c, — new localities; a₁, b₁, c₁ — information from literature

Fig. 7. Distribution of *Chenopodium botrys* (a), *Ch. pumilio* (b), *Atriplex oblongifolium* (c) and *A. tataricum* (d) in Polish sea harbours: a, b, c, d, — new localities; c₁, d₁ — information from literature

Fig. 8. Distribution of *Sisymbrium orientale* (a), *Erucastrum gallicum* (b), *Eruca sativa* (c) and *Sisymbrium wolgens* (d) in Polish sea harbours: a, b, c, d — new localities; a₁, b₁ — information from literature

Fig. 9. Distribution of *Potentilla intermedia* (a), *P. recta* (b) and *Sanguisorba minor* (c) in Polish sea harbours: a, b, c – new localities; a₁, b₁ – information from literature

Fig. 10. Distribution of *Trifolium resupinatum* (a), *Vicia grandiflora* (b), *Lathyrus tuberosus* (c), and *Soja hispida* (d) in Polish sea harbours: a, b, c, d – new localities; c₁, d₁ – information from literature

Fig. 11. Distribution of *Oenothera hoelscheri* (a), *O. octolineata* (b), *O. syrticola* (c) and *O. chicaginisensis* (d) in Polish sea harbours: a, b, c, d – new localities

Fig. 12. Distribution of *Malva pusilla* (a), *Hibiscus trionum* (b), and *Abutilon theophrasti* (c) in Polish sea harbours: a, b, c — new localities; a₁ — information from literature

Fig. 13. Distribution of *Geranium pyrenaicum* (a) and *Geranium dissectum* (b) in Polish sea harbours: a, b — new localities; a₁, b₁ — information from literature

Fig. 14. Distribution of *Eryngium maritimum* (a), *Archangelica litoralis* (b) and *Anthriscus vulgaris* (c) in Polish sea harbours; a, b, c — new localities; b₁, c₁ — information from literature

Fig. 15. Distribution of *Androsace septentrionalis* (a) and *Glaux maritima* (b) in Polish sea harbours: a, b – new localities; a₁, b₁ – information from literature

Fig. 16. Distribution of *Solanum alatum* (a), *S. rostratum* (b), *S. sisymbriifolium* (c) and *Nicandra physaloides* (d) in Polish sea harbours: a, b, c, d – new localities; a₁ – information from literature

Fig. 17. Distribution of *Linaria arvensis* (a), *L. odora* (b) and *Mimulus guttatus* (c) in Polish sea harbours: a, b, c – new localities

Fig. 18. Distribution of *Plantago maritima* (a) and *Centaurium vulgare* (b) in Polish sea harbours: a, b — new localities; a₁, b₁ — information from literature

Fig. 19. Distribution of *Ambrosia artemisiifolia* (a), *A. trifida* (b) and *A. psilostachya* (c) in Polish sea harbours: a, b, c — new localities; a₁, c₁ — information from literature

Fig. 20. Distribution of *Iva xanthiifolia* (a), *Artemisia annua* (a) and *Centaurea solstitialis* (c) in Polish sea harbours: a, b, c — new localities; a₁, b₁, c₁ — information from literature

Fig. 21. Distribution of *Cynodon dactylon* (a), *Setaria faberi* (b), *Anthoxanthum aristatum* (c) and *Alopecurus myosuroides* (d) in Polish sea harbours: a, b, c, d — new localities; a₁, d₁ — information from literature

In the flora storage yards several plant families are abundantly represented such as *Chenopodiaceae*: *Chenopodium album*, *Ch. glaucum*, *Ch. rubrum*, *Atriplex patula* and *Corispermum hyssopifolium*; *Cruciferae*: *Descurainia sophia*, *Erucastrum gallicum*, *Sisymbrium altissimum*, *Berteroa incana*, *Coronopus didymus*; *Amaranthaceae*: *Amaranthus retroflexus*, *A. albus*, *A. blitoides*. Besides, numerous species characteristic for trampled areas are present here as for instance *Plantago maior*, *Polygonum aviculare*, *Poa annua*.

For numerous plants the storage spaces are their first settling site from which they spread further. This observation is confirmed by studies performed on stockyards in other areas (Szotkowski 1972).

On the banks of harbour basins, channels and rivers a total of 289 species have been listed. To the more abundant apophytes belong here: *Polygonum persicaria*, *P. hydropiper*, *Archangelica litoralis*, *Calystegia sepium*, *Lycopus europaeus*, *Phragmites communis* and others. Among anthropophytes noteworthy are the holoagriophytes: *Elodea canadensis* and *Acorus calamus*. In the fissures of concrete quays may be found: *Capsella bursa-pastoris*, *Descurainia sophia*, *Lepidium rudemale*, *Sagina procumbens*, *Cerastium vulgatum*, *Plantago maior*, *Taraxacum officinale*, *Artemisia vulgaris*, *Tripleurospermum inodorum*, *Senecio viscosus*, *Sonchus arvensis*, *Bromus mollis* and others.

Dunes sometimes utilised as storage space for various materials or treated as waste land have the poorest flora consisting of as few as 166 species. This includes, beside species typical for dunes and sandy areas, also anthropophytes such as: *Sisymbrium altissimum*, *Descurainia sophia*, *Melilotus officinalis*, *Oenothera chicaginensis*, *Echium vulgare*.

On streets, squares, parking spaces 326 species have been found. Apart

from the dominating species of the class *Plantaginetea*, juvenile forms of ergasiophytes are frequently found, particularly of cereals.

Lawns, mainly the newly laid out ones are a favourable habitat for the development of many plant species. Therefore, their flora is relatively rich reaching 372 species. Penetration of new species to these habitats proceeds with the earth spread before seeding, for instance numerous species of moist habitats were introduced in this way onto lawns in the Świnoujście, Hel and Gdańsk Northern harbour. The seeds of some species are dispersed and the plants turn wild as for instance in the Szczecin harbour *Trifolium resupinatum* and together with it *Eruca sativa* as an accompanying weed. When greenery was planted in the Gdańsk Northern Port, such crop plants as *Lupinus angustifolius*, and *Ornithopus sativus* were sown beside the commonly used lawn species and trees and shrubs were planted (Siuta 1973a, b). Wind-carried plants also develop on lawns, some of them rare, as for instance *Centaurea solstitialis* in the Northern Port. The numbers of species recorded on lawns in the particular harbours (Table 14) differ widely, they are higher in the harbours where new large lawns were laid out in the period when the present investigations were performed (Gdańsk, Szczecin, Hel).

Brushwood is of the character of seminatural communities. The latter occur along fences, on waste land, edges of channels and railway premises. Their surface may vary widely. A dense vegetal cover does not favour development of new species, nevertheless, a rather high number (273) of species were found in these habitats. They are mostly forest and brushwood apophytes. Among anthropophytes noteworthy are: *Parietaria officinalis*, *Polygonum cuspidatum*, *P. sachalinense*, *Impatiens parviflora*, *I. roylei*, *Lycium halimifolium* and *Reseda luteola*.

A typical habitat of harbours are breakwaters, the joint length of which in the examined harbours is about 8 km. In this habitat 84 species were found (Tables 14 and 16). Most frequent are species characteristic for trampled sites as for instance *Polygonum aviculare*, *Plantago maior*, *Poa annua*, *Taraxacum officinale*. Noteworthy is the very abundant *Puccinella distans* and *Plantago maritima* on the eastern breakwater in Świnoujście, *Glaux maritima* in Świnoujście and Darłówek, *Honckenya peploides*, *Lactuca serriola* and *Archangelica litoralis* in Świnoujście, *Carex contigua* in Kołobrzeg, *Cakile maritima* in Darłówek and *Salsola kali* in Ustka. Table 16 includes the species noted in at least two of the studied harbours. In only one harbour are found: in Świnoujście *Plantago maritima*, *Rosa rugosa*, *Polygonum persicaria*, *Honckenya peploides*, *Solanum nigrum*, *Matricaria chamomilla*, *Tussilago farfara*, *Salix caprea*, *Chenopodium rubrum*, *Cirsium lanceolatum*, *Urtica dioica*, *Linaria vulgaris*, *Solanum lycopersicum*, *Carex hirta*, *Lactuca serriola*, *Acer negundo*, *Atriplex patula*, *Phragmites communis*, *Festuca arundi-*

Table 16
Breakwater flora

Species	Name of harbour							Number of occurrence sites
	S	K	D	U	Ł	W	H ^x	
<i>Poa annua</i>	+	+	+	+	+	+	+	7
<i>Plantago maior</i>	+	+	+	+	+	+	+	7
<i>Taraxacum officinale</i>	+	+	+	+	+	+	+	7
<i>Polygonum aviculare</i>	+	+	+	+	+	+	+	7
<i>Chenopodium album</i>	+	+	+	+	+	+	+	6
<i>Atriplex hastatum</i>	+	+	+	.	+	.	+	5
<i>Artemisia vulgaris</i>	+	+	.	+	.	+	.	4
<i>Sonchus arvensis</i>	+	+	.	+	.	+	.	4
<i>Sagina procumbens</i>	+	+	.	+	.	+	.	4
<i>Agropyron repens</i>	+	+	.	+	.	+	.	4
<i>Lolium perenne</i>	+	+	.	+	.	+	.	4
<i>Rumex crispus</i>	+	+	.	+	.	+	.	4
<i>Cirsium arvense</i>	+	.	.	+	+	.	+	4
<i>Capsella bursa-pastoris</i>	.	+	.	+	+	.	.	4
<i>Potentilla anserina</i>	+	+	.	.	.	+	.	3
<i>Artemisia campestris</i>	+	.	.	+	.	+	.	3
<i>Agrostis stolonifera</i>	+	.	.	+	.	+	.	3
<i>Bromus mollis</i>	+	.	.	+	.	+	.	3
<i>Poa pratensis</i>	+	+	.	.	.	+	.	3
<i>Leontodon autumnalis</i>	+	.	.	.	+	.	+	3
<i>Matricaria discoidea</i>	.	+	.	+	.	+	.	3
<i>Rumex hydrolapathum</i>	+	+	.	.	.	.	.	2
<i>Lycopus europaeus</i>	+	+	.	.	.	.	.	2
<i>Stachys palustris</i>	+	+	.	.	.	.	.	2
<i>Puccinellia distans</i>	+	.	+	.	.	.	.	2
<i>Chenopodium glaucum</i>	+	.	+	.	.	.	.	2
<i>Elymus arenarius</i>	+	.	+	.	.	.	.	2
<i>Glaux maritima</i>	+	.	+	.	.	.	.	2
<i>Achillea millefolium</i>	+	.	.	+	.	.	.	2
<i>Oenothera sp.</i>	+	.	.	+	.	.	.	2
<i>Cerastium vulgatum</i>	+	.	.	.	+	.	.	2
<i>Tripleurospermum inodorum</i>	+	.	.	.	.	+	.	2
<i>Hieracium umbellatum</i>	+	.	.	.	.	+	.	2
<i>Rumex obtusifolius</i>	.	+	.	+	.	.	.	2
<i>Alnus glutinosa</i>	.	+	.	+	.	.	.	2
<i>Rorippa palustris</i>	.	+	.	.	+	.	.	2
<i>Senecio viscosus</i>	.	.	.	+	.	+	.	2
<i>Poa compressa</i>	.	.	.	+	.	+	.	2
<i>Festuca rubra</i>	.	.	.	+	.	+	.	2
<i>Bromus tectorum</i>	.	.	.	+	.	+	.	2

^x Ś – Świnoujście, K – Kołobrzeg, D – Darłowo, U – Ustka, Ł – Łeba, W – Władysławowo, H – Hel

nacea, *Malva neglecta*. *Archangelica litoralis*; In Kołobrzeg: *Ranunculus sceleratus*, *Mentha arvensis*, *Erigeron canadensis*, *Hordeum murinum*, *Carex contigua*, *Rumex maritimus*, *Helianthus annuus*, *Polygonum nodosum*; in Darłówek: *Cakile maritima*, *Carex arenaria*; in Ustka: *Erodium cicutarium*, *Erysimum cheiranthoides*, *Galinsoga parviflora*, *Salsola kali*, *Senecio vernalis*; in Władysławowo: *Melilotus albus*, *Sonchus asper* and *Dactylis glomerata*.

6.8. STATE OF SYNANTHROPIC FLORA IN THE HARBOURS AS COMPARED WITH THAT OF SOME SELECTED REGIONS OF POLAND

From among the synanthropic floras of various cities and railway premises elaborated to date (Sowa and Olaczek 1978) only those were chosen for comparison which lie in the nearest neighbourhood of the studied harbours. Thus, the compared floras are related as regards natural conditions in which they develop. The appearing difference may, therefore, be explained by the different economic factors such as the degree of industrialisation, the kind of transport of goods etc.

In the groups of floras compared the synanthropic ones of the cities with harbours, such as Szczecin (Ćwikliński 1970), Gdańsk (Schwarz 1967) and Świnoujście (Misiewicz et al. 1979) were taken into account. Valuable information was supplied by comparison of the synanthropic flora of railway premises of the Szczecin district (Ćwikliński 1974), part of which is included within the limits of the harbour complex Szczecin-Świnoujście. The river ports of the upper Odra (Sztokowski 1972) which are in immediate contact with the trade port complex of Szczecin-Świnoujście and with similar economic functions were also taken into account.

The number of synanthropic species in the compared floras is given in Table 17, so is the proportion of the particular anthropophyte groups and the indexes for anthropophytes calculated after Kornaś (1977). These indexes are as follows:

1. Index of synanthropisation:

$$S = \text{archaeophytes} + \text{agriophytes} + \text{epoecophytes}.$$

This index expressed by the absolute number of all species of foreign origin (anthropophytes) permanently established illustrates best the degree of synanthropisation of the studied flora.

2. Index of lability of flora:

$$I = \text{ephemerophytes} + \text{ergasiophytes}.$$

Studies to date have proved (Kornaś 1977) that the number of unestablished species, that means brought accidentally from distant regions (ephemerophytes) or crop plants growing wild (ergasiophytes) increases with every year of observation. The joint number of these species, thus describes, though only approximately, but significantly, the degree of lability of the

Table 17

Quantity of synanthropic flora of Polish sea harbours as compared with the flora of selected towns, railway premises and river harbours

Town or other area	Total, species	Anthropophytes						Numerical indexes			Author and year of publication	
		number of anthropophytes	% of anthropophytes	archaeophytes	agriophytes	epocophytes	ephemerophytes	ergasiophytes	S = archaeophytes + agriophytes + epocophytes	I = ephemerophytes + ergasiophytes		M = $\frac{\text{agro.} + \text{epoc.}}{\text{arch.}}$
Polish sea harbours	1055	501	47.5	137	13	83	122	146	233	268	0.70	
Gdańsk and environs	1006	528	52.5	130	6	82	153	157	218	310	0.61	Schwarz 1967
Szczecin — town and harbour	874	449	51.5	89	4	140	97	119	233	216	1.62	Ćwikliński 1970
Ślupsk	854	354	41.5	113	11	65	34	131	189	165	0.67	Misiewicz 1978a
River harbours of upper Odra	811	362	44.7	127	26	60	47	102	213	149	0.68	Szotkowski 1972
Gorzów Wielkopolski	806	375	46.3	93	4	111	24	143	208	167	1.24	Misiewicz 1971
Railway premises of Szczecińskie district	741	265	35.7	80	—	116	19	50	196	69	1.40	Ćwikliński 1974
Świnoujście — town and harbour	728	277	38.1	102	13	53	17	92	168	109	0.65	Misiewicz et al. 1979
Koszalin	550	230	41.8	55	21	96	58	—	172	58	2.13	Ćwikliński 1971b
Elbląg	492	225	45.7	116	14	30	10	55	160	65	0.38	Schwarz 1971
Stargard Szczeciński	459	210	45.8	73	4	54	7	72	131	79	0.79	Szmajda 1974
Pyrzyce	446	195	43.0	72	2	44	5	69	118	74	0.64	Szmajda 1974

100

particular synanthropic floras, especially when in the cases compared the time of the investigations was similar.

3. Index of degree of modernisation of the flora:

$$M = \frac{\text{agriophytes} + \text{epoecophytes}}{\text{archaeophytes}}.$$

The observations performed prove that the number of archaeophytes is almost stable (Zajac and Zajac 1975, Kornaś 1977). Some species, however, have a tendency to disappearance in connection with the progressing synanthropisation process. At the same time an increase is observed in the number of established new species. Therefore, the ratio of agriophytes and epoecophytes to archaeophytes (M) characterises well the degree of modernisation of a given synanthropic flora.

The data in Table 17 show the state of the synanthropic flora of Polish sea harbours and particularly the contribution of anthropophytes as compared with the floras of other regions. These data are not fully comparable, because the authors frequently differed in their opinions as to the historical-geographical background of certain species. Particularly the flora of Szczecin, and Koszalin arouses controversies. Nevertheless, the general state of the synanthropic flora is reflected in these data as well as some of its characteristics.

As far as the number of species is concerned, the synanthropic flora of the Polish sea harbours with 1055 species belongs to the richest synanthropic floras in this country. Numerous are also species in the floras of Gdańsk and environs (1006), Szczecin (874), Słupsk (854), of the upper Odra river ports (811) and of Gorzów (806). These floras also include large numbers of anthropophytes (528, 449, 354, 362, 375, respectively).

The railway premises of the Szczecin district and the town of Świnoujście exhibit markedly poorer floras (741 and 729 species). The number of anthropophytes is also much lower (265 and 277, respectively). In the remaining towns of north-western Poland included in the comparison the number of synanthropic species is much lower and they constitute as little as 50 per cent of the harbour flora. Here too the number of anthropophytes is low (Table 17).

The indexes for anthropophytes calculated for the compared floras indicate in a measurable way their present state and developmental tendencies. The index of synanthropisation (S) of the Polish sea harbour flora amounts to 233 established foreign species, thus it is very high and equal to the corresponding index for Szczecin. The index of lability (I) of the harbour flora is also high reaching 268 species. In this case, although crop species turning wild dominate (146), a considerable number of species introduced (122) from distant areas into the harbours (ephemerophytes) was also found. As regards the latter, a relatively high number (157) was found only in

Gdańsk, whereas in the remaining towns they were less numerous (Table 17).

The index of modernisation of the synanthropic harbour flora amounting to 0.70 reflects, on the one hand, a rather high proportion of archaeophytes (137 species), and on the other hand, it is the result of a large number (96) of species introduced in modern times which with time joined ruderal and segetal communities as epocophytes (83 species) or else settled in natural or seminatural communities as agriophytes (13 species).

Modernisation of the synanthropic flora of Polish sea harbours, according to the general tendencies, will probably proceed towards an increase in the proportion of anthropophytes, with a simultaneous diminution of the archaeophyte number. Among the latter some are already at present extremely rare. On the other hand, the number of the remaining groups of anthropophytes continues to increase. This concerns both established species (epocophytes) and crop plants turning wild (ergasiophytes) as well as transiently introduced plants from distant areas (ephemerophytes). Thus, it is with reason that sea harbours have long been considered as the main entrance gates for many adventitious species (Czubiński 1950, Schwarz 1967).

7. RESULTS AND CONCLUSIONS

On the basis of the investigations carried out in the period 1971-1979 and literature data an inventory of the synanthropic flora of ten Polish sea ports was established (Fig. 1). It numbers 1055 species belonging to 87 families and 457 genera. This flora is exceptionally rich constituting 67.0 per cent of the entire flora of Western Pomerania which includes 1575 species (Czubiński 1950).

The number of species in the particular harbours varies rather widely within the limits of 254 in Władysławowo to 856 in Gdańsk (Table 7). The abundance of synanthropic flora in the particular harbours is the result in the first place of the introduction of diaspores and of favourable habitat conditions, and in some smaller degree of the age of the harbour and its size.

The role of harbours in introduction of plants is indicated by the large number of new species found in the period 1971-1979 in the Gdańsk (273) and Gdynia (243) harbours including 94 species new to Gdańsk and its environs as compared with the data of Schwarz (1967). Among them particularly noteworthy are the new epocophytes: *Amaranthus blitoides*, *A. chlorostachys*, *Chenopodium botrys*, *Atriplex oblongifolium*, *Sisymbrium wolgensse*, *Oenothera syrticola* and *O. hoelscheri* and new ephemerophytes: *Chenopodium pumilio*, *Hibiscus trionum*, *Abutilon theophrasti*, *Oenothera octolineata*, *Trifolium resupinatum*, *Solanum rostratum*, *S. sisymbriifolium*, *Ambrosia trifida*, *Sicyos angulata*, *Setaria faberi* and *Cynodon dactylon*.

The relatively high number of species in the harbours of Gdynia (461), Świnoujście (457) and Kołobrzeg (441) are due to the extensive trade contacts and intensive trans-shipments in these ports, whereas the corresponding number in Władysławowo (254), Łeba (323) and Hel (331) may be attributed to the fact that these are fishery harbours. Moreover, in Władysławowo and Hel wide surfaces have been covered with concrete slabs, asphalt and concrete which make plant development difficult.

There are 198 species common to all the ten harbours, and those found in only one harbour amount to 295 (Table 6). The large number of rare and very rare species is mainly due to the considerable role of ports in their introduction and to the frequent destruction in them of the vegetal cover.

Owing to the intensive economic activity, the number of specimens and species in the particular harbours varied rather widely in the period of the investigations. From among the total number about 30 per cent appeared only transiently. They were usually found only once, or if more frequently, mostly in different years in other parts of the harbours or even in different harbours.

The 1055 species of synanthropic flora in the studied harbours include 554 apophytes (52.5%), 501 anthropophytes (47.5%). The percentual relations of both the groups of plants vary in the particular harbours (Table 7), apophytes prevailing in smaller harbours.

Forest and brushwood species are in majority among apophytes (33.8%), next come the meadow ones (2.8%), the riverside and aquatic ones (15.9%), and those of xerothermic swards (13.4%). These proportions are similar in the particular harbours (Table 10), although in some cases there are deviations caused by different conditions such as for instance dominance of meadow species in the Kołobrzeg and Władysławowo harbours, on account of the absence of forest plantations on their territory and the high proportion of sward and sand species in the Ustka harbour and a low contribution of riverside species in the Ustka (7.1%), Władysławowo (8.1%) and Gdynia (9.5%) harbours. In the latter case the decisive factor is the presence of exclusively concrete quays in these harbours.

In the anthropophyte group ergasiophytes (29.1%) and archaeophytes (27.3%) prevail. An important role is also played by ephemerophytes (24.4%) and epocophytes (16.6%), and a lesser role by hemiagriophytes (2.8%) and holoagriophytes (0.8%).

The studied sea harbours are an area where ephemerophytes are introduced in large numbers (122 species). Most numerous species of this group were noted in large trade harbours, that is Gdańsk (97), Szczecin (21) and Gdynia (21). No representatives of ephemerophytes were discovered in the harbours of Władysławowo, Łeba and Kołobrzeg. In the end period of the research

(1971-1979) the following ephemerophytes were recorded: *Atriplex roseum*, *Coronopus didymus*, *Sisymbrium orientale*, *Hirschfeldia incana*, *Soja hispida*, *Oenothera octolineata*, *Hibiscus trionum*, *Abutilon theophrasti*, *Solanum rostratum*, *S. sisymbriifolium*, *S. alatum*, *Ambrosia trifida*, *A. artemisiifolia*, *Artemisia annua*, *Carthamus tinctorius*, *Centaurea solstitialis*, *Bromus japonicus*, *B. squarrosus*, *Setaria faberi*, *Cynodon dactylon* and *Phalaris canariensis*. Among anthropophytes most numerously represented are south-eastern European and west Asiatic species (32.3%) and south European ones (30.5%). There is also a considerable contribution of American (14.4%) and east-Asiatic species (12.4%). Other regions of the world are represented in smaller numbers (Table 12). Noteworthy is the presence of the Australian species, *Chenopodium pumilio* in the Gdańsk harbour. So far no Australian species had been recorded in the synanthropic flora of Poland (Kornaś 1972).

Perennial species prevail slightly (52.1%) in the examined harbours. Among the annual species anthropophytes dominate (68.5%) and among perennial apophytes (71.8%). The percentual relations as regards perennial and annual species of synanthropic flora are similar in the particular harbours (Table 9).

It results from the analysis of the flora of the studied habitats (Table 14) that most numerous species occur on waste land (686), harbour railway premises (664) and storage yards (582). On lawns and flower beds 372 species were recorded, on the banks of harbour basins 289, on transformed dunes 166 and on breakwaters 84 species.

The synanthropic flora of harbours as compared with other anthropogenic floras (Tables 15 and 17) exhibits the highest number of species (1055) a high contribution of anthropophytes (501) and high indexes: of synanthropisation ($S = 233$), of lability ($I = 268$) and of modernisation ($M = 0.70$). This rich flora of sea harbours is due to:

- a) the introduction of diaspores of new species during trans-shipment of goods from various continents;
- b) import of seeds and fruits of numerous species and with them of seeds of many weeds;
- c) the system of storage of mass goods and some kinds of general cargo on open spaces;
- d) considerable differences in the harbour habitats as regards moisture, edaphic conditions etc.;
- e) the old age of the harbour;
- f) the length of the period of floristic investigations, mainly in the case of Gdańsk;
- g) the situation of the harbours within cities rich in synanthropic flora (Szczecin, Gdańsk, Gdynia, Świnoujście, Kołobrzeg).

Sea harbours were and continue to be the main entrance gates for

many adventitious species. In the latest research (1971-1979) many species new for the synanthropic flora of the towns with harbours, for the flora of Pomerania as well as three species new for the synanthropic flora of Poland have been found: the latter are *Chenopodium pumilio*, *Solanum sisymbriifolium* and *Setaria faberi*.

Doc dr Józef Misiewicz
Department of Botany
Higher Pedagogical School
ul. Arciszewskiego 22b
76-200 Słupsk, Poland

8. STRESZCZENIE

Na podstawie badań przeprowadzonych w latach 1971-1979 oraz danych z literatury, ustalono inwentarz flory synantropijnej 10 polskich portów morskich (ryc. 1). Liczy on 1055 gatunków, należących do 87 rodzin i 457 rodzajów. Jest to wyjątkowo bogata flora, gdyż stanowi ona 67.0% całej flory Pomorza Zachodniego, liczącej 1575 gatunków (Czubiński 1950).

Liczby gatunków w poszczególnych portach są dość zróżnicowane i wahają się w granicach od 254 we Władysławowie do 856 w Gdańsku (tab. 7). Bogactwo flory synantropijnej poszczególnych portów jest wynikiem przede wszystkim zawlekania diaspory oraz występowania odpowiednich warunków siedliskowych, natomiast w mniejszym stopniu zależy od wieku portu i jego wielkości.

O roli portów w zawlekanii roślin mogą świadczyć duże liczby nowych gatunków stwierdzonych w latach 1971-1979 w portach Gdańsk (273) i Gdynia (243), w tym 94 nowe gatunki dla Gdańska i okolicy w porównaniu z opracowaniem Schwarza (1967). Na szczególną uwagę zasługują tu nowe epekofity: *Amaranthus blitoides*, *A. chlorostachys*, *Chenopodium botrys*, *Atriplex oblongifolium*, *Sisymbrium wolgensse*, *Oenothera syrticola* i *O. hoelscheri*; oraz nowe efemerofity: *Chenopodium pumilio*, *Hibiscus trionum*, *Abutilon theophrasti*, *Oenothera octolineata*, *Trifolium resupinatum*, *Solanum rostratum*, *S. sisymbriifolium*, *Ambrosia trifida*, *Sicyos angulata*, *Setaria faberi* i *Cynodon dactylon*.

Stosunkowo duże liczby gatunków w portach Gdynia (461), Świnoujście (457) i Kołobrzeg (441) są wynikiem rozległych kontaktów handlowych i intensywnych przeładunków. Natomiast najmniejsze liczby gatunków w portach Władysławowo (254), Łeba (323) i Hel (331) należy tłumaczyć tym, że są to porty rybackie. Ponadto w portach Władysławowa i Helu znaczne powierzchnie zostały pokryte płytami chodnikowymi, asfaltem i betonem, co utrudnia rozwój roślin.

Gatunków wspólnych dla wszystkich 10 badanych portów jest 198, a znalezionych tylko w jednym z badanych portów 295 (tab. 6). Duża liczba gatunków rzadkich i bardzo rzadkich jest głównie wynikiem dużego udziału portów w ich zawlekanii oraz częstego niszczenia pokroju roślinnej.

Na skutek intensywnej działalności gospodarczej w poszczególnych portach, ilość okazów i gatunków ulegała dość znacznym wahaniom w okresie prowadzonych badań. Z ogólnej liczby gatunków około 30% pojawiało się tylko przejściowo. Spotykano je zwykle jeden raz, a jeśli kilkakrotnie, to przeważnie w różnych latach, w innych częściach portów lub nawet w innych portach.

Spośród 1055 gatunków flory synantropijnej badanych portów, apofity stanowią 554 gatunki (52,5%), a antropofity – 501 gatunków (47,5%). Stosunki procentowe obu grup roślin w poszczególnych portach kształtują się dość różnie (tab. 7), przy czym w małych portach przeważają apofity.

Wśród apofitów najwięcej jest gatunków leśnych i zaroślowych (33,8%), łąkowych (20,8%), nadwodnych i wodnych (15,9%) oraz muraw kserotermicznych (13,4%). Stosunki te w poszczególnych portach kształtują się podobnie (tab. 10), chociaż w kilku przypadkach występują różnice uzasadnione odmiennymi warunkami, jak np. dominacja gatunków łąkowych w porcie

Kołobrzeg i Władysławowo z powodu braku zagajników na ich terenach, znaczny udział gatunków murawowych i piaszczysk w porcie Ustka, mały udział gatunków nadwodnych w portach Ustka (7.1%), Władysławowo (8.1%) i Gdynia (9.5%). W ostatnim przypadku czynnikiem decydującym jest występowanie na terenach wymienionych portów, wyłącznie nabrzeży betonowych.

W grupie antropofitów przeważała ergazjofity (29.1%) i archeofity (27.3%). Istotną rolę odgrywają także efemerofity (24.4%) i epekofity (16.6%), znacznie mniejszą natomiast — hemiagriofity (1.8%) i holoagriofity (0.8%).

Badane porty morskie są miejscem liczego zawlekania efemerofitów (122 gatunki). Najwięcej gatunków z tej grupy zanotowano w dużych portach handlowych tj. w Gdańsku (97), Szczecinie (21) i Gdyni (21). W portach Władysławowa, Łeby i Kołobrzegu nie spotkano przedstawicieli efemerofitów. W ostatnim okresie badań (1971-1979) stwierdzono między innymi następujące efemerofity: *Atriplex roseum*, *Coronopus didymus*, *Sisymbrium orientale*, *Hirschfeldia incana*, *Soja hispida*, *Oenothera octolineata*, *Hibiscus trionum*, *Abutilon theophrasti*, *Solanum rostratum*, *S. sisymbriifolium*, *S. alatum*, *Ambrosia trifida*, *A. artemisiifolia*, *Artemisia annua*, *Carthamus tinctorius*, *Centaurea solstitialis*, *Bromus japonicus*, *B. squarrosus*, *Setaria faberi*, *Cynodon dactylon* i *Phalaris canariensis*.

Wśród antropofitów najliczniej są reprezentowane gatunki z południowo-wschodniej Europy i zachodniej Azji (32.3%) oraz gatunki z południowej Europy (30.5%). Znaczny udział mają także gatunki amerykańskie (14.4%) i wschodnioazjatyckie (12.4%). Pozostałe regiony świata są reprezentowane mniej licznie (tab. 12), jednak zasługuje na uwagę obecność gatunku australijskiego (*Chenopodium pumilio*) w porcie gdańskim. Należy dodać, że dotychczas nie notowano gatunków australijskich we florze synantropijnej Polski (Kornaś 1972).

W badanych portach nieznacznie przeważają gatunki trwałe (52.1%). Wśród gatunków krótkotrwałych dominują antropofity (68.5%), a wśród gatunków trwałych apofity (71.8%). Podobnie kształtują się stosunki procentowe trwałości gatunków flory synantropijnej w poszczególnych portach (tab. 9).

Z analizy flory wyróżnionych siedlisk (tab. 14) wynika, że najczęściej gatunków występuje na nieużytkach (686), portowych terenach kolejowych (664) i placach składowych (582). Na trawnikach i kwietnikach zanotowano 372 gatunki, na brzegach basenów portowych — 289 gatunków, na przekształconych wydmach — 166 gatunków i na falochronach — 84 gatunki.

Flora synantropijna portów, w porównaniu z innymi florami antropogenicznymi (tab. 15 i 17), wyróżnia się największą liczbą gatunków (1055), znacznym udziałem antropofitów (501) oraz wysokimi wskaźnikami: stopnia synantropizacji ($S = 233$), stopnia labilności ($I = 268$) i stopnia modernizacji ($M = 0.70$). To duże bogactwo portów jest wynikiem:

- licznego zawlekania diaspory nowych gatunków podczas przeładunku towarów, pochodzących z różnych kontynentów;
- sprowadzania do portów nasion i owoców wielu gatunków, a wraz z nimi nasion różnych chwastów;
- sposobu magazynowania towarów masowych i drobnicy na otwartych placach;
- znacznego zróżnicowania siedlisk portowych pod względem wilgotności, warunków edaficznych, itp.;
- długiego czasu istnienia portów;
- długiego okresu badań florystycznych w portach, głównie w porcie gdańskim;
- położenia portów w obrębie bogatych we florę synantropijną miast (Szczecin, Gdańsk, Gdynia, Świnoujście, Kołobrzeg).

Porty morskie były i są nadal głównymi bramami wpadowymi dla bardzo wielu gatunków adwentywnych. W ostatnim okresie badań (1971-1979) stwierdzono liczne nowe gatunki we florze synantropijnej miast, w obrębie których są usytuowane porty, we florze Pomorza, a także trzy nowe gatunki we florze synantropijnej Polski: *Chenopodium pumilio*, *Solanum sisymbriifolium* i *Setaria faberi*.

9. REFERENCES

- Abromeit J., 1898-1940. Flora von Ost- und Westpreussen. Vol. I. 1 (1898), 2(1903), II(1926), III(1931), IV(1934), Schlussband (1940). Friedländer u. Sohn, Berlin-Königsberg.
- Andruszkiewicz W., 1960. Porty handlowe. Wydawnictwo Morskie, Gdynia.
- Andruszkiewicz W., Kleban Cz., Obertyński E., 1961. Port Gdynia. Zarząd Portu Gdynia, pp. 1-182.
- Anioł-Kwiatkowska J., 1974. Flora i zbiorowiska synantropijne Legnicy, Lubina i Polkowic. Acta Univ. Wratislaviensis, Prace Botaniczne 19: 1-152.
- Ascherson P., Graebner P., 1898-99. Flora des Nordostdeutschen Flachlandes. Geb. Borntraeger., Berlin, pp. 1-875.
- Bail Th., 1880. Skizze der Flora Danzings und seiner Umgegend. In: H. Conwentz, „Danzig in naturwissenschaftlicher und medicinischer Beziehung“, Kefemann, Danzig, pp. 40-60.
- Balcerkiewicz S., 1971. Zapiski florystyczne z Pomorza Szczecińskiego. Bad. Fizjogr. nad Pol. zach. Ser. B, 24: 253-255.
- Blom C., 1927, 1933, 1936, 1939, 1961. Bidrak till kännedom om Sveriges adventivflora. I. Acta Horti Gothob. 3: 383-403; II. Ibid. 8: 167-188; III. Ibid. 11: 159-182; IV. Ibid. 13: 383-403; V. Ibid. 24: 61-133.
- Blomquist S. G., 1917. Ballastvegetationen vid Kalmar 1912-1914. (Ref.: Die Ballastvegetation bei Kalmar 1912-1914). S. Bot. Tidskr. 11: 289-303.
- Brocki Z., 1959. Port Gdański. Wydawnictwo Morskie, Gdynia.
- Celiński F., 1964. Rośliny naczyniowe Puszczy Bukowej pod Szczecinem. PTPN, Pr. Kom. Biol. 29(2): 135-322.
- Czubiński Z., 1950. Zagadnienia geobotaniczne Pomorza. PTPN, Wydz. Mat.-Przyr. Kom. Fizjogr. Bad. nad Pol. zach. 2(4): 437-660.
- Ćwikliński E., 1965. Rzadsze rośliny Szczecina. Fragm. Flor. Geobot. 11(1): 3-5.
- Ćwikliński E., 1967. Nowy dla Polski gatunek wiesiołka *Oenothera ammodila* Focke. Fragm. Flor. Geobot. 13(1): 51-53.
- Ćwikliński E., 1968. Neolity terenów kolejowych województwa szczecińskiego. Mat. Zakł. Fitosoc. Stos. UW 25: 125-138.
- Ćwikliński E., 1969. Notatki florystyczne z województwa szczecińskiego. Bad. Fizjogr. nad Pol. zach. Ser. B, 23: 261-264.
- Ćwikliński E., 1970. Flora synantropijna Szczecina. Monogr. Bot. 33: 1-103.
- Ćwikliński E., 1971a. Notatki florystyczne z województwa szczecińskiego. Cz. II. Bad. Fizjogr. nad Pol. zach. 24: 247-251.
- Ćwikliński E., 1971b. Flora synantropijna Zielonej Góry i Koszalina na tle warunków przyrodniczych i rozwoju miasta. Mat. Zakł. Fitosoc. Stos. UW 27: 81-113.
- Ćwikliński E., 1974. Flora i zbiorowiska roślinne terenów kolejowych województwa szczecińskiego. AR w Szczecinie, rozprawy. 40: 1-149.
- Dostál J., 1950. Květena ČSR. Přírodovědecké nakladatelství, Praha, pp. 2269.

- Dziedziul B. (ed.), 1975. Port Szczeciński. Dzieje i rozwój do roku 1970. PWN, Warszawa-Poznań.
- Grodek A., 1948. Handel odrzański w rozwoju historycznym. In: Monografia Odry, Inst. Zach., Poznań, pp. 384-418.
- Hansen A., 1959. Gatunki i mieszańce rodzaju *Elytrygia* na polskim wybrzeżu Bałtyku. *Fragm. Flor. Geobot.* 5(2): 181-189.
- Hegi G., 1908-1960. *Illustrierte Flora von Mittel-Europa*, vol. I-VII. Lehmann-Verlag. München.
- Hejny S., Jehlik V., Kopecký K., Kropáč Z., Lhotska M., 1973. Karanténni plevelé Československa. Studie ČSAV, part 8. Praha.
- Heliński Z., 1962. Przeladunki w porcie szczecińskim w latach 1946-1960. *Studium sezonowości*. Szczecin.
- Helm O., 1881. Ballastpflanzen auf der Westerplatte. *Bericht. d. Westpr. Bot.-Zool. Vereins* 4: 23-24.
- Herweg O., 1914. Flora der Kreise Neustadt und Putzig in Westpreussen. *Bericht d. Westpr. Bot.-Zool. Vereins* 37: 85-331.
- Hilbert H., 1910. *Schrift d. Phys. Ökonom. Gesell.* 51: 87-91.
- Hoffa M., 1968. Niektóre cechy fizyczno-geograficzne rejonu Kołobrzegu i ich znaczenie dla portu i uzdrowiska. *PTPN, Wydz. Mat.-Przyr., Pr. Kom. Biol.* 9(3): 1-116.
- Holzfuß E., 1916. Die Gattung *Potentilla* in Pommern. *Allgem. Bot. Zeitschrif. Syst., Flor., Pflanzengeogr. Jg. XXII*, nr 1-4.
- Holzfuß E., 1920. Beitrag zur Flora der Insel Wollin. *Abhandl. Ber. Pom. Naturforsch. Ges. Jg. I. Stettin*.
- Holzfuß E., 1933. Nachtrag zur Flora von Stolp und Stolpmünde. *Dohrniana* 13: 95.
- Holzfuß E., 1936. Die Pflanzenwelt der Schuttplätze in Pommern. *Dohrniana* 15: 116-126.
- Holzfuß E., 1937. Beitrag zur Adventivflora von Pommern. *Dohrniana* 16: 94-130.
- Holzfuß E., 1941. Mitteilungen aus der Flora der Schuttplätze in Pommern. *Dohrniana* 20: 185-187.
- Hupke H., 1933. Adventiv- und Ruderalpflanzen der Kölner Güterbahnhöfe, Hafenanlagen und Schuttplätze. *Wissenschaftlichen Mitteilungen des Vereins für Natur- und Heimatkunde in Köln a Rh.* 3(1): 3-21.
- Hupke H., 1935. Adventiv- und Ruderalpflanzen der Kölner Güterbahnhöfe Hafenanlagen und Schuttplätze. I Nachtrag. *Verb Natur. Rheinlande Westfalens*. Bd. 91: 187-204.
- Kalkreuth P., 1941. Zur Analyse der heimischen Pflanzendecke. *Bericht Westpr. Bot.-Zool. Vereins* 63: 14-22.
- Klinggraeff C., 1848. *Flora von Preussen*. Baumann (ed.), Marienwerder.
- Klinggraeff C., 1854. Nachtrag zur Flora von Preussen. Lovysohn (ed.), Marienwerder.
- Klinggraeff C., 1866. Die Vegetationsverhältnisse der Provinz Preussen. Lovysohn (ed.), Marienwerder.
- Klinggraeff H., 1880. Versuch eine topographischen der Flora Provinz Westpreussen. *Schrift. Naturf. Gesell. Danzig* 5(2): 82-232.
- Klinsmann E., 1836. Beiträge zur Flora von Danzig nebst einigen andern Preussischen Gegenden. *Preuss. Prov.-Bl.* 15: 467-477.
- Klinsmann E., 1843. Novita atque defectus florae Gedanensis. (Neueste Schr. Naturf. Ges. Dzg.). *Schrift. Naturf. Gesell. Danzig, N.F.* 4(1): 103-120.
- Klinsmann E., 1865. Ergänzungen und Berichtigungen zu Novita atque defectus florae Gedanensis. *Schrift. Naturf. Gesell. Danzig, N.F.* 1(2): 1-8.
- Kornaś J., 1968a. Geograficzno-historyczna klasyfikacja roślin synantropijnych. *Mat. Zakł. Fitosoc. Stos. UW* 25: 33-42.
- Kornaś J., 1968b. Prowizoryczna lista nowszych przybyszów synantropijnych (kenofitów) zadomowionych w Polsce. *Mat. Zakł. Fitosoc. Stos. UW* 25: 43-53.

- Kornaś J., 1972. Wpływ człowieka i jego gospodarki na szatę roślinną Polski. In: Szata roślinna Polski. Eds.: W. Szafer and K. Zarzycki, vol. I, pp. 95-128. PWN, Warszawa.
- Kornaś J., 1977. Analiza flor synantropijnych. Wiad. Bot. 21(2): 85-91.
- Kornaś J., Leśniewska J., Skrzywanek A., 1959. Obserwacje nad florą linii kolejowych i dworców towarowych w Krakowie. Fragm. Flor. Geobot. 5(2): 199-216.
- Koselnik B., 1965. Porty morskie Wybrzeża Środkowego. In: Województwo koszalińskie. Monografia geograficzno-gospodarcza. Eds.: F. Barciński, B. Krygowski and S. Zajchowska. Instytut Zachodni, Poznań, pp. 471-498.
- Kowalenko W., 1950. Starosławiańskie grody portowe na Bałtyku. Prz. Zach. 6: 378-419.
- Kownas S., 1958. Roślinność Ziemi Szczecińskiej. Szczecin (Mies. Pom. Zach.) nr 6.
- Kownas S., 1959. Drzewa i krzewy egzotyczne w Szczecinie. Szczecin (Mies. Pom. Zach.) nr 3: 61-66.
- Kownas S., Piskorski Cz., 1958. Szczecin — miasto parków i zieleni. PWN, Poznań.
- Kownas S., Sienicka A., 1962. Obecny stan zadrzewienia miasta Szczecina. Szczecin. Tow. Nauk. Wydz. Nauk Przyr.-Roln. Vol. 11, fasc. 1.
- Krauze H., 1934. Weitere Beiträge zur Flora von Stolp und Umgegend. Dohrniana 13: 217-218.
- Leick E., 1926. Die Pflanzendecke der Provinz Pommern. Sonderabdruck aus dem Pom. Heimatbuch. Berlin, pp. 95-210.
- Lehmann E., 1895. Flora von Polnisch Livland. Mittiesen and Jurjew (ed.). Dorpad.
- Lejmbach B., Rurka Z., Siedlecka B., Sijka J., 1975. Flora torowisk kolejowych Pobrzeża Wschodniopomorskiego. Fragm. Flor. Geobot. 19(1): 53-66.
- Lesiński H. (ed.), 1965. Dzieje Kołobrzegu (X-XX wiek). Biblioteka Słupska, vol. 14, Wyd. Poznańskie, Poznań.
- Lhotská M., Hejný S., 1979. *Chenopodium pumilio* in Czechoslovakia: its strategy of dispersal and domestication. Folia Geobot. Phytotax. 14: 367-375.
- „Locja Bałtyku. Część południowa. Wybrzeże polskie”, 1973. Vth edition. Biuro Hydrograficzne Marynarki Wojennej, Gdynia.
- Lützw C., 1892. Seltene und zweifelhafte Pflanzen in Westpreussen. Bericht Westpr. Bot.-Zool. Vereins 15: 16-20.
- Magiera W., 1966. Zespół Szczecin-Świnoujście portem rzeczno-morskim ujścia Odry. Technika i Gospodarka, nr 15.
- Marsson Th. Fr., 1869. Flora von Neu-Vorpommern und den Insel Rügen und Usedom. Leipzig.
- Michalak S., 1970. Florą synantropijna miasta Opola. Rocznik Muzealny 4: 1-181.
- Mieczyski T., 1946. Gleby byłego terytorium Gdańska. Inst. Bałt., Gdańsk-Bydgoszcz-Szczecin.
- Mikołajski J., 1960. W sprawie warunków rozwoju polskich portów morskich. Szczecin (Mies. Pom. Zach.) nr 4-5: 126.
- Misiewicz J., 1970. Masowe występowanie *Anthoxanthum aristatum* Boiss. w zasiewach żyta ozimego (*Secale cereale* L.) Fragm. Flor. Geobot. 16(2): 317-318.
- Misiewicz J., 1971. Flora i zbiorowiska synantropijne Gorzowa Wlkp. i okolicy. Mat. Zakł. Fitosoc. Stos. UW 27: 65-79.
- Misiewicz J., 1976. Flora synantropijna i zbiorowiska ruderalne polskich portów morskich. WSP, Słupsk, pp. 1-321.
- Misiewicz J., 1977a. Interesujące gatunki we florze synantropijnej terenów portowych Wybrzeża Środkowego. In: Ochrona i Kształtowanie Środowiska Przyrodniczego. WSP, Słupsk, pp. 227-233.
- Misiewicz J., 1977b. Wpływ portów Pomorza Środkowego na zawlekanie obcych gatunków roślin. Koszalińskie Studia i Materiały 3(19): 92-98.

- Misiewicz J., 1977c. Nieznane zbiory zielnikowe flory Pomorza zachowane w Muzeum w Darłowie. In: Ochrona i Kształtowanie Środowiska Przyrodniczego Pomorza Środkowego. WSP, Słupsk, pp. 195-226.
- Misiewicz J., 1978a. Flora synantropijna Słupska na tle warunków przyrodniczych i rozwoju miasta. WSP, Słupsk, pp. 1-143.
- Misiewicz J., 1978b. Rola polskich portów morskich w zawlekanii obcych gatunków roślin. *Studia i Materiały Oceanologiczne* 27: 97-108.
- Misiewicz J., Rudkowska J., Szłapak M., 1979. Badania nad florą synantropijną Świnoujścia. *Fragm. Flor. Geobot.* (in press).
- Moniak J., 1961. Środowisko geograficzne województwa gdańskiego. *Zesz. Nauk. WSE*, seria B, 9.
- Müller W., 1904. Flora von Pommern. Stettin.
- Nowiński M., 1962. Spostrzeżenia florystyczne z okolic Łeby. *Przyr. Pol. zach.* 6(1-4): 55-59.
- Nowiński M., 1965. Chwasty segetalne Łeby i okolicy w województwie gdańskim. *PTPN*. Wyd. Nauk Roln. i Leśn., Pr. Kom. Roln. i Kom. Leśn. 19(2): 303-352.
- Piotrowska H., 1957. Z badań nad roślinnością halofilną wysp Wolina i Uznamu. *Przyr. Pol. zach.* 1/2.
- Piotrowska H., 1966a. Rośliny naczyniowe wysp Wolina i południowo-wschodniego Uznamu. *PTPN*, Wyd. Mat.-Przyr., Prace Kom. Biol. 30(4): 1-283.
- Piotrowska H., 1966b. Stosunki geobotaniczne wysp Wolina i południowo-wschodniego Uznamu. *Monogr. Bot.* 22.
- „Port Gdański”, 1929. Opracowane przez Radę Portu i Dróg Wodnych w Gdańsku. Przemysł i Handel, Warszawa.
- „Port Gdynia”, 1936. Instytut. Wyd. Państw. Szkoły Morskiej, Gdynia, pp. 1-98.
- Prawdź K., 1961. Klimat województwa szczecińskiego w świetle potrzeb rolnictwa. *Wiad. Zach.*, Szczecin.
- Prawdź K., 1962. Zarys klimatu rolniczego województwa koszalińskiego. *Soc. Scien. Stetinensis* 12(1).
- Preuss H., 1910. Die Vegetationsverhältnisse der westpreussischen Ostseeküste. Danzig.
- Preuss H., 1928. Das Herbar Klinsmann bes Berücks der Danziger Adventivflora. *Bericht Westpr. Bot.-Zool. Vereins* 50: 201-230.
- Probst R., 1949. Wolladventivflora Mitteleuropas. Vogt-Schild AG, Solothurn.
- Radomski J., Jasnowska J., 1964. Rośliny zbiorowisk otwartych na zachodniej krawędzi doliny Odry. Cz. I. *Zesz. Nauk. WSR w Szczecinie*, 17.
- Radomski J., Jasnowska J., 1965. Roślinność zbiorowisk murawowych na zachodniej krawędzi doliny Odry. Cz. II. *Zesz. Nauk. WSR w Szczecinie*, 19.
- Rocznik statystyczny 1978. GUS, Warszawa.
- Romer E., 1949. Regiony klimatyczne Polski. Pr. Wrocł. Tow. Nauk., ser. B, 16.
- Rostański K., Szotkowski P., 1973. Gatunki rodzaju *Oenothera* L. w portach rzecznych górnej Odry. *Frag. Flor. Geobot.* 19(4): 373-378.
- Runge F., 1956. Adventivpflanzen der beiden Kanalhäfen in Münster während der Jahre 1950 bis 1956. *Natur und Heimat* 16: 74-79.
- Runge F., 1965. Adventivpflanzen der beiden Kanalhäfen während der Jahre 1957 bis 1964. *Natur und Heimat* 25: 61-64.
- Sarrisalo-Taubert A., 1963. Die Flora ihrer beziehung zur siedlung und siedlungsgeschichte in den südfinnischen Staden Porvoo, Lovisa und Hamina. *Ann. Bot. Soc. „Vanamo”* 35(1): 1-190.
- Scheuermann R., 1956. Beitrag zur Adventivflora in Pommern. *Decheniana* 108.
- Schmidt W. L. E., 1840. Flora von Pommern und Rügen. Stettin.
- Schoeps H., 1931. Beitrage zur Flora von Stolpmünde. *Dohrniana* 11(1): 148-150.

- Schwarz Z., 1961. Nowe i bardziej interesujące gatunki we florze synantropijnej Pomorza Gdańskiego. *Fragm. Flor. Geobot.* 7(1): 59-75.
- Schwarz Z., 1967. Badania nad florą synantropijną Gdańska i okolicy. *Acta Biol. Med. Soc. Sc. Gedan.* 11: 363-494.
- Schwarz Z., 1971. Flora synantropijna miasta Elbląga. *Mat. Zakł. Fitosoc. Stos. UW* 27: 145-166.
- Sendek A., 1971. Flora synantropijna miasta Kluczborka w związku z jego funkcją. *Mat. Zakł. Fitosoc. Stos. UW* 27: 217-229.
- Siuta J., 1973a. Projekt rekultywacji gruntu oraz urządzenie zieleni w Porcie Północnym i w strefie ochronnej. Cz. I. Projekt rekultywacji gruntu. IUNG, Puławy (mscr.).
- Siuta J., 1973b. Port Północny. Studium kształtowania środowiska przyrodniczego. Cz. II. IUNG, Puławy, (mscr.).
- Siuta J., 1974. Jak zazielenić Port Północny w Gdańsku. *Przyroda polska* 6: 5.
- Sowa R., 1971. Flora i roślinne zbiorowiska ruderalne na obszarze województwa łódzkiego ze szczególnym uwzględnieniem miast i miasteczek. *UŁ. Łódź*, pp. 1-282.
- Sowa R., Olaczek R., 1978. Stan badań szaty roślinnej miast Polski. *Wiad. Ekol.* 24(1): 25-42.
- Stricker W., 1960. Floristische Beobachtungen in der Gegend von Leipzig. *Berichte der Arbeitsgemeinschaft sächsischer Botaniker* 2: 168-186.
- Stricker W., 1961. Floristische Beobachtungen in der Gegend von Leipzig. *Ibidem*, nr 3: 178-188.
- Sulma T., 1959. Główne problemy botaniczne Pomorza. *Acta Biol. Med. Soc. Sc. Gedan.* 3: 10-73.
- Szmajda P., 1974. Flora synantropijna Stargardu Szczecińskiego i Pyrzyc. *Bad. Fizjogr. nad Pol. zach.*, ser. B, 27: 227-261.
- Szopowski Z., 1962. Małe porty Pomorza Zachodniego w okresie do drugiej wojny światowej. PWN, Warszawa-Poznań.
- Szotkowski P., 1972. Flora synantropijna portów rzecznych górnej Odry (Koźla, Gliwice, Opola). *UJ, Instytut Botaniki, Kraków*, pp. 299 (Ph. D. Thesis).
- Trzcińska-Tacik H., 1963. Badania nad zasięgami roślin synantropijnych. Cz. II. *Rumex confertus* Willd. w Polsce. *Fragm. Flor. Geobot.* 9(1): 71-84.
- Urbański J., 1958. Sałata tatarska (*Lactuca tatarica* (L.) C. A. Meyer) koło Świnoujścia. *Bad. Fizjogr. nad Pol. zach.* 4.
- Wangerin W., 1916. Beiträge zur Kenntnis der Vegetationsverhältnisse der Moore Westpreussens. *Bericht Westpr. Bot.-Zool. Vereins* 38: 77-136.
- Wangerin W., 1921. Beiträge zur Kenntnis der Verbreitung der Gefäßpflanzen im nordostdeutschen Flachlande. *Bericht Westpr. Bot.-Zool. Vereins* 43: 46-55.
- Wangerin W., 1925. Beiträge zur Kenntnis der Verbreitung der Gefäßpflanzen im nordostdeutschen Flachlande. *Ver. Bot. Prov.* 67: 91-99.
- Wangerin W., 1930-1931. Die pflanzengeographischen Verhältnisse der Umgegend von Danzig. *Schrift. Naturf. Gesell. in Danzig*, N. F. 19: 37-122.
- Wangerin W., 1937. Beiträge zur Kenntnis der Verbreitung der Gefäßpflanzen im nordostdeutschen Flachlande. *Ver. Bot. Prov.* 77: 58-85.
- Wisniewski W., 1953. Atlas opadów atmosferycznych w Polsce 1891-1930. PIHM, Warszawa.
- Zajac E. U., Zajac A., 1975. Lista archeofitów występujących w Polsce. *Zesz. Nauk. UJ, Pr. Bot.* 3: 7-16.
- Zaremba P., Orlińska H., 1965. Urbanistyczny rozwój Szczecina. Wydawnictwo Poznańskie, Poznań.
- Zbierski A., 1962. Port gdański w X-XV wieku. In: *Z przeszłości Pomorza Wschodniego*. PTA, Wrocław-Warszawa-Kraków.

- Żukowski W., 1960. Nowe stanowiska roślin synantropijnych ze szczególnym uwzględnieniem Polski północno-zachodniej. *Fragm. Flor. Geobot.* 6(4): 481-488.
- Żukowski W., 1961. Notatki florystyczne z Pomorza Zachodniego. *Bad. Fizjogr. nad Pol. zach.* 8: 219-230.
- Żukowski W., Piaszyk M., 1971. Rozmieszczenie niektórych gatunków synantropijnych z rodzaju *Artemisia* L. w Polsce. *Bad. Fizjogr. nad Pol. zach.* 24: 107-128.

CONTENTS

1. Abstract	5
2. Introduction	5
3. Outline of physiographic conditions	7
3.1. Territory investigated	7
3.2. Climate	10
3.3. Habitat conditions in harbour areas	13
4. Trade relations of Polish sea harbours	15
5. History of the research on synanthropic flora of Polish sea harbours	17
6. Analysis of synanthropic flora	20
6.1. State of flora	20
6.2. Flora of the particular harbours	22
6.3. Biological durability of species	31
6.4. Analysis of apophytes	33
6.5. Analysis of anthropophytes	36
6.6. Geographical and genetic anthropophyte spectrum	39
6.7. Flora of harbour habitats	41
6.8. State of synanthropic flora in the harbours as compared with that of some selected regions of Poland	52
7. Results and conclusions	55
8. Streszczenie	59
9. References	61